

ANSWERS XI

CHEMISTRY

1.d 2.a 3.b 4.d 5.d 6.c 7.c 8.a 9.c 10.b 11.b 12.c 13.a
14.d 15.b 16.c 17.a 18.b 19.c 20.a 21.d 22.c 23.b 24.d 25.a 26.c
27.d 28.a 29.d 30.c

PHYSICS

1.d 2.b 3.c 4.b 5.c 6.c 7.b 8.d 9.b 10.c 11.c 12.a 13.d
14.b 15.a 16.a 17.a 18.b 19.a 20.a 21.d 22.c 23.c 24.d 25.c 26.d
27.a 28.b 29.a 30.b

MATHEMATICS

1.d 2.a 3.b 4.b 5.d 6.c 7.d 8.b 9.a 10.a 11.b 12.b 13.a
14.c 15.b 16.d 17.c 18.a 19.d 20.c 21.a 22.d 23.b 24.d 25.a 26.d
27.b 28.c 29.d 30d

HINTS AND EXPLANATIONS XI

CHEMISTRY


Sol.1

10% solution means 10g in 100 mL; 1 mol of glucose = 180 g

10 g of glucose is present in 100 mL 180 g of glucose will be present in $\frac{100 \times 180}{10} = 1800$ mL or 1.8 L


Sol.2

Formation of I_3^- involves sp^3d hybridization.




Sol.3

Structure of a phospholipid is shown below



Sol.4

Dipole moment of CCl_4 is zero.



Sol.5

Microwave has lowest frequency, γ -rays has highest frequency (lowest wavelength). Decreasing order of frequency: γ -rays > X-rays > visible > microwave.

Sol.6

In the given reaction oxidation state of Cr changes from $+6 \rightarrow +3$, i.e., 3 electrons per Cr atom. For $\text{Cr}_2\text{O}_7^{2-}$, the oxidation state changes by 6, therefore its equivalent wt = $M/6$

Sol.7

The formula of potassium dicyanobis(oxalate) nickelate (11) is $\text{K}_4[\text{Ni}(\text{CN})_2(\text{Ox})_2]$

Sol.8

$$K_{sp} = [\text{Ba}^{2+}][\text{CrO}_4^{2-}]$$

$$2.4 \times 10^{-10} = [\text{Ba}^{2+}] \times 6 \times 10^{-4}$$

$$[\text{Ba}^{2+}] = 0.4 \times 10^{-6} = 4 \times 10^{-7}$$

Sol.9

$$Q = I \times t = 1 \times 60 = 60 \text{ C}$$

96500 c delivers 1 mol or 6.023×10^{23} electrons at cathode

$$60 \text{ C will deliver electrons} = \frac{6.023 \times 10^{20}}{96500} \times 60 = 3.74 \times 10^{20}$$

Sol.10


The given reaction $\text{Cu}_2\text{O} + \text{FeS} \rightarrow \text{FeO} + \text{Cu}_2\text{s}$ takes place during smelting

Sol.11

pH of a buffer solution is given as


$$pH = pKa + \log \frac{[\text{salt}]}{[\text{acid}]}$$

$$= 4.75 + \log \frac{0.1}{0.1} = 4.75$$

Sol.12

Sol.13

Addition of HBr on acetylene gives ethylidene bromide


**Sol.14**

Number of electrons in the given species are :

$$\text{Cl}^- = 2,8,8; \text{F}^- = 2,8; \text{Na}^+ = 2,8; \text{Mg}^{2+} = 2,8$$

Sol.15

Electrolysis of sodium salt of succinic acid gives ethylene.

**Sol.16**


All the elements of lanthanide and actinide series are not radioactive

Sol.17

Natural gas is a mixture of gaseous paraffins

Sol.18


Action of heat on mixture of anhydrous sodium propanoate and soda lime produces ethane.

**Sol.19**

The presence of electron withdrawing atoms or groups at the α -carbon of a carboxylic acid increases its acidity. Fluorine is more electronegative than both Br and Cl.


Sol.20

From the options given the rotation about C=C in 1,1,2,2-tetrabromoethylene is most sterically hindered.



Sol.21

Pent-2-ene on ozonolysis gives $\text{CH}_3\text{CH}_2\text{CHO}$ and CH_3CHO .

**Sol.22**

n-hexane can be prepared by Wurtz reaction of *n*-propyl bromide $\text{CH}_3\text{CH}_2\text{CH}_2\text{Br} + 2\text{Na} + \text{BrCH}_2\text{CH}_2\text{CH}_3 \rightarrow \text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$

Sol.23

Fog is a colloidal system of liquid in a gas

Sol.24

$$\frac{V_1}{T_1} = \frac{V_2}{T_2} \text{ or } V_2 = \frac{V_1}{T_1} T_2 = \frac{V \times 500}{300} = \frac{5V}{3};$$


$$\text{If } V = 100 \text{ mL then } V_2 = \frac{500}{300} = 166.6 \text{ mL}$$

$$\text{Air escaped during heating} = 66.6 \text{ mL or } \frac{66.6 \times 100}{166.6} = 40\%$$

Sol.25

$$8.8 \text{ g CO}_2 = 0.2 \text{ mol}; \quad 5.4 \text{ g H}_2\text{O} = 0.3 \text{ mol.}$$

Number of moles of oxygen required to produce 0.2 mol of CO_2 and 0.3 moles of H_2O can be calculated from the balanced equation.



$$3\text{g} \quad \quad \quad 0.35 \text{ mol} \quad \quad \quad 0.2 \text{ mol} \quad \quad \quad 0.3 \text{ mol}$$

$$3\text{g} \quad \quad \quad 11.2 \text{ g} \quad \quad \quad 8.8 \text{ g} \quad \quad \quad 5.4 \text{ g}$$

$$\text{Mass of reactants} = 3 + 11.2 = 14.2 \text{ g}$$

$$\text{Mass of products} = 8.8 + 5.4 = 14.2 \text{ g}$$

$$\text{Mass of reactants} = \text{Mass of products}$$


Thus the data illustrate law of conservation of mass.

Sol.26

Noble gases do form some chemical compounds, e.g., XeF₅, XeF₄, XeO₃ etc

Sol.27

CaF₂ is ionic solid; H₂O and Cl₂ are molecular solids ;SiC is a covalent solid.

Sol.28**Sol.29**

M. O. energy level diagram for O₂ molecule indicates that two electrons are unpaired so it is paramagnetic.

Sol.30

CuF₂, d⁹ system is coloured due to d-d transitions

PHYSICS**Sol.1**

$$[T] \propto [\rho]^x [r]^y [s]^z$$

$$[T] \propto [ML^{-3}]^x [L]^y [MT^{-2}]^z$$

$$\Rightarrow 0 = x + z$$

$$0 = -3x + y$$

$$1 = 2z \Rightarrow z = \frac{1}{2}$$

$$\text{Hence } x = \frac{1}{2}, y = \frac{3}{2}$$

$$\therefore T = \left[\frac{\rho r^3}{5} \right]^{1/2}$$

Sol.2

Using $R = \frac{u^2 \sin 2\theta}{2g}$ we get

$$\frac{g}{u^2} = \frac{\sin 2\theta}{R}$$

As range $R = 6 + 18 = 24\text{m}$

$$\therefore \frac{g}{u^2} = \frac{\sin 2\theta}{24}$$

$$\text{Again } y = x \tan \theta - \frac{gx^2}{2u^2 \cos^2 \theta}$$

For $x = 6\text{m}$, $y = 3\text{m}$ and $\frac{g}{u^2} = \frac{\sin 2\theta}{24}$, we get $3 = 6 \tan \theta - \frac{\sin 2\theta}{2 \times 24} \cdot \frac{6^2}{\cos^2 \theta}$

Again $\sin \theta = 2 \sin \theta \cos \theta$

We get $\tan \theta = \frac{2}{3}$ or $\theta = \tan^{-1} \left(\frac{2}{3} \right)$

Sol.3

As impulse = change in momentum

$$\therefore \text{Force of machine gun} = \frac{40}{1000} \times 1200 = 48\text{N}$$

$$\text{Force of Man} = 144\text{N} \Rightarrow \text{No. of bullets} = \frac{144}{48} = 3$$

Sol.4

$$\text{As } K \propto \frac{1}{L} \Rightarrow K' \times \frac{2L}{3} = KL \text{ or } K' = \frac{3}{2}K$$

Sol.5

No horizontal force will act on the rod as the surface is smooth. The only vertical forces acting on it are its own weight and normal reaction. Therefore centre of mass should fall vertically downwards towards negative y-axis. The path will be a straight line.

Sol.6

Total workdone = $4 \times$ Potential energy along sides + $2 \times$ Potential energy along diagonals

$$= 4 \times \left[-\frac{Gm_2m_2}{0.2} \right] + 2 \left[-\frac{Gm_1m_2}{0.2\sqrt{2}} \right]$$

$$= 4 \times \left[-\frac{(6.67 \times 10^{-11})(0.1)^2}{0.2} \right] + 2 \left[-\frac{(6.67 \times 10^{-11})(0.1)^2}{0.2\sqrt{2}} \right]$$

$$= -1.33 \times 10^{-11} - 0.47 \times 10^{-11} = -1.8 \times 10^{-11}\text{J}$$

Sol.7

As we know $V_c = R \frac{n}{\rho D}$, $R = \text{Reynold's number}$

For laminar flow, Reynold's number

$$R = 2000, \eta = 10^{-3} \text{ Nm}^{-2} \text{ s}^{-1}, \rho = 10^3 \text{ kgm}^{-3}$$

$$D = 2 \text{ cm} = 2 \times 10^{-2} \text{ m}$$

$$\Rightarrow V_c = \frac{2000 \times 10^{-3}}{10^3 \times 2 \times 10^{-2}} = 0.1 \text{ m/s}$$

Sol.8

As internal energy $U = n \left[\frac{F}{2} RT \right]$

$F = \text{degrees of freedom}$

Given $U = V_0 + U_{Ar}$

$$= 2 \times \frac{5}{2} RT + 4 \times \frac{3}{2} RT = 11RT$$

Sol.9

Given $\frac{d\delta}{dx} = -\delta a$

$\delta = \text{density}$, $a = \text{acceleration}$

If gas is accelerated in positive x direction, pressure will decrease, thereby implying that pressure is lower on the front side.

Sol.10

The amplitude and frequency are two independent characteristics of simple Harmonic Motion.

Sol.11

The frequency of open pipe

$$n_1 = \frac{v}{2(l+2e)} = \frac{330}{2(l+2 \times 0.3d)} \quad (\text{Ase} = 0.3d) = \frac{330}{2(l+0.6d)}$$

Frequency of closed pipe

$$n^2 = \frac{v}{4(l+e)} = \frac{330}{4(l+0.3d)}$$

$$\Rightarrow \frac{n_2}{n_1} = \frac{2(l+0.6d)}{4(l+0.3d)} = \frac{1(l+0.6d)}{2(l+0.3d)}$$

Sol.12

From the relation

kinetic Energy = $q\Delta V$, we get

$$K.E. = 2 \times 1.6 \times 10^{-19} (70 - 50)J = 40eV$$

Sol.13

Making use of relation

$$V = IR \Rightarrow \frac{V}{R}$$

Here $V = 15V$ and $R = 4 + 6 + 10 + 2 = 22\Omega$

$$\therefore I = \frac{15}{22} = 0.6A \text{ i.e. } r = 5\Omega$$

Sol.14

Magnetic field at the middle of solenoid

$$B = \mu_0 nI = \mu_0 \frac{N}{L} I = 4\pi \times 10^{-7} \times \frac{500}{0.4} \times 3 = 4.713 \times 10^{-3} T$$

Magnetic dipole moment of the coil

$$M = NIA = NI\pi r^2 = 10 \times 0.4 \times 3.142 \times (0.01)^2 = 1.26 \times 10^{-3} Am^2$$

Torque acting on the coil

$$\tau = MB \sin\theta \text{ As } \theta = 90^\circ$$

$$\tau = MB = 1.26 \times 10^{-3} \times 4.713 \times 10^{-3} = 5.94 \times 10^{-6} Nm$$

Sol.15

Here the proton has no acceleration so $E = 0, B = 0$

Sol.16

The input and output powers should be same for 100% efficiency.

Sol.17

$$\text{As we know that } E = \frac{LdI}{dt} \Rightarrow L = \frac{Edt}{dI} = \frac{20 \times 0.05}{18.2} = 62.5 \times 10^{-3} H = 62.5mH$$

Sol.18

$$\text{As } \lambda = \frac{c}{\nu} = \frac{3 \times 10^8}{2 \times 10^{10}} = 1.5 \times 10^{-2}$$

Sol.19

Path difference for a point

$$= (d^2 + b^2)^{1/2} - d = \frac{b^2}{2d}$$

Path difference for a dark bonds = $(2n - 1)\frac{\lambda}{2}$

$$\Rightarrow (2n - 1)\frac{\lambda}{2} = \frac{b^2}{2d} \text{ or } (2n - 1)\lambda = \frac{b^2}{d}$$

$$\text{for } n = 1, \lambda = \frac{b^2}{d}$$

Sol.20

$$\text{As } P = \frac{1}{f} = \frac{1}{f_1} - \frac{1}{f_2} = \frac{1}{0.5} - \frac{1}{1} = 1.0D$$

Sol.21

The length of telescope tube would increase by an amount equal to $4f$

Sol.22

X-rays have a certain minimum wavelength and also the wavelength larger than this minimum value.

Sol.23

$$\text{As } \mu = \text{current} \times \text{area} = \frac{q\omega}{2\pi} \times \pi r^2 = \frac{1}{2}\omega q r^2$$

Orbital angular momentum

$$L = m\omega r^2 = \frac{h}{2\pi} = h$$

$$\Rightarrow \omega r^2 = \frac{h}{m}$$

$$\therefore \mu = \frac{1}{2} \frac{qh}{m} = \frac{1.6 \times 10^{-19} \times 1.05 \times 10^{-34}}{2 \times 9.1 \times 10^{-31}} = 9.2 \times 10^{-24} \text{ Am}^2$$

Sol.24

As per the penetrating power

$$P_\gamma < P_\beta < P_\alpha$$

Sol.25

$$Eg = hv = \frac{hc}{\lambda}$$

$$\text{Given } \lambda = 2480nm = 2480 \times 10^{-9}m = 248 \times 10^{-8}m$$

$$\Rightarrow Eg = \frac{6.6 \times 10^{-34} \times 3 \times 10^8}{248 \times 10^{-8}} = 7.984 \times 10^{-20} J$$

$$= \frac{7.984 \times 10^{-20}}{1.6 \times 10^{-19}} eV = 0.499 eV = 0.5 eV$$

Sol.26

$$\text{Power gain} = \frac{a^2 R_L}{R_{in}} = \left(\frac{25}{26}\right)^2 \times \frac{800}{200} = 3.69$$

Sol.27

$$\text{Emitter Current, } I_e = \frac{n_e \times e}{t} = \frac{10^{10} \times 1.6 \times 10^{-19}}{10^{-6}} = 1.6 \times 10^{-3} A = 1.6 mA$$

Sol.28

$$[Y] = \left[\frac{x}{z^2}\right] = \left[\frac{\text{capacitance}}{(\text{magnetic induction})}\right]$$

$$= \frac{[M^{-1}L^{-2}Q^2T^2]}{[M^2Q^{-2}T^{-2}]} = [M^{-3}L^{-2}T^4Q^4]$$

Sol.29

A person has to swim perpendicular to the river current in order to cross the river in shortest time.


Sol.30

$$L = m \frac{v}{\sqrt{2}} r \perp$$

$$\text{Here } r \perp = h = \frac{v^2 \sin^2 45^\circ}{2g} = \frac{v^2}{4g} \text{ or } L = m \left(\frac{v}{\sqrt{2}}\right) \left(\frac{v^2}{4g}\right) = \frac{mv^3}{4\sqrt{2}g}$$

MATHEMATICS

Sol.1



Sol.2

When the left most integer is subtracted from its corresponding real number, the result will always range between 0 and 1 (0 included but 1 not included).

Sol.3

Let the given Relation be R. In the Relation R from $A \rightarrow B$, $x \in A$ and $y \in B$ and (x,y) . Hence A is $\{2,4,6\}$.

Sol.4

As α is the root of the equation $ax^2 + bx + c = 0$.

Therefore, $a\alpha^2 + b\alpha + c = 0$.

$$\text{Hence } \frac{1}{a\alpha+b} = -\frac{\alpha}{c} \quad (\text{i})$$

$$\text{Similarly } \frac{1}{a\beta+b} = -\frac{\beta}{c} \quad (\text{ii})$$

$$S = \text{Sum of roots} = \left(-\frac{\alpha}{c}\right) + \left(\frac{\beta}{c}\right) = -\frac{(\alpha+\beta)}{c} = \frac{b}{ac} \quad (\text{because } \alpha + \beta = -b/a) \quad \text{Product of roots} = \frac{\alpha\beta}{c^2} = \frac{1}{ac}$$

Equation is $x^2 - Sx + P = 0$

$$x^2 - \frac{b}{ac}x + \frac{1}{ac} = 0$$

$$acx^2 - bx + 1 = 0.$$

Sol.5

$$x = \sqrt{a+x} \text{ or } x^2 = a+x$$

$$\Rightarrow x^2 - x - a = 0$$

$$\Rightarrow x = \frac{1 \pm \sqrt{1+4a}}{2}$$

Sol.6

$$\begin{vmatrix} 1 + \sin^2 \theta & \cos^2 \theta & 4 \sin 4\theta \\ \sin^2 \theta & 1 + \cos^2 \theta & 4 \sin 4\theta \\ \sin^2 \theta & \cos^2 \theta & 1 + 4 \sin 4\theta \end{vmatrix} = 0$$

$$[C_1 \rightarrow C_1 + C_2]$$

$$\Rightarrow \begin{vmatrix} 2 & \cos^2 \theta & 4 \sin 4\theta \\ 2 & 1 + \cos^2 \theta & 4 \sin 4\theta \\ 1 & \cos^2 \theta & 1 + 4 \sin 4\theta \end{vmatrix} = 0$$

$$[R_2 \rightarrow R_2 - R_1, R_3 \rightarrow R_3 - R_1]$$

$$\Rightarrow \begin{vmatrix} 2 & \cos^2 \theta & 4 \sin 4\theta \\ 0 & 1 & 0 \\ -1 & 0 & 1 \end{vmatrix} = 0$$

$$\Rightarrow 2 + 4 \sin 4\theta = 0$$

$$\Rightarrow \sin 4\theta = -\frac{1}{2}$$

$$\Rightarrow 4\theta = \frac{7\pi}{24}, \frac{11\pi}{24}$$

Sol.7

For non-zero solution $\Delta=0$

$$\Rightarrow \begin{vmatrix} 1 & -K & -1 \\ K & -1 & -1 \\ 1 & 1 & -1 \end{vmatrix}$$

On solving, we get $K^2 - 1 = 0 \Rightarrow K = \pm 1$

Sol.8

We have total number of persons = 3 girls + 9 boys = 12

The total number of numbered seats = $2 \times 3 + 4 \times 3 = 14$ So, the total number of ways in which 12 persons can be seated on 14 seats = number of arrangements of 14 seats by taking 12 at a time = ${}^{14}P_{12}$. Three girls can be seated together in a back row on adjacent seats in the following way:

1, 2, 3 or 2, 3, 4 of first van and 1, 2, 3 or 2, 3, 4 of second van.

In each way the three girls can interchange among themselves in $3!$ ways. So, the total number of ways in which three girls can be seated together in a back row on adjacent seats = $4 \times 3!$

Now, 9 boys are to be seated on remaining 11 seats, which can be done in ${}^{11}P_9$ ways. Hence, by the fundamental principle of counting, the total number of seating arrangement is ${}^{11}P_9 \times 4 \times 3! = {}^{11}P_9 \times 4!$

Sol.9

Arrangement of n things in circle irrespective of the direction = $\frac{(n-1)!}{2} = \frac{4!}{2} = 12$

Sol.10

$$\begin{aligned} \left(\frac{a}{a+x}\right)^{\frac{1}{2}} + \left(\frac{a}{a-x}\right)^{\frac{1}{2}} &= \left(\frac{a+x}{a}\right)^{-\frac{1}{2}} + \left(\frac{a-x}{a}\right)^{-\frac{1}{2}} \\ &= \left(1 + \frac{x}{a}\right)^{-\frac{1}{2}} + \left(1 - \frac{x}{a}\right)^{-\frac{1}{2}} \\ &= \left[1 + \left(-\frac{1}{2}\right)\left(\frac{x}{a}\right) + \frac{\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)}{2 \cdot 1}\left(\frac{x}{a}\right)^2 + \dots\right] \\ &+ \left[1 + \left(-\frac{1}{2}\right)\left(-\frac{x}{a}\right) + \frac{\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)}{2 \cdot 1}\left(-\frac{x}{a}\right)^2 + \dots\right] \\ &= 2 + \frac{3x^2}{4a^2} + \dots \end{aligned}$$

Sol.11

Put $x = 1$ in the expansion of $(1 - 3x + 10x^2)^n$,

We get

$$(1 - 3 + 10(1)^2)^n = a$$

$$\Rightarrow a = (1 - 3 + 10)^n = 8^n$$

$$\Rightarrow a = 2^{3n} \quad (\text{i})$$

Put $x = 1$ in the expansion of $(1 + x^2)^n$, we get $(1 + 1)^n = b$

$$\Rightarrow b = 2^n \quad (\text{ii})$$

From (i) and (ii), we get $a = b^3$

Sol.12

Let T_n be the n^{th} term of the series $3 + 10 + 17 + \dots$

$$\text{Therefore } T_n = 3 + (n - 1)7 = 7n - 4$$

Let T_n' be the n^{th} term of the series $63 + 65 + 67 + \dots$

$$\text{Therefore } T_n' = 63 + (n - 2)2 = 2n + 61$$

$$\text{Now } T_n = T_n' \Rightarrow 7n - 4 = 2n + 61 \Rightarrow n = 13$$

Sol.13

Given that a, b, c are in A.P, and $(b - a), (c - b), a$ are in G.P, therefore

$$\text{we get } 2b = a + c \text{ and } (c - b)^2 = (b - a) a$$

$$\Rightarrow (b - a)^2 = (b - a) a \quad (\text{using } c - b = a - b)$$

$$\Rightarrow (b - a) = a$$

$$\Rightarrow b = 2a$$

$$\Rightarrow c = 3a \quad (\text{using } 2b = a + c).$$

Therefore, $a : b : c = 1 : 2 : 3$

Sol.14

$$\text{We have } \lim_{x \rightarrow 0} \frac{(1-x)^n - 1}{x}$$

On applying the L'Hospital rule, we get

$$\lim_{x \rightarrow 0} \frac{-n(1-x)^{n-1}}{1} = -n$$

Hence, option (c) is correct.

Sol. 15

$$Rf'(2) = \lim_{h \rightarrow 0} \frac{f(2+h) - f(2)}{h}$$

$$\lim_{h \rightarrow 0} \frac{|2+h| - |2|}{2}$$

$$\lim_{h \rightarrow 0} \frac{2+h-2}{h} = \lim_{h \rightarrow 0} \frac{h}{h} = 1$$

$$Lf'(2) = \lim_{h \rightarrow 0} \frac{f(2-h) - f(2)}{-h}$$

$$= \lim_{h \rightarrow 0} \frac{|2-h| - |2|}{-h}$$

$$= \lim_{h \rightarrow 0} \frac{2-h-2}{-h} = \lim_{h \rightarrow 0} \frac{h}{h} = 1$$

$$\text{Hence } Lf'(2) = Rf'(2) = 1$$

$$\text{Therefore } f'(2) = 1$$

Sol.16

We have $y - e^{xy} + x = 0$ (i)

Hence, on differentiating (i), we get

$$y' - e^{xy}(xy' + y) + 1 = 0$$

$$\text{or } y' = \frac{ye^{xy}-1}{1-xe^{xy}}$$

At any point the vertical tangent would have its slope as infinite. Now equating the denominator with 0, we get $1 - xe^{xy} = 0$ (ii)

(1, 0) satisfies (ii)

Therefore only option (d) is correct.

Sol.17

$$I = \int e^{\log(x^{-1})} dx = \int x^{-1} dx = \int \frac{1}{x} dx = \log|x|$$

Sol.18

$$I = \frac{1}{2} \int_{\frac{\pi}{2}}^{\pi} (1 - \cos 2x) dx = \frac{1}{2} \left[x - \frac{\sin 2x}{2} \right]_{\frac{\pi}{2}}^{\pi} = \frac{\pi}{4}$$

Sol.19

$$\frac{dr}{dt} = -rt \Rightarrow \frac{dr}{r} = -tdt \Rightarrow \log r = -\frac{t^2}{2} + c \quad (i)$$

Putting $t = 0$ and $r = r_0$ in (i), we get $c = \log r_0$

$$\Rightarrow \log r = -\frac{t^2}{2} + \log r_0$$

$$\text{Hence, } r = r_0 \left(e^{-t^2/2} \right)$$

Sol.20

$$\frac{dy}{dx} + \left(\frac{1-y^2}{1-x^2} \right)^{1/2} = 0$$

$$\Rightarrow \frac{dy}{\sqrt{1-y^2}} + \frac{dx}{\sqrt{1-x^2}} = 0$$

$$\Rightarrow \sin^{-1} y + \sin^{-1} x = \sin^{-1} c$$

$$\Rightarrow \sin^{-1} \left[y\sqrt{1-x^2} + x\sqrt{1-y^2} \right] = \sin^{-1} c \Rightarrow \left[y\sqrt{1-x^2} + x\sqrt{1-y^2} \right] = c$$

Sol.21

$$(x\sqrt{1+x^2})dx + (y\sqrt{1+y^2})dy = 0$$

$$\Rightarrow \frac{xdx}{\sqrt{1+x^2}} + \frac{ydy}{\sqrt{1+y^2}} = 0$$

$$\Rightarrow \sqrt{1+x^2} + \sqrt{1+y^2} = c$$

Sol.22 we know that $y = mx + c$ touches $y^2 = 4ax$, iff $c = a/m$ and $m \neq 0$

Therefore the line $y = 2x + c$ touches the curve $y^2 = 16x$ only if $c = 4/2 = 2$

Sol.23

We know that if α, β, γ are the directional angles with x-axis, y-axis and z-axis, then

$$\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1$$

$$\Rightarrow (1 - \sin^2 \alpha) + (1 - \sin^2 \beta) + (1 - \sin^2 \gamma) = 1$$

$$\Rightarrow \sin^2 \alpha + \sin^2 \beta + \sin^2 \gamma = 2$$

Sol.26

The total number of ways of choosing two numbers out of 1, 2, 3, ..., 30 is ${}^{30}C_2 = 435$

\Rightarrow Exhaustive number of cases = 435

Since $a^2 - b^2$ is divisible by 3 if either a and b both are divisible by 3 or none of a and b is divisible by 3.

Thus, the favourable number of cases

$$= {}^{10}C_2 + {}^{20}C_2 = 235$$

Hence, the required probability = $\frac{235}{435} = \frac{47}{87}$

Sol.29

$$b^2 = a^2(e_1^2 - 1) \text{ and } a^2 = b^2(e_2^2 - 1)$$

$$\Rightarrow b^2 = b^2(e_2^2 - 1)(e_1^2 - 1)$$

$$\Rightarrow 1 = (e_2^2 - 1)(e_1^2 - 1)$$

$$\Rightarrow e_1^2 e_2^2 - e_1^2 - e_2^2 = 0$$

$$\Rightarrow \frac{1}{e_1^2} + \frac{1}{e_2^2} = 1$$

Sol.30

The slope of the given tangent at any point (x, y) is $\frac{dy}{dx} = \frac{2x}{4-y^2}$

For a vertical tangent the slope must be infinite.

$$\text{Therefore } \frac{2x}{4-y^2} = \infty$$

$$\Rightarrow 4 - y^2 = 0$$

$$\Rightarrow y = \pm 2$$

$$\Rightarrow x = \pm \frac{4}{\sqrt{3}}$$