COMMON ENTRANCE TEST - 2007

DATE	SUBJECT	TIME
09 - 05 - 2007	PHYSICS & CHEMISTRY	10.00 AM to 12.30 PM
	(COMBINED PAPER)	

MAXIMUM MARKS	TOTAL DURATION	MAXIMUM TIME FOR ANSWERING
120	150 MINUTES	140 MINUTES

CET NUI	WBER	VERSION CODE	SERIAL NUMBER 370241
		Λ_1	370241

IMPORTANT INSTRUCTIONS TO CANDIDATES

(Candidates are advised to read the following instructions carefully, before answering on OMR answer sheet.)

- 1. Ensure that you have entered your Name and Register Number of 2^{nd} PUC Annual Examination / 12^{th} Std. in the space provided on the OMR answer sheet.
- 2. Ensure that CET No. has been entered and shaded the respective circles on the OMR answer sheet.
- 3. ENSURE THAT THE TIMING, MARKS PRINTED ON THE OMR ANSWER SHEET ARE NOT DAMAGED/MUTILATED/SPOILED.
- 4. This Question Booklet is issued to you by the invigilator after the 2nd Bell. i.e., after 10.00 a.m.
- 5. Enter the Serial Number of this question booklet on the OMR answer sheet.
- 6. Carefully enter the Version Code of this question booklet on the OMR answer sheet and SHADE the respective circles completely.
- 7. As answer sheets are designed to suit the Optical Mark Reader (OMR) system, please take special care while filling and shading the CET NO. & Version Code of this question booklet.
- 8. DO NOT FORGET TO SIGN AT THE BOTTOM PORTION OF OMR ANSWER SHEET IN THE SPACE PROVIDED.
- 9. Until the 3rd Bell is rung at 10.10 a.m.:
 - Do not remove the seal present on the right hand side of this question booklet.
 - Do not look inside this question booklet.
 - Do not start answering on the OMR answer sheet.
- 10. After the 3rd Bell is rung at 10.10 a.m., remove the seal present on the right hand side of this question booklet and start answering on the OMR answer sheet.
- 11. This question booklet contains 120 questions and each question will have four different options / choices.
- 12. During the subsequent 140 minutes:
 - Read each question carefully.
 - Determine the correct answer from out of the four available options / choices given under each question.
 - Completely darken / shade the relevant circle with a BLUE OR BLACK INK BALLPOINT PEN against the question number on the OMR answer sheet.

CORRECT METHOD OF SHADING THE CIRCLE ON THE OMR SHEET IS AS SHOWN BELOW:



- 13. Please note that even a minute unintended ink dot on the OMR sheet will also be recognised and recorded by the scanner. Therefore, avoid multiple markings of any kind on the OMR answer sheet.
- 14. Use the space provided on each page of the question booklet for Rough work AND do not use the OMR answer sheet for the same.
- 15. After the last bell is rung at 12.30 p.m., stop writing on the OMR answer sheet and affix your LEFT HAND THUMB IMPRESSION on the OMR answer sheet as per the instructions.
- 16. Hand over the OMR ANSWER SHEET to the room invigilator as it is.
- 17. After separating and retaining the top sheet (CET Cell Copy), the invigilator will return the bottom sheet replica (Candidate's copy) to you to carry home for self-evaluation.
- 18. Preserve the replica of the OMR answer sheet for a minimum period of One year.





PHYSICS

A ray of light is travelling from glass to air. (Refractive index of glass = 1.5) The angle of 1. incidence is 50°. The deviation of the ray is

1)
$$Sin^{-1} \left[\frac{Sin \, 50^0}{1.5} \right] - 50^0$$

2)
$$50^0 - Sin^{-1} \left[\frac{Sin50^0}{1.5} \right]$$

 $3) 80^0$

4)

A vessel of height 2 d is half filled with a liquid of refractive index $\sqrt{2}$ and the other half 2. with a liquid of refractive index n. (The given liquids are immiscible). Then the apparent depth of the inner surface of the bottom of the vessel (neglecting the thickness of the bottom of the vessel) will be

$$1) \quad \frac{nd}{d + \sqrt{2n}}$$

2)
$$\frac{\sqrt{2n}}{d(n+\sqrt{2})}$$
4)
$$\frac{n}{d(n+\sqrt{2})}$$

$$3) \quad \frac{d\left(n+\sqrt{2}\right)}{n\sqrt{2}}$$

4)
$$\frac{n}{d(n+\sqrt{2})}$$

- 3. A ray of light is incident normally on one face of a right angled isosceles prism. It then grazes the hypotenuse. The refractive index of the material of the prism is
 - 1) 1.732

2) 1.5

3) 1.414

- 1.33 4)
- 4. Two thin equiconvex lenses each of focal length 0.2 m are placed coaxially with their optic centres 0.5 m apart. Then the focal length of the combination is
 - 1) 0.1 m

 $2) - 0.1 \,\mathrm{m}$

3) 0.4 m

- 4) 0.4 m
- 5. A prism of a certain angle deviates the red and blue rays by 80 and 120 respectively. Another prism of the same angle deviates the red and blue rays by 10° and 14° respectively. The prisms are small angled and made of different materials. The dispersive powers of the materials of the prisms are in the ratio
 - 1) 11:9

2) 6:5

3) 9:11

4) 5:6

The electro magnetic theory of light failed to explain

	1)	Interference	2)	Diffraction
	· .	Polarisation	4)	Photo electric effect
7.	the scree	om two coherent sources of the en. The intensity of the central y at the same point will be	e same ai maximu	mplitude A and wavelength λ illuminates m is I_0 . If the sources were incoherent, the
	1)	$\frac{I_0}{2}$		I_0
	3)	$2I_0$	4)	$4I_0$
8.	In Youn slits 0.5	g's double slit experiment with 89 mm apart, the half angular	sodium v	vapour lamp of wavelength 589 nm and the the central maximum is
•		$Sin^{-1}0.1$		$Sin^{-1}0.001$
	3)	$Sin^{-1}0.0001$	4)	$Sin^{-1}0.01$
9.	A single	slit Fraunhoffer diffraction pat	tern is fo	rmed with white light. For what wavelength
	of light	the third secondary maximum	in the d	iffraction pattern coincides with the second
	seconda	ry maximum in the pattern fo	r red ligh	nt of wavelength $6500 \stackrel{0}{A}$?
	1)	9100A	2)	$^{0}_{4642.8A}$
	3)	$^0_{4100A}$	4)	$4400 \overset{0}{A}$
10.	The he	ad lights of a jeep are 1.2 m	apart. If	the pupil of the eye of an observer has a
	diamete	er of 2 mm and light of wavele	ngth 589	$\overset{0}{6A}$ is used, what should be the maximum
	distanc	e of the jeep from the observer	if the tv	vo head lights are just separated?
	1)			3.39 km
	3)) 33.9 m	4)	33.9 km

(Space for Rough Work)

6.

11.	When the angle of incidence is 60° on the surface of a glass slab, it is found that the r	eflected
	ray is completely polarised. The velocity of light in glass is	

1)
$$3 \times 10^8 \, \text{ms}^{-1}$$

2)
$$2 \times 10^8 \,\mathrm{ms}^{-1}$$

3)
$$\sqrt{3} \times 10^8 \,\mathrm{ms}^{-1}$$

4)
$$\sqrt{2} \times 10^8 \,\mathrm{ms}^{-1}$$

12. A 20 cm length of a certain solution causes right handed rotation of 38°. A 30 cm length of another solution causes left handed rotation of 24°. The optical rotation caused by 30 cm length of a mixture of the above solutions in the volume ratio 1:2 is

- 1) right handed rotation of 30
- 2) left handed rotation of 30
- 3) right handed rotation of 140
- 4) left handed rotation of 140

13. Two identical charges repel each other with a force equal to 10 mgwt when they are 0.6 m apart in air.($g = 10 \text{ ms}^{-2}$) The value of each charge is

1) $2\mu C$

2) 2nC

3) $2 \times 10^{-7} C$

4) 2mC

14. The potential of the electric field produced by a point charge at any point (x, y, z) is given by $V = 3x^2 + 5$, where x, y, z are in metres and V is in volts. The intensity of the electric field at (-2, 1, 0) is

1) $-12Vm^{-1}$

2) $+12Vm^{-1}$

3) $-17Vm^{-1}$

4) $+17Vm^{-1}$

15. The potential of a large liquid drop when eight liquid drops are combined is 20V. Then the pontential of each single drop was

1) 2.5 V

2) 5 V

3) 7.5 V

4) 10 V

- 16. Two indentical capacitors each of capacitance $5\mu F$ are charged to potentials 2 kV and 1 kV respectively. The –ve ends are connected together. When the +ve ends are also connected together, the loss of energy of the system is
 - 1) 1.25 J

2) 5 J

3) 0 J

- 4) 160 J
- 17. A parallel plate capacitor with air as the dielectric has capacitance C. A slab of dielectric constant K and having the same thickness as the separation between the plates is introduced so as to fill one-fourth of the capacitor as shown in the figure. The new capacitance will be



1) $\frac{KC}{4}$

 $2) \quad (K+1)\frac{C}{4}$

3) $(K+2)\frac{C}{4}$

- 4) $(K+3)\frac{C}{4}$
- 18. A current of 5 A is passing through a metallic wire of cross-sectional area $4 \times 10^{-6} \,\mathrm{m}^2$. If the density of charge carriers of the wire is $5 \times 10^{26} \,\mathrm{m}^{-3}$, the drift velocity of the electrons will be
 - 1) $1 \times 10^{-2} \, \text{ms}^{-1}$


2) $1.56 \times 10^{-3} \,\mathrm{ms}^{-1}$

3) $1.56 \times 10^{-2} \text{ms}^{-1}$

- 4) $1 \times 10^2 \text{ms}^{-1}$
- 19. Two bulbs rated 25 W 220 V and 100 W 220 V are connected in series to a 440 V supply. Then,
 - 1) both the bulbs fuse
- 2) neither of the bulbs fuses

3) 25 W bulb fuses

- 4) 100 W bulb fuses
- 20. The current passing through the ideal ammeter in the circuit given below is



21. In the Wheatstone's network given below,

$$P = 10 \Omega$$

$$Q = 20 \Omega$$

$$R = 15 \Omega$$
,

$$S = 30\Omega$$

The current passing through the battery (of negligible internal resistance) is



2) 0.18 A

Q

4) 0.36 A

22. A circular coil carrying a certain current produces a magnetic field *Bo* at its centre. The coil is now rewound so as to have 3 turns and the same current is passed through it. The new magnetic field at the centre is

$$2) \quad \frac{B_0}{3}$$

4)
$$\frac{Bc}{9}$$

23. A proton and a deuteron with the same initial kinetic energy enter a magnetic field in a direction perpendicular to the direction of the field. The ratio of the radii of the circular trajectories described by them is

3)
$$1:\sqrt{2}$$

24. Two tangent galvanometers A and B have coils of radii 8 cm and 16 cm respectively and resistance 8Ω each. They are connected in parallel with a cell of emf 4 V and negligible internal resistance. The deflections produced in the T.G's A and B are 30° and 60° respectively. If A has 2 turns, then B must have

- 4) 18 turns
- 25. A charged particle is moving in a magnetic field of strength B perpendicular to the direction of the field. If q and m denote the charge and mass of the particle respectively. Then the frequency of rotation of the particle is


$$1) \quad f = \frac{2\pi m}{qB}$$

$$2) \quad f = \frac{2\pi^2 m}{qB}$$


$$3) \quad f = \frac{qB}{2\pi m^2}$$

$$f = \frac{qB}{2\pi m}$$

26. A and B are two infinitely long straight parallel conductors. C is another straight conductor of length 1 m kept parallel to A and B as shown in the figure. Then the force experienced by C is



- 2) towards A equal to $5.4 \times 10^{-5} N$
- 3) towards B equal to $5.4 \times 10^{-5} N$
- 4) towards A equal to $0.6 \times 10^{-5} N$



27. An electric bulb has a rated power of 50 W at 100 V. If it is used on an a.c. source 200 V, 50Hz, a choke has to be used in series with it. This choke should have an inductance of

1) 1.1 H

2) 0.1 H

3) 1 mH

4) 0.1 mH

28. An inductance of $\frac{200}{\pi}$ mH, a capacitance of $\frac{10^{-3}}{\pi}$ F and a resistance of 10Ω are connected in series with an a.c. source 220 V, 50Hz. The phase angle of the circuit is

1) $\frac{\pi}{3}$

 $2) \quad \frac{\pi}{2}$

3) $\frac{\pi}{4}$

4) $\frac{\pi}{6}$

29. A stepdown transformer reduces the voltage of a transmission line from 2200 V to 220 V. The power delivered by it is 880 W and its efficiency is 88%. The input current is

1) 4.65 A

2) 0.465 A

3) 0.0465 A

(4) 4.65 mA

30. Current in a coil changes from 4 A to zero in 0.1 second and the emf induced is 100 V. The self inductance of the coil is

1) 4 H

2) 2.5 H

3) 0.4 H


4) 0.25 H


- 31. All components of the electromagnetic spectrum in vacuum have the same
 - 1) Frequency


2) Wavelength


3) Velocity

- 4) Energy
- 32. Which one of the following graphs represents the variation of maximum kinetic energy (E_K) of the emitted electrons with frequency γ in photoelectric effect correctly?









- 1) 1
- 3) 3

- 2) 2
- 4) 4
- 33. A and B are two metals with threshold frequencies $1.8\times10^{14}\,\mathrm{Hz}$ and $2.2\times10^{14}\,\mathrm{Hz}$. Two identical photons of energy 0.825 eV each are incident on them. Then photoelectrons are emitted in
 - 1) A alone

2) B alone

3) in both A and B

4) in neither A nor B

(Take $h = 6.6 \times 10^{-34} \text{ Js}$)

- **34.** The ionization energy of L_i^{++} is equal to
 - 1) hcR

2) 2 hcR

3) 6 hcR

- 4) 9 hcR
- **35.** Electrons in a certain energy level n = n, can emit 3 spectral lines. When they are in another energy level $n = n_2$. They can emit 6 spectral lines. The orbital speeds of the electrons in the two orbits are in the ratio
 - 1) 1:2

2) 2:1

3) 3:4

4) 4:3

- 36. The deBroglie wavelength of a proton (charge = $1.6 \times 10^{-19} C$, mass = $1.6 \times 10^{-27} kg$) accelerated through a p.d of 1 kV is
 - 1) 0.9 nm

2) $7A^{0}$

3) $0.9 \times 10^{-12} \text{ m}$

- 4) $600 \stackrel{0}{A}$
- 37. A radio active element forms its own isotope after 3 consecutive disintegrations. The particles emitted are
 - 1) 2α particles and 1β particle
- 2) 2β particles and 1γ particle
- 3) 2β particles and 1α particle
- 4) 3β particles
- **38.** A radio active substance contains 10,000 nuclei and its half life period is 20 days. The number of nuclei present at the end of 10 days is
 - 1) 7,500

2) 8,000

3) 9,000

- 4) 7,070
- 39. In Raman effect, Stokes' lines are spectral lines having
 - 1) wavelength greater than that of the original line.
 - 2) wavelength less than that of the original line.
 - 3) wavelength equal to that of the original line.
 - 4) frequency greater than that of the original line.
- 40. The principle of LASER action involves
 - 1) Stimulated emission
 - 2) Population inversion
 - 3) Amplification of particular frequency emitted by the system
 - 4) All of these

41. The volume of a nucleus is directly proportional to

1) $A^{\frac{1}{3}}$

2) \sqrt{A}

3) A^3

4) A

(Where A = mass number of the nucleus)

42. An electron is

1) A lepton

2) A nucleon

3) Baryon

4) Hadron

43. Minority carriers in a p-type semiconductor are

1) Holes

- 2) Free electrons
- 3) Both holes and free electrons
- 4) Neither holes nor free electrons

44. In a reverse biased diode when the applied voltage changes by 1 V, the current is found to change by $0.5\,\mu A$. The reverse bias resistance of the diode is

1) 2Ω

2) 200Ω

3) $2\times10^6\,\Omega$

4) $2\times10^5\,\Omega$

45. The truth table given below is for

B	Y
. 0	1
1	1
0	1
1	0
	0 1 0

(A and B are the inputs, Y is the output)

1) NAND

2) XOR

3) AND

4) NOR

- 46. The dimensional formula for impulse is
 - 1) $ML^{-1}T^{-1}$

2) $M^{-1}LT^{-1}$

3) $ML^{-1}T$

- 4) MLT^{-1}
- 47. The maximum height attained by a projectile when thrown at an angle θ with the horizontal is found to be half the horizontal range. Then θ =
 - 1) $Tan^{-1}\frac{1}{2}$

 $2) \frac{\pi}{4}$

3) $\frac{\pi}{6}$

- 4) $Tan^{-1}2$
- 48. A shell of mass 20 kg at rest explodes into two fragments whose masses are in the ratio 2:3. The smaller fragment moves with a velocity of 6 ms⁻¹. The kinetic energy of the larger fragment is
 - 1) 360 J

2) 144 J

3) 216 J

- 4) 96 J
- 49. Water rises in plant fibres due to
 - 1) Osmosis

2) Fluid pressure

3) Viscosity

- 4) Capillarity
- **50.** The acceleration due to gravity becomes $\left(\frac{g}{2}\right)$ where g = acceleration due to gravity on the surface of the earth at a height equal to
 - 1) $\frac{R}{2}$

2) 2*R*

3) $\frac{R}{4}$

4) 4*R*

- 51. The cylindrical tube of a spray pump has a cross-section of 8 cm², one end of which has 40 fine holes each of area 10⁻⁸ m². If the liquid flows inside the tube with a speed of 0.15 m.min⁻¹, the speed with which the liquid is ejected through the holes is
 - 1) 0.5 ms^{-1}

 $2) 0.05 \text{ ms}^{-1}$


3) 5 ms^{-1}

- 4) 50 ms^{-1}
- 52. During an adiabatic process, the cube of the pressure is found to be inversely proportional to the fourth power of the volume. Then the ratio of specific heats is
 - 1) 1.4

2) 1.67

3) 1.33

- 4) 1
- **53.** Two identical rods AC and CB made of two different metals having thermal conductivities in the ratio 2: 3 are kept in contact with each other at the end C as shown in the figure. A is at 100° C and B is at 25° C. Then the junction C is at



1) 50° C

2) 75° C

 $3) 60^{0}$ C

- 4) 55° C
- 54. 310 J of heat is required to raise the temperature of 2 moles of an ideal gas at constant pressure from 25°C to 35°C. The amount of heat required to raise the temperature of the gas through the same range at constant volume is
 - 1) 452 J

2) 276 J

3) 144 J

- 4) 384 J
- 55. A Carnot's engine operates with source at 127°C and sink at 27°C. If the source supplies 40 kJ of heat energy, the work done by the engine is
 - 1) 1 kJ

2) 4 kJ

3) 10 kJ

4) 30 kJ

- 56. The maximum particle velocity in a wavemotion is half the wave velocity. Then the amplitude of the wave is equal to
 - 1) λ

 $2) \quad \frac{\lambda}{2\pi}$

3) $\frac{2\lambda}{\pi}$

- 4) $\frac{\lambda}{4\pi}$
- 57. The ratio of the velocity of sound in hydrogen $\left(r = \frac{7}{5}\right)$ to that in helium $\left(r = \frac{5}{3}\right)$ at the same temperature is
 - $1) \quad \frac{\sqrt{21}}{5}$

 $2) \quad \frac{\sqrt{42}}{5}$

 $3) \quad \sqrt{\frac{5}{21}}$

- $4) \quad \sqrt{\frac{5}{42}}$
- 58. An engine is moving towards a wall with a velocity 50 ms⁻¹ emits a note of 1.2 kHz. Speed of sound in air = 350 ms⁻¹. The frequency of the note after reflection from the wall as heard by the driver of the engine is
 - 1) 1.2 kHz

2) 1.6 kHz

3) 0.24 kHz

- 4) 2.4 kHz
- 59. A glass tube is open at both the ends. A tuning fork of frequency f resonates with the air column inside the tube. Now the tube is placed vertically inside water so that half the length of the tube is filled with water. Now the air column inside the tube is in unison with another fork of frequency f'. Then
 - $1) \quad f' = \frac{f}{2}$

2) f' = 2f

3) f' = 4f

- 4) f'=f
- **60.** The surface temperature of the Sun which has maximum energy emission at 500 nm is 6000 K. The temperature of a star which has maximum energy emission at 400 nm will be
 - 1) 6500 K

2) 7500 K

3) 4500 K

4) 8500 K

16 A-1

A - 1

CHEMISTRY

18

A - 1

61. During the extraction of gold the following reactions take place -

$$Au + CN^{-} + H_{2}O \xrightarrow{O_{2}} [X]$$
$$[X] + Zn \xrightarrow{} [Y] + Au$$

5 . j.

X and Y are respectively -

1)
$$\left[Au\left(CN\right)_{2}\right]^{-}$$
 and $\left[Zn\left(CN\right)_{4}\right]^{2-}$ 2) $\left[Au\left(CN\right)_{4}\right]^{3-}$ and $\left[Zn\left(CN\right)_{4}\right]^{2-}$

3)
$$\left[Au\left(CN\right)_{4}\right]^{2-}$$
 and $\left[Zn\left(CN\right)_{4}\right]^{2-}$ 4) $\left[Au\left(CN\right)_{2}\right]^{-}$ and $\left[Zn\left(CN\right)_{6}\right]^{4-}$

- 62. The number of gram molecules of chlorine in 6.02×10^{25} hydrogen chloride molecules is
 - 1) 5

2) 50

3) 100

- 4) 10
- **63.** Graphite is a soft solid lubricant extremely difficult to melt. The reason for this anomalous behaviour is that graphite
 - 1) has molecules of variable molecular masses like polymers.
 - 2) has carbon atoms arranged in large plates of rings of strongly bound carbon atoms with weak interplate bonds.
 - 3) is a non-crystalline substance.
 - 4) is an allotropic form of carbon.
- 64. Paracetamol is a / an
 - 1) antimalarial

2) antipyretic

3) analgesic

- 4) both 2 and 3
- 65. Which one of the following has maximum number of atoms of oxygen?
 - 1) 2 g of water

- 2) 2 g of sulphur dioxide
- 3) 2 g of carbon dioxide
- 4) 2 g of carbon monoxide.

00.	which one of the following shows functional isomerism?					
	1)	CH_2Cl_2	2)	C_2H_5OH		
	3)	C_3H_6	4)	C_2H_4		•
67.	In the ic	onic equation $-BiO_3^-+6$	$H^+ + Xe^-$	$\rightarrow Bi^{3+} + 3H_2O$,		
	the valu	es of X is $-$		–		
	1)	3	2)	4		
	3)	2	4)	6		•
68.	Molarity	of a given orthophospho	oric acid solut	ion is 3M. It's noi	cmality is –	
	1)	1 N	2)	3 N	T	
	3)	0.3 N	4)	9 N		
69.	Acidified colourati	l sodium fusion extract on which confirms the p	on addition o presence of –	f ferric chloride s	solution gives bloo	d red
	1)	S	2)	N		-
	3)	N and S	4)	S and Cl		
70.		f mass 10 mg is moving v sociated with it would be		of 100 ms^{-1} . The	wavelength of de-B	roglie
	(Note: h	$=6.63\times10^{-34} \text{Js}$)			•	
	1)	$6.63 \times 10^{-37} \text{m}$	2)	6.63×10^{-31} m		
	3)	6.63×10^{-34} m	4)	6.63×10^{-35} m	•	

71. $M\sigma^{2+}$ is isoelectronic w	$I_{\mathcal{O}}^{2+}$ is isoelectronic	with
---------------------------------------	---	------

1) Ca^{2+}

2) Na^{-1}

3) Zn^{2+}

4) Cu^{2+}

72. Gram molecular volume of oxygen at STP is -

1) 11200 cm³

2) 22400 cm³

 $3) 5600 \text{ cm}^3$

4) 3200 cm^3

73. Presence of halogen in organic compounds can be detected using -

1) Beilstien's test

2) kjeldahl test

3) Duma's test

4) Leibig's test

74. The electronic configuration of
$$Cr^{3+}$$
 is

1) $[Ar]3d^54s^1$

 $2) \quad [Ar] 3d^2 4s^1$

3) $[Ar]3d^34s^0$

4) $[Ar]3d^44s^2$

75. The mass of a metal, with equivalent mass 31.75, which would combine with 8 g of oxygen is

1) 31.75

2) 3.175

3) 8

4) 1

1)	C_6H_5Cl $C_6H_6Cl_6$ eriodic table metals usually used s - block d - block	4) l as ca	atalysts belong to	,
In the p	eriodic table metals usually used s - block	l as ca	atalysts belong to	,
1)	s - block			
		2)	m blook	
3)	d - block		p - block	
	W NICOIL	4)	f - block	
Dalton's	law of partial pressures is appli	cable	to which one of the following syst	ems S
1)	$CO+H_2$	2)	$H_2 + Cl_2$	•
3)	$NO + O_2$	4)	$NH_3 + HCl$	
The gen	eral formula of a cycloalkane is			
1)	C_nH_{2n+2}	2)	C_nH_{2n-2}	
3)	C_nH_{2n}	4)	C_nH_n	
In acety	lene molecule, between the carbo	n ato	oms there are –	
1)	three sigma bonds	2)	two sigma and one pi bonds	
3)	one sigma and two pi bonds	4)	three pi bonds	
	(Space for F	lough	Work)	
	Dalton's 1) 3) The gen 1) 3) In acety 1)	1) $CO + H_2$ 3) $NO + O_2$ The general formula of a cycloalkane is 1) C_nH_{2n+2} 3) C_nH_{2n} In acetylene molecule, between the carbo 1) three sigma bonds 3) one sigma and two pi bonds	Dalton's law of partial pressures is applicable 1) $CO + H_2$ 2) 3) $NO + O_2$ 4) The general formula of a cycloalkane is 1) C_nH_{2n+2} 2) 3) C_nH_{2n} 4) In acetylene molecule, between the carbon atom 1) three sigma bonds 2) 3) one sigma and two pi bonds 4)	Dalton's law of partial pressures is applicable to which one of the following system: 1) $CO + H_2$ 2) $H_2 + Cl_2$ 3) $NO + O_2$ 4) $NH_3 + HCl$ The general formula of a cycloalkane is 1) C_nH_{2n+2} 2) C_nH_{2n-2} 3) C_nH_{2n} 4) C_nH_n In acetylene molecule, between the carbon atoms there are $-$ 1) three sigma bonds 2) two sigma and one pi bonds

ni. Deliaulieu alconor	atured	81.	d alcohol	is
------------------------	--------	-----	-----------	----

- 1) Rectified spirit
- 2) Undistilled ethanol
- 3) Rectified spirit + methanol + naphtha
- 4) Ethanol + methanol

82. During the formation of a chemical bond

- 1) energy decreases
- 2) energy increases
- 3) energy of the system does not change
- 4) electron-electron repulsion becomes more than the nucleus-electron attraction
- 83. One mole of oxygen at 273 k and one mole of sulphur dioxide at 546 k are taken in two separate containers, then,
 - 1) kinetic energy of O_2 > kinetic energy of SO_2 .
 - 2) kinetic energy of O_2 < kinetic energy of SO_2 .
 - 3) kinetic energy of both are equal.
 - 4) None of these

84. +I effect is shown by

 $1) -NO_2$

2) *-Cl*

3) -Br

4) -CH₃

85. Formation of coloured solution is possible when metal ion in the compound contains

1) paired electrons

- 2) unpaired electrons
- 3) lone pair of electrons
- 4) none of these

86.	wnich o	i the following is an intensive p	ropert	у:
	1)	temperature	2)	surface tension
	3)	viscosity	4)	all of these
87.	Hofman	n's bromamide reaction is to con	vert	
	1)	amine to amide	2)	amide to amine
	3)	alcohol to acid	4)	acid to alcohol
88.	IUPAC :	name of $Na_3igl[{\it Co(NO_2)}_6igr]$ is		
	1)	sodium cobaltinitrite	2)	sodium hexanitrito cobaltate (III)
	3)	sodium hexanitro cobalt (III)	4)	sodium hexanitrito cobaltate (II)
89.	Thermo	dynamic standard conditions of	tempe	rature and pressure are
	1)	0^0 C and 1 atm	2)	273 k and 101.3 k Pa
	3)	298 k and 1 atm	4)	0^0 C and 101.3 k Pa
90.	How ma	ny chiral carbon atoms are prese	ent in	2, 3, 4 - trichloropentane?
	1)	3	2)	2
	3)	1	4)	4
		<u> </u>	-	

- 91. The number of unidentate ligands in the complex ion is called
 - 1) EAN

2) Coordination number

3) primary valency

- 4) oxidation number
- **92.** $2SO_{2(g)} + O_{2(g)} \xrightarrow{V_2O_5}$ is an example for
 - 1) irreversible reaction
- 2) heterogenous catalysis
- 3) homogenous catalysis
- 4) neutralisation reaction
- 93. The amino acid which is not optically active is
 - 1) glycine

2) alanine

3) serine

- 4) lactic acid
- **94.** For a stable molecule the value of bond order must be
 - 1) negative
 - 2) positive
 - 3) zero
 - 4) there is no relationship between stability and bond order.
- 95. Which one of the following is a second order reaction?
 - 1) $CH_3COOCH_3 + NaOH \longrightarrow CH_3COONa + H_2O$
 - 2) $H_2 + Cl_2 \xrightarrow{\text{sunlight}} 2HCl$
 - 3) $NH_4NO_3 \longrightarrow N_2 + 3H_2O$
 - 4) $H_2 + Br_2 \longrightarrow 2HBr$

96.	Accordin	ng to Bayer's strain theory which	is hi	ghly stable?	
	1)	cyclohexane	2)	cycloheptane	
	3)	cyclopentane	4)	cyclobutane	
97.	The nun	nber of antibonding electron pairs	$\sin O_2^2$	2– molecular ion on the bas	is of molecular
	orbital t	heory is			
	[Note - A	Atomic number of O is 18]	•		
	1)	2	2)	3,	
	3)	4	4)	5	
98.	Hydroxy	rl ion concentration of 1M HCl is	S		
	1)	$1\times10^{-14}\mathrm{mol}\;\mathrm{dm}^{-3}$	2)	1×10^{-1} mol dm ⁻³	
	. 3)	$1\times10^{-13}\mathrm{mol}\;\mathrm{dm}^{-3}$	4)	$1\times10^1\mathrm{mol\ dm}^{-3}$. 3
99.	Geometr	rical isomerism is shown by			
	1)	-C-C-	2)	$-C \equiv C$ –	
	3)	-C - C - $C = C'$	4)	None of these	•
100.	The oxid	dation state of iron in $K_4[Fe(C)]$	$N)_6$	is	
	1)	2	. 2)	3	
	3)	4	4)	1	

101. In which of the following process, a maximum increase in entropy is observed?

- 1) dissolution of salt in water
- 2) condensation of water
- 3) sublimation of naphthalene
- 4) melting of ice

102. Decomposition of benzene diozonium chloride by using Cu_2Cl_2/HCl to form chlorobenzene is

- 1) Cannizarro's reaction
- 2) Kolbe's reaction
- 3) Sandmeyer's reaction
- 4) Raschig's reaction

103. Which complex can not ionise in solution?

1) $[pt(NH_3)_6]Cl_4$

2) $K_2[pt(F_6)]$

3) $K_4[Fe(CN)_6]$

4) $\left[CoCl_3 \left(NH_3 \right)_3 \right]$

104. Considering the reaction $C_{(s)} + O_{2(g)} \rightarrow CO_{2(g)} + 393.5 \text{ kJ}$ the signs of ΔH , ΔS and ΔG respectively are

1) -, +, -

2) -,-,-

3) - + +

4) +, -, -

105. The product formed when hydroxylamine condenses with a carbonyl compound is called

1) hydrazone

2) hydrazine

3) oxime

4) hydrazide

106. Whi	ch o	f the following forms a colourles	s solu	ition in aqueous medium?
	1)	Ti ³⁺	2)	Sc^{3+}
	3)	_V 3+	4)	Cr^{3+}
		sulphur sol is evaporated sulphu rmed. The sol is	r is ob	otained. On mixing with water sulphur sol
	1)	hydrophilic	2)	hydrophobic
	3)	reversible	4)	lyophilic
108. An a	-	halide reacts with alcoholic am	monia	a in a sealed tube, the product formed
	1)	a primary amine	2)	a secondary amine
	3)	a tertiary amine	4)	a mixture of all the three
109. Whe	en co	onc. H_2SO_4 is heated with P_2O_5	, the	acid is converted into
	1)	sulphur		
	2)	sulphur dioxide		
	3)	sulphur trioxide		
	4)	a mixture of sulphur dioxide as	nd sul	lphur trioxide
110. Ent	ropy.	of the universe is		
•	1)	continuously increasing	2)	continuously decreasing
•	3)	zero	4)	constant
		(Space for	Rough	ı Work)

		•	20			
111.	Which o	f the following salts o	on being dissolved	in water give	es pH > 7 at 25° C ?	
	1)	NH_4CN	2)	NH_4Cl		
	3)	KNO ₃	4)	KCN		
112.	The reas	gent used in Clemme	nson's reduction	is		
	1)	alc. KOH	2)	aq. KOH		
	3)	Zn-Hg / con. HCl	4)	Conc. H_2SO_4	•	
113.	When K	Br is dissolved in wa	ter, K^+ ions are			,
	1)	oxidised	2)	reduced		
	3)	hydrolysed	4)	hydrated		
114.	The nob		ed in a coconut bu	ılb at 173 <i>K</i> . 7	The gases that are not	
	1)	He and Ne	2)	Ar and Kr		
	3)	He and Xe	4)	Ne and Xe	·	
115.	The volu	me of $10N$ and $4NH$	${\it Cl}$ required to ma	ke 1 litre of 7	N HCl are	
	1)	0.75 litre of 10 <i>N HC</i>	l and 0.25 litre of	4N~HCl		
	2)	0.80 litre of 10 <i>N HC</i>	l and 0.20 litre of	4N~HCl		
	3)	0.60 litre of 10 <i>N HC</i>	$\it l$ and 0.40 litre of	4N~HCl		
	4)	0.50 litre of 10 <i>N HC</i>	l and 0.50 litre of	4N HCl		
			(Space for Dough	Worls)		

116.	A metal	present in insulin is	<u> </u>	Carlotte Contraction	
•	1)	copper	2)	iron	
	3)	zinc	4)	aluminium	
117.		forms two oxides which have differmains constant?	erei	nt compositions. The	equivalent mass of
	1)	carbon	2)	oxygen	
	3)	neither carbon nor oxygen	4)	both carbon and oxy	gen
118.	Maximu	m number of molecules of CH_3I th	at c	an react with a molec	ule of <i>CH</i> ₃ <i>NH</i> ₂ are
	1)	1	2)	2	4
	3)	4	4)	3	•
119.	Ellingha	um diagram represents a graph of			
	1)	$\Delta G \operatorname{Vs} T$	2)	$\Delta G^0 \mathrm{Vs} T$	
	3)	$\Delta S \operatorname{Vs} P$	4)	$\Delta G \operatorname{Vs} P$	
120.	Identify	the ore not containing iron			
	1)	chalcopyrites	2)	carnallite	
	3)	siderite	4)	limonite	
		(C		TTT 1\	

31

A - 1

TIS

COMMON ENTRANCE TEST - 2007

DATE	SUBJECT	TIME .
09 - 05 - 2007	BIOLOGY	04.00 PM to 05.20 PM

MAXIMUM MARKS	TOTAL DURATION	MAXIMUM TIME FOR ANSWERING
60	80 MINUTES	70 MINUTES

AILS	OKLET DETAI	N BOO	ESTIO	Qτ	R	YOUI	ION	ÆNT	I I
JMBER	SERIAL NUI	ODE	ION C	VERS	, ,	IBER	NUM	CET	
41	18424		. 1	Δ					
			- 1	A					

IMPORTANT INSTRUCTIONS TO CANDIDATES

(Candidates are advised to read the following instructions carefully, before answering on OMR answer sheet.)

- Ensure that you have entered your Name and Register Number of 2nd PUC Annual Examination / 12th Std. in the space provided on the OMR answer sheet.
- 2. Ensure that CET No. has been entered and shaded the respective circles on the OMR answer sheet.
- 3. ENSURE THAT THE TIMING, MARKS PRINTED ON THE OMR ANSWER SHEET ARE NOT DAMAGED / MUTILATED / SPOILED.
- 4. This Question Booklet is issued to you by the invigilator after the 2nd Bell. i.e., after 04.00 p.m.
- 5. Enter the Serial Number of this question booklet on the OMR answer sheet.
- 6. Carefully enter the Version Code of this question booklet on the OMR answer sheet and SHADE the respective circles completely.
- 7. As answer sheets are designed to suit the Optical Mark Reader (OMR) system, please take special care while filling and shading the CET NO. & Version Code of this question booklet.
- 8. DO NOT FORGET TO SIGN AT THE BOTTOM PORTION OF OMR ANSWER SHEET IN THE SPACE PROVIDED.
- 9. Until the 3rd Bell is rung at 04.10 p.m.:
 - Do not remove the seal present on the right hand side of this question booklet.
 - Do not look inside this question booklet.
 - Do not start answering on the OMR answer sheet.
- 10. After the 3rd Bell is rung at 04.10 p.m., remove the seal present on the right hand side of this question booklet and start answering on the OMR answer sheet.
- 11. This question booklet contains 60 questions and each question will have four different options / choices.
- 12. During the subsequent 70 minutes:
 - Read each question carefully.
 - Determine the correct answer from out of the four available options / choices given under each question.
 - Completely darken / shade the relevant circle with a BLUE OR BLACK INK BALLPOINT PEN against the question number on the OMR answer sheet.

CORRECT METHOD OF SHADING THE CIRCLE ON THE OMR SHEET IS AS SHOWN BELOW:



- 13. Please note that even a minute unintended ink dot on the OMR sheet will also be recognised and recorded by the scanner. Therefore, avoid multiple markings of any kind on the OMR answer sheet.
- 14. Use the space provided on each page of the question booklet for Rough work AND do not use the OMR answer sheet for the same.
- 15. After the last bell is rung at 05.20 p.m., stop writing on the OMR answer sheet and affix your LEFT HAND THUMB IMPRESSION on the OMR answer sheet as per the instructions.
- 16. Hand over the OMR ANSWER SHEET to the room invigilator as it is.
- 17. After separating and retaining the top sheet (CET Cell Copy), the invigilator will return the bottom sheet replica (Candidate's copy) to you to carry home for self-evaluation.
- 18. Preserve the replica of the OMR answer sheet for a minimum period of Oné year

· · · · · · · · · · · · · · · · · · ·	

1.	The terr	ns 'cytoplasm' and 'nucleoplasm' were given by
•	1)	Brown 2) Flemming
8	3)	Purkinje 4) Strasburger
2.	Which o	of the following experiment is called physiological demonstration of Osmosis?
,	1)	Potometer
	. 2)	Bell jar experiment
	3)	Thistle funnel - whose mouth is tied with egg membrane.
	4)	Thistle funnel - whose mouth is tied with parchment paper.
3.	The net	gain of ATP during glycolysis is
	1)	Two 2) Four
	3)	Six 4) Eight
4.	Coronar	y heart disease is due to
	1)	Weakening of the heart valves.
	2)	Insufficient blood supply to the heart muscles.
	3)	Streptococci bacteria.
	4)	Inflammation of pericardium.
5.	Manas s	sanctuary is located at
	. 1)	Bihar 2) Gujarath
	3)	Rajasthan 4) Assam

- 6. In which of the following organisms, self fertilization is seen?
 - 1) Earth worm

2) Liver fluke

3) Fish

4) Round worm


- 7. Rauwolfia serpentina belongs to family.
 - 1) Liliaceae

2) Fabaceae

3) Apocynaceae

4) Solanaceae

8.



$$2H_{2}O$$
 $\begin{cases} 4e^{-} \\ 4H^{+} + O_{2} \end{cases}$

In the above schematic diagram, which is plastocyanin?

1) A

2) B

3) C

- 4) D
- 9. In ABO blood groups, how many phenotypes are found?
 - 1) 1

2) 4

3) 6

- 4) 8
- 10. The tumor inducing capacity of <u>Agrobacterium</u> <u>tumaefaciens</u> is located in large extra chromosomal plasmids called
 - 1) pBR 322

2) Ti plasmid

3) Ri plasmid

4) lambda phage


- 11. Name the class of the Mycota, which is commonly called 'fungi imperfecti'.
 - 1) Zygomycota

2) Basidiomycota

3) Deuteromycota

- 4) Ascomycota
- 12. Which one is not correct about Krebs' cycle?
 - 1). It occurs in mitochondria.
 - 2) It starts with six carbon compound.
 - 3) It is also called citric acid cycle.
 - 4) The intermediate compound which links glycolysis with Krebs' cycle is malic acid.
- 13. Which would do maximum harm to a tree?
 - 1) Loss of all its bark.
- 2) Loss of half of its leaves.
- 3) Loss of half of its branches.
- 4) Loss of all of its leaves.

14.



RA - Right Auricle

RV - Right Ventricle

LA - Left Auricle

LV - Left Ventricle

In the above given diagram which blood vessel represents vena cava?

1) A

2) E

3) C


- 4) I
- 15. Rh ve person donated blood to Rh +ve person for the second time. Then,
 - 1) Rh +ve blood starts reacting to Rh -ve blood.
 - 2) Rh +ve person will die.
 - 3) Rh -ve person will die.
 - 4) Nothing happens to Rh +ve person.

16.	Checkin	ig of reradiating heat by at	mospheric dust O_3 , CO_2 and water vapours is –
	1)	Ozone layer effect	2) Radioactive effect
	3)	Green house effect	4) Solar effect
17.	Mutatio	n can not change	
	1)	Enzyme	2) DNA
	3)	RNA	4) Environment
18.	Liberati suitable	on of O_2 when green cells acceptor is called –	in water are exposed to sunlight in the presence of
	1)	Blackmann's reaction	2) Hill's reaction
	3)	Arnon's reaction	4) Emerson's enhance effect
19.	Guttatio	on is mainly due to	
	1)	Osmosis	2) Transpiration
*. 	3)	Root pressure	4) Imbibition
20.	• State	ment A : All Metatherian a	re placental mammals.
	• State:	ment B : All placental man	amals have menstrual cycle.
	1)	Both the statements A and	d B are true.
	2)	Both the statements A and	d B are false.
	3)	Statement A is true and S	tatment B is false.
	4)	Statement B is true and S	tatment A is false.
<u>·</u>	· · · · · · · · · · · · · · · · · · ·		

21.	Population density of terrestrial organisms is measured in terms of individual per
	1) Meter 2) Meter ²
	3) Meter ³ 4) Meter ⁴
22.	Nitrogenous waste products are eliminated mainly as -
	1) urea in tadpole as well as in adult frog.
	2) urea in tadpole and ammonia in adult frog.
	3) urea in tadpole and uric acid in adult frog.
	4) urea in adult frog and ammonia in tadpole.
23.	In man, the blue eye colour is recessive to the brown eye colour. If the boy has brown eye and his mother is blue eyed, what would be the phenotype of his father?
,	1) Green eye 2) Blue eye
	3) Black eye 4) Brown eye
24.	Munch hypothesis is based on
	1) Translocation of food due to Turgor Pressure (TP) gradient.
. •	2) Translocation of food due to imbibition force.
	3) Translocation of food due to TP gradient and imbibition force.
	4) None of these
25.	Interferons are
_	1) Complex protein 2) Anti-clotting protein.
	3) Anti-bacterial protein 4) Anti-viral protein.
	(Space for Rough Work)

In the diagram of lenticel

identify the parts as A, B, C, D.



- 1) A-Complementary cells, B- Phellogen, C- Phelloderm, D- Periderm
- 2) A- Complementary cells, B- Phellum, C- Periderm, D- Phelloderm
- 3) A- Phellum, B- Periderm, C- Phellogen, D- Phelloderm
- 4) A- Phellum, B- Complementary cells, C- Phellogen, D- Phelloderm

Sterlization of tissue culture medium is done by -

- 1) Mixing the medium with antifungal agents.
- 2) Keeping the medium at -20° C.
- Autoclaving of medium at 120°C for 15 minutes.
- Filtering the medium through fine sieve.

Match the following:

- <u>Leishmania</u> <u>dorovani</u>
- В. <u>Wuchereria</u> – <u>bancrofti</u>
- C. <u>Trypanosoma</u> – <u>gambiense</u>
- D. Entamoeba - histolytica

- p. Malaria
- g. Amoebiosis.
- r. Kala azar
- s. Sleeping sickness
- t. Filariasis
- $\begin{bmatrix} 1 \end{bmatrix}$ A-r B-t C-s D-q = 2) A-r B-t C-q D-p
- 3) A-s B-r C-q D-p 4) A-r B-s C-t D-t

29. The idea of Natural selection as the fundamental process of evolutionary changes was reached

- Independently by Charles Darwin and Alfred Russel Wallace in 1900
- 2) By Charles Darwin in 1866.
- 3) By Alfred Russel Wallace in 1901.
- Independently by Charles Darwin and Alfred Russel Wallace in 1859.

30. Auxins originates at the tip of the stem and controls growth elsewhere. The movement of auxins is largerly

- 1) Acropetal and basipetal
- Centropetal

3) Basipetal

4) Acropetal

31.	If a lengtake?	th of DNA has 45,000 ba	ase pairs, ho	ow m	nany complete trans will the DNA molecule
	1) .	45		2)	450
	3)	4,500		4)	45,000
32.		cess in which mature lus is called	differentia	ted	cells reverse to meristematic activity to
,	1)	Cyto differentiation		2)	Redifferentiation
	3)	Dedifferentiation		4)	Differentiation
33.	The late	eral roots originate from	m		
	1)	Epiblema		2)	Cortical cells below root hairs
. ' ·	3)	Endoderm cells		4)	Pericycle cells
34.	Which a	ccessory genital gland	occurs only	in r	nammalian male ?
	1)	Cowper's gland	2	. 2)	Bartholian gland
	3)	Prostate gland		4)	Perineal gland
35.	When th	ne concentration of the	soil solutes	s is l	ow, the absorption of water is
	1)	Increased		2)	Decreased
	3)	Remain normal		4)	Stopped
	٠		1		

36 .	Edaphology is	
	1) Study of Snakes	2) Study of Amphibians
	3) Study of Elephants	4) None of these
37.	Pineal gland of human brain secretes r	melatonin concerned with
	1) Colouration of skin	2) Sleep
	3) Anger	4) Body temperature
38.		(TTRR) crossed with a dwarf plant with wrinkle is of tall plants with round seeds. What would be rinkle seeds in \mathbf{F}_1 generation?
	1) 0 3) $\frac{1}{4}$	2) $\frac{1}{2}$ 4) $\frac{1}{16}$
39	Cell wall consists of	, 10
	1) Lignin, hemi cellulose, pectin	n and lipid
	2) Lignin, hemi cellulose, pectin	n and cellulose
	3) Lignin hemi cellulose, protein	n and lipid
	4) Hemi cellulose, cellulose, tubu	oulin and lignin.
40.	The post and tail is present in -	
	1) Invertebrates	2) Vertebrates
	3) Chordates	4) In all of them

A -1

- 41. Synthesis of food in C_4 pathway occurs in Chlorophyll of
 - 1) Spongy mesophyll
- 2) Palisade cells

11

3) Guard cells

- 4) Bundle sheath
- 42. The sequence of structural gene in lac operon concept is
 - , 1) lac Y, lac Z, lac A

2) lac Z, lac Y, lac A


3) lac A, lac Y, lac Z

- 4) lac A, lac Z, lac Y
- 43. Pericarp and placentae are edible part of simple fleshy berry fruit
 - 1) Tomato

· 2) Date palm

3) Jack fruit

- 4) Banana
- 44. In the diagram, which of the following processes are shown in Amoeba?



Molecules in solution

1) Phagocytosis

2) Pinocytosis

3) Exocytosis

- 4) All of these
- 45. An essential element is that which
 - 1) is found in plant ash.
 - 2) is available in the soil.
 - 3) improve health of the plant.
 - 4) is irreplaceable and indispensable for growth of plants.

- 46. Nucleic acid occurs in
 - 1) Cytoplasm

2) Mitochondria and chloroplast

3) Golgibody


- 4) Lysosomes.
- 47. The number of mitotic cell division required to produce 256 cells from single cell would be
 - 1) 6

2)

3) 10

- 4) 12
- 48. The central dogma of protein synthesis in teminious is
 - 1) $DNA \rightarrow DNA \rightarrow m-RNA \rightarrow Protein$
 - 2) $m-RNA \rightarrow g.RNA \rightarrow DNA \rightarrow Protein$
 - 3) $g.RNA \rightarrow DNA \rightarrow m-RNA \rightarrow Protein$
 - 4) $DNA \rightarrow G-RNA \rightarrow m-RNA \rightarrow Protein$
- 49. In tissue culture roots can be induced by
 - 1) No cytokinin and only auxins.
 - 2) Higher concentration cytokinin and lower concentration auxins.
 - 3) Lower concentration of cytokinin and higher concentration of auxins.
 - 4) Only cytokinin and no auxins.

50.



- 1) A- uterus, B- uterine cavity, C- oviducal funnel, D- ovary
- 2) A-cervix, B- uterine cavity, C- fallopian tube, D- ovary
- 3) A- oviduct, B- uterus, C- outduct, D- ovary
- 4) A-cervix, B-uterus, C-ovary, D-tumour

51.	•	environment for germination is		the seed coat when a seed is placed in
٠.	1)	Absorption	2)	Imbibition
	3)	Osmosis	4)	Active transport
52.		is a taxon, which is likely to mons	ve in	to endangered category in near future, if
	1)	Rare	2)	Extinct
	3)	Vulnerable	4)	Endanger
53.		ed inflammatory response appears d heat due to certain chemical, th		ne site of infection causes redness, swelling, re
,	1).	Histamin and cerumen	2)	Prostaglandins and cerumen
	3)	Histamin and prostaglandins	4)	Cerumen and mucus.
54.	Non ker	atimised stratified epithelium oc	curs	in
•	1)	Vagina and cervix	2)	Buccal cavity and anus
	3)	Vagina, cervix and buccal cavity	4)	Vagina, cervix, buccal cavity and anus
55.	Succus	entericus is secreted by ?	àr.	
•	· 1)	Crypts of Leiberkuhn	2)	Brunner's gland
	3)	Both (1) and (2)	4)	None of these
	•			

- 56. Residual volume is
 - 1) Greater than vital capacity
- 2) Greater than fidal volume.
- 3) Lesser than tidal volume.
- 4) Greater than inspiratory volume.

- **57.** Find the odd example.
 - 1) Sea cucumber


2) Sea urchin

3) Sea lily

4) Sea fan

14

- 58. Which one is correct?
 - 1) Neuron = Cyton + Dendrite + Axon + Synapse
 - 2) Lymph = Plasma + RBC + WBC
 - 3) Blood = Plasma + RBC + WBC + Blood platelets
 - 4) Plasma = Blood lymphocytes
- 59. In the given diagram name the parts A, B, C and D.



- 1) A- Intine, B- Exine, C- Germ pore, D- Generative cell, E- Vegetative cell
- 2) A- Exine, B- Intine, C- Vegetative cell, D- Germ pore, E- Generative cell
- 3) A- Germ pore, B- Generative cell, C- Intine, D- Exine, E- Vegetative cell
- 4) A- Germ pore, B- Generative cell, C- Exine, D- Intine, E- Vegetative cell
- 60. The largest RBC's have been seen in
 - 1) Amphibia

2) Man


3) Elephant

4) Whale

15 A -1

(Space for Rough Work)

SR - 1 Turn Over



€ -