Page 1 of 10

RUHS

Medical college Recruitment Exam Nurse Grade II

	-
Q. No. 1 0131001	The innermost layer of the heart is
Option 1	Epicardium
Option 2	Endocardium
Option 3	Myocardium
Option 4	Pericardium
Correct Option	2
Q. No. 2 0131002	What is a sociological definition of the word drug ?
Option 1	A substance that changes mood, behavior or consciousness.
Option 2	A dysfunction
Option 3	A chemical
Option 4	Ecology
Correct Option	1
	·
O No 3	

Q. No. 3 0131003	Hypnotherapy is least effective on people suffering from
Option 1	Anxiety
Option 2	Hypertension
Option 3	Pain
Option 4	Schizophrenia
Correct Option	4

Q. No. 4 0131004	Which of the following stain is used to detect Mycobacterium species ?
Option 1	Gram stain
Option 2	Capsule stain
Option 3	Acid fast stain
Option 4	Endospore stain
Correct Option	3

Q. No. 5 0131005	The best method of sterilization of dressing powder is
Option 1	Autoclaving
Option 2	Hot air oven
Option 3	Inspissation
Option 4	Chemical method
Correct Option	2

-	Which of the following results of the lab test is indicative of the bacterial infection ?

Option 1	RBC : 4.0million/cmm
Option 2	Hemoglobin : 13.2gm/dl
Option 3	Platelets : 350,000/cmm
Option 4	WBC : 15,000/ cmm
Correct Option	4

Q. No. 7 0131007	If after surgery the patient's abdomen becomes distended and no bowel sounds appreciated, what would be the most suspected complication?
Option 1	Intussusception
Option 2	Paralytic Ileus
Option 3	Hemorrhage
Option 4	Ruptured colon
Correct Option	2

Q. No. 8 0131008	A registered nurse, witnessed an old woman hit by a motorcycle while crossing a train railway. The old woman fell at the railway. She was rushed at the scene. As a registered nurse, the first thing that that needs to be done is
Option 1	Stay with the person, encourage her to remain still and immobilize the leg while waiting for the ambulance
Option 2	Leave the person for a few moments to call for help
Option 3	Reduce the fracture manually
Option 4	Move the person to a safer place
Correct Option	4

Q. No. 9 0131009	Good source of vitamin D include all except
Option 1	blueberries
Option 2	sunlight
Option 3	salmon, tuna sardines and mackerel
Option 4	fortified milk and other dairy products
Correct Option	2

Q. No. 10 0131010	A 48-year-old woman presents with intermenstrual bleeding for two months and episodes of bleeding occurring any time in the cycle. There is no associated pain. Differential diagnosis for intermenstrual bleeding does not include
Option 1	endocervical polyp
Option 2	ovarian teratoma
Option 3	cervical malignancy
Option 4	endometrial polyp
Correct Option	2

Q. No. 11 0131011	Following are the complications of premature menopause except
Option 1	osteoporosis
Option 2	infertility

Option 3	decreased cardiovascular risk
Option 4	vasomotor symptoms
Correct Option	3
r	
Q. No. 12 0131012	Which nursing diagnosis should receive priority in the plan of care for a three- month-old infant who has a salmonella infection ?
Option 1	Weight loss
Option 2	Fluid volume deficit
Option 3	Sleepiness
Option 4	Shouting and Crying
Correct Option	2

Q. No. 13 0131013	Which of the following is a non-communicable disease ?
Option 1	Diabetes Mellitus
Option 2	Tuberculosis
Option 3	Typhoid
Option 4	Small Pox
Correct Option	1

Q. No. 14 0131014	New cases in population during a fixed period is determined by
Option 1	Prevalence Ratio
Option 2	Attack Rate
Option 3	Disability Rates
Option 4	Incidences Rates
Correct Option	4

Q. No. 15 0131015	Factors affecting parenting behaviour does not include a parent's own
Option 1	personal experiences and trauma
Option 2	mental health
Option 3	family experiences
Option 4	dietary habits
Correct Option	4

Q. No. 16 0131016	A 32-year-old woman is presented to the gynaecology clinic with infrequent periods. A hormone profile is done and all of the following are consistent with polycystic ovarian syndrome except
Option 1	increased androgen levels
Option 2	normal FSH
Option 3	normal oestradiol
Option 4	decreased LH
Correct Option	4

Q. No. 17	
Q. NO. 17 0131017	Which drug is responsible for "red men syndrome"?
Option 1	Vancomycin
Option 2	Linezolid
Option 3	Chloramphenicol
Option 4	Cefadroxyl
Correct Option	1
Q. No. 18 0131018	A male patient's daughter telephones an adult-gerontology primary care nurse practitioner to inquire about the patient's illness. The daughter reports that her father relies upon her to explain everything to him. The nurse practitioner's initial response is to
Option 1	ask the daughter to provide a copy of the patient's advance directive document.
Option 2	tell the daughter that the nurse practitioner can discuss the information only with the patient.
Option 3	assure the daughter that the nurse practitioner can disclose the requested information.
Option 4	inform the daughter that she must come to the clinic to discuss her father's case.
Correct Option	2
Q. No. 19 0131019	Which one of these is used in bed wetting in children ?
Option 1	Imipramine
Option 2	Chlorpromazine
Option 3	Levodopa
Space 5	
Option 4	Tacrine
Option 4	Tacrine
Option 4 Correct Option Q. No. 20	Tacrine 1 What is the most common site of rupture of a cerebral aneurysm with resulting
Option 4 Correct Option Q. No. 20 0131020	Tacrine 1 What is the most common site of rupture of a cerebral aneurysm with resulting subarachnoid hemorrhage (SAH) ?
Option 4 Correct Option Q. No. 20 0131020 Option 1	Tacrine 1 What is the most common site of rupture of a cerebral aneurysm with resulting subarachnoid hemorrhage (SAH) ? Anterior cerebral artery
Option 4 Correct Option Q. No. 20 0131020 Option 1 Option 2	Tacrine 1 What is the most common site of rupture of a cerebral aneurysm with resulting subarachnoid hemorrhage (SAH) ? Anterior cerebral artery Middle cerebral artery
Option 4 Correct Option Q. No. 20 0131020 Option 1 Option 2 Option 3	Tacrine 1 What is the most common site of rupture of a cerebral aneurysm with resulting subarachnoid hemorrhage (SAH) ? Anterior cerebral artery Middle cerebral artery Anterior communicating artery
Option 4 Correct Option Q. No. 20 0131020 Option 1 Option 2 Option 3 Option 4	Tacrine 1 What is the most common site of rupture of a cerebral aneurysm with resulting subarachnoid hemorrhage (SAH) ? Anterior cerebral artery Middle cerebral artery Anterior communicating artery Posterior communicating artery
Option 4 Correct Option Q. No. 20 0131020 Option 1 Option 2 Option 3 Option 4 Correct Option Q. No. 21	Tacrine 1 What is the most common site of rupture of a cerebral aneurysm with resulting subarachnoid hemorrhage (SAH) ? Anterior cerebral artery Middle cerebral artery Middle cerebral artery Anterior communicating artery Posterior communicating artery 3
Option 4 Correct Option Q. No. 20 0131020 Option 1 Option 2 Option 3 Option 4 Correct Option Q. No. 21 0131021	Tacrine 1 What is the most common site of rupture of a cerebral aneurysm with resulting subarachnoid hemorrhage (SAH) ? Anterior cerebral artery Middle cerebral artery Middle cerebral artery Posterior communicating artery Posterior communicating artery 3 Which of these used in multidrug resistant tuberculosis ?
Option 4 Correct Option Q. No. 20 0131020 Option 1 Option 2 Option 3 Option 4 Correct Option Q. No. 21 0131021 Option 1	Tacrine 1 What is the most common site of rupture of a cerebral aneurysm with resulting subarachnoid hemorrhage (SAH) ? Anterior cerebral artery Middle cerebral artery Middle cerebral artery Anterior communicating artery Posterior communicating artery 3 Which of these used in multidrug resistant tuberculosis ? Ciprofloxacin
Option 4 Correct Option Q. No. 20 0131020 Option 1 Option 2 Option 3 Option 4 Correct Option Q. No. 21 0131021 Option 1 Option 2	Tacrine 1 What is the most common site of rupture of a cerebral aneurysm with resulting subarachnoid hemorrhage (SAH) ? Anterior cerebral artery Middle cerebral artery Middle cerebral artery Anterior communicating artery Posterior communicating artery 3 Which of these used in multidrug resistant tuberculosis ? Ciprofloxacin Penicillin
Option 4 Correct Option Q. No. 20 0131020 Option 1 Option 2 Option 3 Option 4 Correct Option Q. No. 21 0131021 Option 1 Option 2 Option 2 Option 3	Tacrine 1 What is the most common site of rupture of a cerebral aneurysm with resulting subarachnoid hemorrhage (SAH) ? Anterior cerebral artery Middle cerebral artery Middle cerebral artery Anterior communicating artery Posterior communicating artery 3 Which of these used in multidrug resistant tuberculosis ? Ciprofloxacin Penicillin Metronidazole
Option 4 Correct Option Q. No. 20 0131020 Option 1 Option 2 Option 3 Option 4 Correct Option Q. No. 21 0131021 Option 1 Option 1 Option 2 Option 3 Option 3 Option 3 Option 4	Tacrine 1 What is the most common site of rupture of a cerebral aneurysm with resulting subarachnoid hemorrhage (SAH) ? Anterior cerebral artery Middle cerebral artery Middle cerebral artery Anterior communicating artery Posterior communicating artery 3 Which of these used in multidrug resistant tuberculosis ? Ciprofloxacin Penicillin Metronidazole Linezolide

Option 2	Pleural effusion
Option 3	Flail chest
Option 4	Pneumothorax
Correct Option	4

Q. No. 23 0131023	A patient comes to the clinic with reports of acute pain and loss of vision in the right eye. The adult-gerontology primary care nurse practitioner informs the patient that these symptoms require
Option 1	immediate referral to an ophthalmologist.
Option 2	assessment with a split lamp.
Option 3	examination with an otoscope.
Option 4	treatment with several ophthalmologic agents.
Correct Option	1

Q. No. 24 0131024	Which of the following blood group is universal recipient?
Option 1	A
Option 2	В
Option 3	АВ
Option 4	0
Correct Option	3

Q. No. 25 0131025	Deficiency of which of the following vitamin during pregnance may cause neural tube defects ?
Option 1	Folic acid
Option 2	Riboflavin
Option 3	Niacin
Option 4	Thiamine
Correct Option	1

Q. No. 26 0131026	During the initial care of rape victims the following are to be considered by a nurse except
Option 1	maintain a non-judgmental approach.
Option 2	assure privacy
Option 3	touch the client to show acceptance and empathy.
Option 4	accompany the client in the examination room.
Correct Option	3

Q. No. 27 0131027	Shick test is carried out to determine
Option 1	polio
Option 2	pertusis
Option 3	meningitis
Option 4	diphtheria
Correct Option	4

Q. No. 28 0131028	Which of the following is most common side effect of aspirin ?
Option 1	Gastrointestinal bleeding
Option 2	Constipation
Option 3	Nystagmus
Option 4	Urinary incontinence
Correct Option	1
Q. No. 29 0131029	The intestinal juice secreted in the intestine containing digestive enzymes is known as
Option 1	Succusentericus
Option 2	Juicy fluid
Option 3	Succus juice
Option 4	Chyme
Correct Option	1
Q. No. 30 0131030	All are the non-modifiable risk factors for atherosclerosis except
Option 1	high cholesterol intake
Option 2	age
Option 3	heredity
Option 4	gender
Correct Option	1
Q. No. 31 0131031	The first drug of choice for a patient admitted with status epilepticus is
Option 1	phenytoin
Option 2	carbamezepine
Option 3	diazepam
Option 4	sodium valproate
Correct Option	3
Q. No. 32	Which of the following is not a saturated fatty acid ?
0131032	Oleic acid
0131032 Option 1	Oleic acid Palmic acid
0131032 Option 1 Option 2	
0131032 Option 1 Option 2 Option 3 Option 4	Palmic acid
0131032 Option 1 Option 2 Option 3 Option 4	Palmic acid Lauric acid
0131032 Option 1 Option 2 Option 3	Palmic acid Lauric acid Stearic acid

Г

Τ

Option 2	topical hormonal cream.
Option 3	topical antibiotic cream.
Option 4	topical antifungal cream.
Correct Option	2

Q. No. 34 0131034	Which of the following is antibody occurs as pentameric structure ?
Option 1	IgA
Option 2	IgG
Option 3	IgE
Option 4	IgM
Correct Option	4

Q. No. 35 0131035	Which of the following is not typical anti-psychotic and does not have marked extra pyramidal side effects ?
Option 1	Chlorpromazine
Option 2	Fluphenazine
Option 3	Clozapine
Option 4	Haloperidol
Correct Option	3

Q. No. 36 0131036	In case of heparin over dose, which agent should be used ?
Option 1	Diltiazem
Option 2	Protamine
Option 3	Imferon
Option 4	Warfarin
Correct Option	2

Q. No. 37 0131037	In case of development of urticaria one-half hour after the blood transfusion has begun, the first action by the nurse should be
Option 1	to slow the rate of infusion
Option 2	to administer anti-histaminic agent
Option 3	to note vital signs
Option 4	to stop the infusion
Correct Option	4

Q. No. 38 0131038	When haemoglobin combines with oxygen, it is known as
Option 1	oxyhemoglobin
Option 2	deoxyhemoglobin
Option 3	reduced hemoglobin
Option 4	oxidized haemoglobin
Correct Option	1

Q. No. 39 0131039	Which of the following monoclonal antibody is approved for the treatment of asthma ?
Option 1	Bevacizumab
Option 2	Omalizumab
Option 3	Cetuximab
Option 4	Trastuzumab
Correct Option	2
Q. No. 40 0131040	A 55-year-old man was admitted after a motor vehicle accident with multiple injuries. His friend reported that he had been seeing a psychiatrist for an unknown diagnosis. He stated that the patient sometimes gets "hyper" for no reason, and becomes violent and sometimes is depressed. Of the following, which general psychiatric disorder is characterized by a pattern of aggression or violence which includes irritability, agitation, and violent behavior during manic or psychotic episodes ?
Option 1	Schizophrenia
Option 2	Bipolar disorder
Option 3	Delusional disorder
Option 4	Depression
Correct Option	2
Q. No. 41 0131041	Which of the following is not filtered out during glomerular filtration ?
Option 1	Glucose
Option 2	Water
Option 3	Salts
Option 4	Proteins
Correct Option	4
Q. No. 42 0131042	For a 6 month pregnant lady, which of the following are contraindicated ?
Option 1	Retinol
Option 2	Tetanus toxoid
Option 3	Folic acid
Option 4	Ferrous sulphate
Correct Option	1
Q. No. 43 0131043	Recommended site of administration of DPT in infants is
Option 1	deltoid
Option 2	gluteal
Option 3	forearm
Option 4	intravenous
Correct Option	1
Q. No. 44 0131044	Spironolactone produces

Option 1	hypernatremia
Option 2	hyperkalaemia
Option 3	hyponatremia
Option 4	hyprkalaemia
Correct Option	2

Q. No. 45 0131045	The agent contraindicated with warfarin treatment is
Option 1	aspirin
Option 2	ranitidine
Option 3	codeine
Option 4	isaphgol
Correct Option	1

Q. No. 46 0131046	After obtaining 3 L of fluid from a client via parecentesis, the nurse would be alert for which complication ?
Option 1	Encephalopathy
Option 2	Bleeding from site
Option 3	Vascular collapse
Option 4	Respiratory distress
Correct Option	3

Q. No. 47 0131047	Which of the following is the most prominent feature of public health nursing ?
Option 1	Public health nursing focuses on preventive, not curative, services.
Option 2	Services are provided free of charge to people within the catchment area.
Option 3	The public health nurse functions as part of a team providing a public health nursing services.
Option 4	It involves providing home care to sick people who are not confined in the hospital
Correct Option	1

Q. No. 48 0131048	The lung function tests are determined by
Option 1	tonometer
Option 2	spirometer
Option 3	sphygmomanometer
Option 4	barometer
Correct Option	2

Q. No. 49 0131049	Which of the following part of eye contains rhodopsin pigment ?
Option 1	Rods
Option 2	Cones
Option 3	Lens
Option 4	Iris

Correct Option	1
Q. No. 50 0131050	Which of the following agent causes discolouration of teeth if taken with milk ?
Option 1	Tetracycline
Option 2	Penicillin
Option 3	Metronidazole
Option 4	Ciprofloxacin
Correct Option	1