

Question Paper of MP MCA - 2014

This Test was MP Pre - MCA 2014 (Set - A)

1. The multiplication inverse of the matrix :

$$A = \begin{bmatrix} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

is :

(a) $\begin{bmatrix} -\cos \theta & \sin \theta & 0 \\ -\sin \theta & -\cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix}$

(c) $\begin{bmatrix} -\cos \theta & -\sin \theta & 0 \\ \sin \theta & -\cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix}$



(b) $\begin{bmatrix} \cos \theta & \sin \theta & 0 \\ -\sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix}$

(d) $\begin{bmatrix} \cos \theta & \sin \theta & 0 \\ \sin \theta & -\cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix}$

2. The equations :

$$x + 2y + 3z = 1$$

$$x - y + 4z = 0$$

$$2x + y + 7z = 1$$

have :

(a) Only one solution

(c) No solutions

(b) Only two solutions

(d) Infinitely many solutions

3. For positive number x, y, z the value of the determinant :

$$\begin{vmatrix} 1 & \log_x y & \log_x z \\ \log_y x & 1 & \log_y x \\ -\log_z x & \log_z y & 1 \end{vmatrix} \text{ is :}$$


(a) 0

(b) 1

(c) -1

(d) $\log 3$

4. The centre of a circle passing through the points $(0, 0)$, $(1, 0)$ and touching the circle $x^2 + y^2 = 9$ is :
- (a) $\left(\frac{3}{2}, \frac{1}{2}\right)$ (b) $\left(\frac{1}{2}, -\sqrt{2}\right)$ (c) $\left(\frac{1}{2}, \frac{3}{2}\right)$ (d) $\left(\frac{1}{2}, \frac{1}{2}\right)$
5. Two perpendicular tangents to the circle $x^2 + y^2 = a^2$ meet at a point P. Then the locus of point P has the equation :
- (a) $x^2 + y^2 = a^2$ (b) $x^2 + y^2 = 2a^2$ (c) $x^2 + y^2 = 3a^2$ (d) $x^2 + y^2 = 4a^2$
6. One of the limit point of the coaxial system of a circles containing $x^2 + y^2 - 6x - 6y + 4 = 0$, $x^2 + y^2 - 2x - 4y + 3 = 0$ is :
- (a) $(1, -1)$ (b) $(-1, 1)$ (c) $(-1, 2)$ (d) $(1, -2)$
7. The locus of the centre of circle which cuts the circles $x^2 + y^2 + 2g_i x + 2f_i y + c_i = 0$, $(i = 1, 2)$ orthogonally is
- (a) an ellipse (b) another circle
(c) the radical axis of given circles (d) a conic
8. The locus of the centre of a circle of radius 2 unit which rolls on the outside of the circle $x^2 + y^2 + 3x - 6y - 9 = 0$ is :
- (a) $x^2 + y^2 + 3x - 6y + 5 = 0$ (b) $x^2 + y^2 + 6x - 3y + 24 = 0$
(c) $x^2 + y^2 + 3x - 6y - 31 = 0$ (d) $x^2 + y^2 + 3x + 6y + 41 = 0$
9. The mid-point of the chord intercepted on the line $4x - 3y + 4 = 0$ by the parabola $y^2 = 8x$, is :
- (a) $(5, 3)$ (b) $\left(5, \frac{3}{2}\right)$ (c) $\left(\frac{5}{2}, 3\right)$ (d) $\left(\frac{5}{4}, 3\right)$
10. The polar of a point with respect to $y^2 = 4ax$ touches $x^2 = 4by$, then the locus of this point is .
- (a) a circle (b) a parabola (c) an ellipse (d) a rectangular hyperbola
11. The line $x + y = 6$ is a normal to the parabola $y^2 = 8x$ at the point
- (a) $(18, -12)$ (b) $(4, 2)$ (c) $(2, 4)$ (d) $(3, 3)$
12. The length of the major axis of an ellipse is three time the length of its minor axis, its eccentricity is :
- (a) $\frac{2\sqrt{2}}{3}$ (b) $\frac{1}{3}$ (c) $\frac{1}{\sqrt{3}}$ (d) $\frac{1}{\sqrt{2}}$
13. If CP and CD are the semi-conjugate diameter of the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, then $CP^2 + CD^2 =$
- (a) $a + b$ (b) $a^2 + b^2$ (c) $a^2 - b^2$ (d) $\sqrt{a^2 + b^2}$



14. The ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ and the straight line $y = mx + c$ intersect in real points only, if :
- (a) $a^2m^2 < c^2 - b^2$ (b) $a^2m^2 > c^2 - b^2$ (c) $a^2m^2 \geq c^2 - b^2$ (d) $c \geq b$
15. The curve represented by $x = a(\cosh\theta + \sinh\theta)$, $y = b(\cosh\theta - \sinh\theta)$ is :
- (a) a hyperbola (b) an ellipse (c) a parabola (d) a circle
16. The foci of the ellipse $\frac{x^2}{16} + \frac{y^2}{b^2} = 1 (b < 4)$ and the hyperbola $\frac{x^2}{144} - \frac{y^2}{81} = \frac{1}{25}$ coincide, then the value of b^2 is
- (a) 1 (b) 5 (c) 7 (d) 9
17. The equation of a tangent parallel to $y = x$ drawn to $\frac{x^2}{3} - \frac{y^2}{2} = 1$ is :
- (a) $x - y + 1 = 0$ (b) $x - y + 2 = 0$ (c) $x + y - 1 = 0$ (d) $x + y + 2 = 0$
18. The equation of ellipse whose one focus is at $(4, 0)$ and whose eccentricity is $\frac{4}{5}$, is
- (a) $\frac{x^2}{9} + \frac{y^2}{25} = 1$ (b) $\frac{x^2}{25} + \frac{y^2}{9} = 1$ (c) $\frac{x^2}{25} + \frac{y^2}{16} = 1$ (d) $\frac{x^2}{16} + \frac{y^2}{25} = 1$
19. If $f(x) = \log |\log x|$, then $f'(e)$ is equal to :
- (a) e (b) $-e$ (c) e^2 (d) e^{-1}
20. If $y = x + e^x$, then the value of $\frac{d^2x}{dy^2}$ is :
- (a) e^x (b) $-e^x(1 + e^x)^{-3}$ (c) $-e^x(1 + e^x)^{-2}$ (d) $(1 + e^x)^{-2}$
21. If $x\sqrt{1+y} + y\sqrt{1+x} = 0$ then $\frac{dy}{dx}$ is equal to :
- (a) $\frac{1}{(1+x)^2}$ (b) $\frac{1}{(1+x^2)}$ (c) $-\frac{1}{(1+x)^2}$ (d) $-\frac{1}{(1+x^2)}$
22. The differentiation of $\log \sqrt{\frac{1+\sin x}{1-\sin x}}$ with respect to x is :
- (a) $\cos x$ (b) $\log \cos x$ (c) $\frac{1}{2}(1 + \sin x)$ (d) $\sec x$

23. If $x = a \left(\cos t + \log \tan \frac{t}{2} \right)$, $y = a \sin t$, then $\frac{dy}{dx}$ is equal to :
- (a) $\cos t$ (b) $\log \tan t$ (c) $\operatorname{cosec} t$ (d) $\tan t$
24. The angle of intersection of the curves $y = x^2$ and $y = 7 - \frac{1}{6}x^3$ at $(1, 1)$ is :
- (a) $\frac{\pi}{4}$ (b) $\frac{\pi}{3}$ (c) $\frac{\pi}{2}$ (d) $\frac{\pi}{6}$
25. The curve $y - e^{xy} + x = 0$ has a vertical tangent at the point :
- (a) $(1, 1)$ (b) no point (c) $(0, 1)$ (d) $(1, 0)$
26. The minimum value of the function $y = x^3 - 12x$, on the interval $0 \leq x \leq 3$, is :
- (a) 0 (b) -9 (c) -16 (d) -19
27. If $u = x^2y + y^2z + z^2x$, then the value of $\frac{\partial u}{\partial x} + \frac{\partial u}{\partial y} + \frac{\partial u}{\partial z}$ is equal to :
- (a) $x + y + z$ (b) $(x + y + z)^2$ (c) $(x + y + z)^3$ (d) $x^2y^2z^2$
28. The value of $\int e^x \left(\frac{1}{x} - \frac{1}{x^2} \right) dx$ is equal to :
- (a) $\frac{e^x}{x} + c$ (b) $\frac{e^x}{x^2} + c$ (c) $e^x \left(1 + \frac{1}{x} \right) + c$ (d) $e^x \log \left(1 - \frac{1}{x} \right) + c$
29. $\int_b^1 \frac{\tan^{-1} x}{1+x^2} dx$ is equal to :
- (a) $\frac{\pi}{4}$ (b) $\frac{\pi^2}{16}$ (c) $\frac{\pi^3}{32}$ (d) $\frac{\pi^2}{18}$
30. Area lying in the first quadrant and bounded by the circle $x^2 + y^2 = 4$, the line $x = \sqrt{3}y$ and x-axis is
- (a) π (b) $\frac{\pi}{2}$ (c) $\frac{\pi}{3}$ (d) $\frac{\pi}{4}$
31. The value of $\int_b^1 \frac{dx}{1+x}$ by the trapezoidal rule, taking $n = 2$, is equal to :
- (a) $\frac{7}{24}$ (b) $\frac{11}{24}$ (c) $\frac{13}{24}$ (d) $\frac{17}{24}$

32. In Simpson $\frac{1}{3}$ rule, we approximate the integrand over every pairs of adjacent subintervals by curves of degree

- (a) 0 (b) 1 (c) 2 (d) 3

33. If $u = \sin^{-1}\left(\frac{\sqrt{x}-\sqrt{y}}{\sqrt{x}+\sqrt{y}}\right)$, then u_x is equal to :

- (a) $-\frac{x}{y}u_y$ (b) $-\frac{y}{x}u_y$ (c) $\frac{x}{y}u_y$ (d) $\frac{y}{x}u_y$

34. The solution of $\frac{dy}{dx} = \frac{3e^{2x} + 3e^{4x}}{e^x + e^{-x}}$ is given by :

- (a) $y = e^{4x} + x^2 + c$ (b) $ye^x = \frac{x^3}{3} + c$ (c) $y = e^{3x} + c$ (d) $y = e^{-3x} + 6x + c$

35. The solution of $\frac{dy}{dx} = \cos(x+y)$ is :

- (a) $\log\left|1 + \tan\left(\frac{x+y}{2}\right)\right| = x + c$ (b) $\tan\left(\frac{x+y}{2}\right) = x + c$
(c) $y = \sin(x+y) + c$ (d) $\tan(y+x) = x + \sec x + c$



36. The differential equation of all non-vertical lines in a plane ($ax + by = 1, b \neq 0$) is :

- (a) $\frac{dy}{dx} = 0$ (b) $\frac{dx}{dy} = 0$ (c) $\frac{d^2x}{dy^2} = 0$ (d) $\frac{d^2y}{dx^2} = 0$

37. The solution of $(x+y) dx + xdy = 0$ is given by :

- (a) $x^2 + y^2 = c$ (b) $2x^2 - y^2 = c$ (c) $x^2 + 2xy = c$ (d) $y^2 + 2xy = c$

38. The solution of differential equation

$$y \frac{dy}{dx} = x - 1, \text{ satisfying } y(1) = 1 \text{ is :}$$

- (a) $y^2 = x^2 - 2x + 2$ (b) $y^2 = 2x^2 - x - 1$ (c) $y = x^2 - 2x + 2$ (d) $y^2 = x + 2$

39. The integrating factor of the differential equation :

$$\frac{dy}{dx} + \frac{2}{x}y = 3x^2y^{4/3}, x > 0 \text{ is given by :}$$

- (a) $2 \log x$ (b) x^2 (c) $x^{2/3}$ (d) $x^{-2/3}$

40. The solution of the differential equation :

$$(x + y) \frac{dy}{dx} + 1 = e^{x-y} \text{ is given by :}$$

- (a) $e^y = e^x + c$ (b) $(x + y) e^y = e^x + c$ (c) $e^y = (x + 1) e^x + c$ (d) $y = (x + 1) \log x + c$

41. The solution of differential equation :

$$\sqrt{1+x^2} dy + \sqrt{1+y^2} dx = 0 \text{ is given by :}$$

- (a) $\sqrt{1+y^2} = \sqrt{1+x^2} + c$ (b) $(y + \sqrt{1+y^2}) = (x + \sqrt{1+x^2}) + c$
 (c) $(y + \sqrt{1+y^2})(x + \sqrt{1+x^2}) = c$ (d) $y = (\sqrt{1+x^2} + \sqrt{1+y^2}) + c$

42. The solution of differential equation :

$$a \left(x \frac{dy}{dx} + 2y \right) = xy \frac{dy}{dx} \text{ is given by :}$$

- (a) $yx^2 = e^{\frac{y-c}{a}}$ (b) $y = x^2 e^{\frac{y-c}{a}}$ (c) $y = x^2 + e^{\frac{y-c}{a}}$ (d) $e^y = e^2 + \frac{x^2}{2} + c$

43. The solution of differential equation :

$$\left(x \sin \frac{y}{x} \right) dy = \left(y \sin \frac{y}{x} - x \right) dx \text{ is given by :}$$

- (a) $y = \sin \left(\frac{y}{x} \right) \log |x| + c$ (b) $\cos \left(\frac{y}{x} \right) = \log |x| + c$
 (c) $\sin \left(\frac{y}{x} \right) = \cos \left(\frac{y}{x} \right) + c$ (d) $y = \sin \left(\frac{y}{x} \right) \cos \left(\frac{y}{x} \right) + \log |x| + c$

44. The order of difference equation :

$$u_{x+3} - 5u_{x+2} = 2^x \text{ is :}$$

- (a) 2 (b) 3 (c) 1 (d) 0

45. The general solution of difference equation :

$$u_{x+2} - 7u_{x+1} + 10u_x = 12(4^x) \text{ is :}$$

- (a) $u_x = c_1 2^x + c_2 5^x - 6 \cdot 4^x$ (b) $u_x = c_1 3^x + c_2 4^x - 6 \cdot 2^x$
 (c) $u_x = c_1 x + c_2 x^2 - 12 \cdot 4^x$ (d) $u_x = c_1 e^x + c_2 x^3 - 4^x$

46. The particular integral of the difference equation :

$$u_{x+2} - 7u_{x+1} + 10u_x = 12 \cdot 5^x \text{ is :}$$

- (a) $3 \cdot 4^{x-1} \cdot x$ (b) $4 \cdot 5^x$ (c) $4 \cdot 5^{x-1}$ (d) $4 \cdot 5^{x-1} \cdot x$

47. The general solution of the differential equation $e^{dy/dx} = x^x$ is :
- (a) $y = e^x + c$ (b) $y = \log x + c$
 (c) $y = \frac{x^2}{2} \log x + c$ (d) $y = \frac{x^2}{2} \left(\log x - \frac{1}{2} \right) + c$
48. The slope of the tangent at a point $P(x, y)$ on a curve is $\left\{ -\frac{(y+3)}{(x+2)} \right\}$. If the curve passes through the origin, then equation of curve is :
- (a) $x^2 + 2x + 3 = 0$ (b) $x^2 + 2xy + y^2 = 0$ (c) $xy + 2y + 3x = 0$ (d) $xy + 2x + 3y = 0$
49. If $P(A)$ denotes the probability of an event A in a sample space, then the correct assertion is :
- (a) $P(A) \leq 0$ (b) $P(A) \geq 1$ (c) $0 \leq P(A) \leq 1$ (d) $-1 \leq P(A) \leq 1$
50. Let A and B be two events belonging to a sample space. Then the probability that exactly one of A, B occurs, is equal to :
- (a) $P(A) + P(B)$ (b) $P(A) + P(B) - P(A \cap B)$
 (c) $P(A \cup B) - P(A \cap B)$ (d) $P(A) \cdot P(B)$
51. A letter is selected at random from the word 'PROBABILITY'. The probability that it is a vowel, is
- (a) $2/11$ (b) $3/11$ (c) $4/11$ (d) 4
52. A number is chosen at random from among the first 30 natural numbers. The probability of the number chosen being a prime is :
- (a) $1/3$ (b) $3/10$ (c) $1/30$ (d) $11/30$
53. A point C divides the line segment AB in the ratio $1 : 3$. A car covers a distance AC with a speed of 10 km/hr and the distance CB with a speed of 20 km/hr. If V is the average speed of car, then V is equal to :
- (a) 13.33 km/h (b) 16 km/hr (c) 15 km/hr (d) 17.5 km/hr
54. If S.D. of a variate X is σ , then the S.D. $aX + b$ is :
- (a) $a\sigma + b$ (b) σ (c) $|a|\sigma$ (d) $(a + b)\sigma$
55. Of the following, the only one that is not a measure of dispersion, is
- (a) S.D. (b) Variance (c) Mean Deviation (d) Mode
56. Karl-Pearson's coefficient of skewness of a distribution is 0.32 . Its S.D. is 6.5 and mean is 39.6 . The median of the distribution is given by :
- (a) 28.61 (b) 38.91 (c) 29.13 (d) 28.31
57. The sum of ten number is 12 and the sum of their squares is 16.9 , then their variance is equal to :
- (a) 0.01 (b) 0.10 (c) 0.25 (d) 2.50

58. The Binomial distribution, whose mean is 3 and variance is 2, is given by :

- (a) $\left(\frac{2}{3} + \frac{1}{3}\right)^3$ (b) $\left(\frac{2}{3} + \frac{1}{3}\right)^5$ (c) $\left(\frac{2}{3} + \frac{1}{3}\right)^6$ (d) $\left(\frac{2}{3} + \frac{1}{3}\right)^9$

59. For a bi-variate distribution (x, y) if $\Sigma x = 50$, $\Sigma y = 60$, $\Sigma xy = 350$, $\bar{x} = 5$, $\bar{y} = 6$ variance of x is 4, variance of y is 9, then r_{xy} (Karl-Pearson's Correlation Coefficient) is :

- (a) 5/36 (b) 5/6 (c) 11/3 (d) 11/18

60. Given that $y(0) = 1$, $y(1) = 0$, $y(2) = 1$, $y(3) = 10$, then the value of $y(4)$ is given by :

- (a) 20 (b) 30 (c) 33 (d) 36

61. The best fit straight line (in the sense of least squares) to the following data points

x :	0	1	2	3	4
y :	1	1.8	3.3	4.5	6.3

is given by :

- (a) $y = 1.02x + 5.7$ (b) $y = x + 2$ (c) $y = 1.22 + 0.65$ (d) $y = 1.33x + 0.72$

62. If $P(A) = \frac{3}{8}$, $P(B) = \frac{1}{2}$ and $P(A \cap B) = \frac{1}{4}$, then $P\left(\frac{B}{A}\right)$ is equal to

- (a) 3/8 (b) 3/5 (c) 3/4 (d) 5/8

63. The value of mean and variance are equal in :

- (a) Normal distribution (b) Binomial distribution
(c) Poisson distribution (d) Uniform distribution

64. A set of instructions in a sequential manner telling the computer what to do is called :

- (a) Instructor (b) Comiler (c) Program (d) Key-Board

65. Which of the following performs simple maths for CPU ?

- (a) DIMM (b) ALU (c) BUS (d) Register

66. Which of the following is not a Hardware of a Computer ?

- (a) Monitor (b) Key-Board (c) Winows (d) Mouse

67. Which of the following is called brain of the Computer ?

- (A) Motherboard (b) RAM (c) CPU (d) Memory

68. 'WWW' stands for :

- (a) World Word Web (b) World Wide Web (c) Word Wide Web (d) World Work Web

69. The type of software used to carry out tasks, such as writing a letter, is called :
- (a) GUI software (b) Utility software (c) Application software (d) System software
70. Which of the following key is pressed to work as directional arrow from number pad ?
- (a) Num lock (b) Caps Lock (c) Arrow Lock (d) Shift
71. An Octal number 237 is equal to the Binary number :
- (a) 010 011 111 (b) 010 111 011 (c) 011 101 101 (d) 011 000 001
72. One Gigabyte is approximately equal to :
- (a) 1000 bytes (b) 100,000 bytes (c) 1000, 000, 000 bytes (d) 1000, 000, 000, 000, bytes
73. Which of the following is not a computer language ?
- (a) BASIC (b) COBOL (c) LOTUS (d) FORTRAN
74. In how many different ways can the letters of the word 'ABILITY' be arranged ?
- (a) 1260 (b) 2520 (c) 2420 (d) 720
75. The value of $(73)^2 - (41)^2 + (29)^2$ is equal to :
- (a) 4344 (b) 4321 (c) 4489 (d) 4649
76. The difference between 58% of a number and 37% of the same number is 399. The 72% of that number is equal to :
- (a) 1913 (b) 1330 (c) 1425 (d) 1368
77. The least number to be added to 4700 to make it a perfect square, is :
- (a) 61 (b) 74 (c) 69 (d) 76
78. The cost of 10 pens and 12 pencils is Rs. 138. Then the cost of 15 pens and 18 pencils will be equal to
- (a) Rs. 276 (b) Rs. 878 (c) Rs. 268 (d) Rs. 207
79. If in a certain code RAIL is written as 5796 and TAPE is written as 3748, the PAIR is written in that code as
- (a) 4795 (b) 4785 (c) 8795 (d) 3795
80. How many such pairs to letters are there in the word 'PHYSICAL' each of which has as many letters between them in the word as they have in the English alphabet ?
- (a) One (b) Two (c) Three (d) None
81. If one-fourth of one-fifth of a number is 7, then three - fourteenth of that number will be :
- (a) 42 (b) 60 (c) 70 (d) 30

82. Ram has some hens and some cows. If the total number of animal heads are 43 and total number of feet are 142, then the number of hens Ram has equal to :
- (a) 28 (b) 21 (c) 15 (d) 27
83. The ratio of the present age of Sita and Gita is 3 : 8 respectively. seven years hence the respectively ratio of the their ages will be 4 : 9. Gita's present age is :
- (a) 63 years (b) 42 years (c) 56 years (d) 49 years
84. The average of five consecutive odd number A, B, C, D and E is 45. The product of B and D is equal to
- (a) 2107 (b) 2205 (c) 2021 (d) 1935
85. What should be the compund intereset on an amount of Rs. 5,500 at the rate of 5% per annum after 2 years ?
- (a) Rs. 588 (b) Rs. 645 (c) Rs. 454.50 (d) Rs. 563.75
86. Two candidates contested an election. If one got 520 votes which was 65% of votes, then total number of votes polled i s:
- (a) 858 (b) 800 (c) 780 (d) 754
87. If $x + y = 20$ and $xy = 84$, then the value of $x^2 + y^2$ is equal to .
- (a) 232 (b) 400 (c) 128 (d) 168
88. The number of coins of diameter 1.5 cm and 0.2 cm thick, made out of a right circular cylinder of height 10 cm and diameter 4.5 cm, is
- (a) 500 (b) 4750 (c) 450 (d) 425
89. In the binomial expansio of $(a + b)^n$, the coefficients of 4th and 13th terms are equal to each other. The value of n is :
- (a) 12 (b) 14 (c) 15 (d) 16
90. The coefficient of x^4 in the expansion of $(1 + 2x + 3x^2 + 4x^3 + \dots)^{1/2}$ is :
- (a) 4 (b) 1 (c) 6 (d) 16
91. $\sum_{k=1}^n (-1)^{k^n} C_k$ is :
- (a) -1 (b) 2^* (c) 2^n (d) 0
92. If n is multiple of 3, then the coefficient of x^n in the expansion of $\log(1 + x + x^2)$, is :
- (a) $\frac{1}{n}$ (b) $\frac{2}{n}$ (c) $-\frac{1}{n}$ (d) $-\frac{2}{n}$

93. The sum of the series :

$$\frac{2}{1!} + \frac{6}{2!} + \frac{12}{3!} + \frac{20}{4!} + \dots \infty$$

is :

- (a) e (b) $2e$ (c) $3e$ (d) $\frac{3e}{2}$

94. If $x = \frac{1}{1.2} - \frac{1}{2.3} + \frac{1}{3.4} - \frac{1}{4.5} + \dots \infty$, then e^x is equal to :

- (a) $\frac{4}{e}$ (b) $\frac{e}{4}$ (c) $\log\left(\frac{4}{e}\right)$ (d) $\log\left(\frac{e}{4}\right)$

95. A matrix $A = [a_{ij}]$ is an upper triangular matrix if :

- (a) it is a square matrix and $a_{ij} = 0, i < j$
 (b) it is a square matrix and $a_{ij} = 0, i > j$
 (c) it is not a square matrix and $a_{ij} = 0, i < j$
 (d) it is not a square matrix and $a_{ij} = 0, i > j$

96. If A and B are two invertible matrices, then the inverse of AB is equal to

- (a) AB (b) BA (c) $A^{-1}B^{-1}$ (d) $B^{-1}A^{-1}$

97. The matrix A satisfying the equation :

$$\begin{bmatrix} 1 & 3 \\ 0 & 1 \end{bmatrix} A = \begin{bmatrix} 1 & 1 \\ 0 & -1 \end{bmatrix}$$

is :

- (a) $\begin{bmatrix} 1 & 4 \\ -1 & 0 \end{bmatrix}$ (b) $\begin{bmatrix} 1 & -4 \\ 1 & 0 \end{bmatrix}$
 (c) $\begin{bmatrix} 1 & 4 \\ 0 & -1 \end{bmatrix}$ (d) $\begin{bmatrix} -1 & -4 \\ 1 & 0 \end{bmatrix}$

98. The solution set of equation

$$\begin{vmatrix} 1 & 3 & x \\ 1 & 1 & x^2 \\ 3 & 7 & 3 \end{vmatrix} = 0$$

is :

- (a) ϕ (b) $\{0, 1\}$ (c) $\{1, -1\}$ (d) $\{1, -3\}$

99. If the Cramer's rule fails to solve the system of equations

$$\lambda x + y - z = 1$$

$$2x + 3y + z = 2$$

$$x - y + z = 0$$

then λ is equal to :

(a) 0

(b) -1

(c) 1

(d) -2

100. The value of the determinant :

$$\begin{vmatrix} 1 & x & y+z \\ 1 & y & z+x \\ 1 & z & x+y \end{vmatrix} \text{ is}$$

(a) 0

(b) $x + y + z$

(c) $1 + x + y + z$

(d) $(x - y)(y - z)(z - x)$