

PART A

1. A watch which gains 5 seconds in 3 minutes was set right at 7 a.m. In the afternoon of the same day, when the watch indicated quarter past 4 hrs, the true time is
(A) 3 hrs $59\frac{7}{12}$ min (B) 4 hrs (C) 4 hrs $2\frac{3}{11}$ min (D) 3 hrs $58\frac{7}{11}$ min
2. In how many different ways can the letters of the word 'LEADER' be arranged in such a way that the vowels always come together ?
(A) 72 (B) 144 (C) 48 (D) 720
3. A rectangular park 60m long and 40m wide has two concrete crossroads running in the middle of the part and rest of the park has been used as a lawn. If the area of the lawn is 2109 sq.m., then what is the width of the road
(A) 5.82 m (B) 2.91 m (C) 3 m (D) 1.91 m

JITENDRA MISHRA ACADEMY (JMA), INDORE (India's No. 1 Institute for All India MCA Entrance Training)

JMA HOUSE - 7, CHANDRALOK COLONY, INDORE (M.P.) Ph.: 0731 - 4236844 / 2566799 Visit us : www.jmaindore.com

4. Insert the missing number in the series 16, 33, 65, 131, 261,
(A) 521 (B) 721 (C) 523 (D) 613
5. Find the wrong number in the series 7, 8, 18, 57, 228, 1165, 6996
(A) 228 (B) 8 (C) 1165 (D) 57
6. If $D =$ the day $(1 - 366)$ in year Y , then the day of the week for D can be calculated using the following formula :
$$d = \left(D + Y + \frac{Y-1}{4} - \frac{Y-1}{100} + \frac{Y-1}{400} \right) \text{ mod } 7$$
Where $d = 1$ would mean Sunday, $2 =$ Monday and so on. Which of the following days can the first day of a century NOT be ?
(A) Thursday (B) Monday (C) Saturday (D) Friday
7. If A and B run a race, then A wins by 60 seconds. If B and C run the same race, then B wins by 30 seconds. Assuming that C maintains a uniform speed to find the time taken by C to finish the race, which of the following statement are needed ?
 - I. A and C run the same race and A wins by 375 metres.
 - II. The length of the race is 1 km.

(A) Either I or II is sufficient to answer the question
(B) Both I and II are needed to answer the question
(C) Both are not sufficient to answer the question
(D) Not enough information to answer the question

8. Consider the following statements :
- There are six village A, B, C, D, E and F.
- F is 1 km to the west of D
 - B is 1 km to the east of E
 - A is 2 km to the north of E
 - C is 1 km to the east of B
 - D is 1 k to the south of A
- Which three villages are in a straight line ?
- (A) A, D, E (B) A, C, B (C) E, B, D (D) C, B, F
9. Find the odd man out in the following series ? 844, 211, 120, 752, 541, 303
- (A) 303 (B) 211 (C) 844 (D) 120
10. There are five hobby clubs in a college viz photography, yachting, cheess, electronics and gardening. The gardening group meets every second day, the electronics groups meets every third day, the chess group meets every fourth day, the yachting group meets every fifth day and the photography group meets every sixth day. How many time do all the five groups meet on the same day within 360 days ?
- (A) 18 (B) 12 (C) 6 (D) 30

JITENDRA MISHRA ACADEMY (JMA), INDORE (India's No. 1 Institute for All India MCA Entrance Training)

JMA HOUSE - 7, CHANDRALOK COLONY, INDORE (M.P.) Ph.: 0731 - 4236844 / 2566799 Visit us : www.jmaindore.com

11. Four political parties W, X, Y and Z decided to set up a joint candidate for the coming elections. The formula agreed by them was the acceptance of a candidate of the most of the parties. Four aspiring candidates A, B, C and D approached the parties for thicket.
- A was acceptable to W but not Z.
 - B was acceptable to Y but not X.
 - C was acceptable to W and Y.
 - D was acceptable to W and X.
- Further, candidates B was preferred by W and Z, candidates C was preferred by X and Z and candidate A was acceptable to X but not Y; who got the ticket.
- (A) B (B) A (C) D (D) C
12. Raman walks 100 m from his house towards North. From there he goes 100 m towards West. Here is the house of Shyam. From there they both go to the market which is in the perfect South – West direction from Shyam's house. If the market is exactly to the West of Raman's house, then the how far is the market from Raman's house ?
- (A) 100 m (B) 200 m (C) 250 m (D) 150 m
13. How many three digit numbers can generated from 1, 2, 3, 4, 5, 6, 7, 8, 9 such that the digits are in ascending order.
- (A) 81 (B) 80 (C) 84 (D) 83

JITENDRA MISHRA ACADEMY (JMA), INDORE (India's No. 1 Institute for All India MCA Entrance Training)

JMA HOUSE - 7, CHANDRALOK COLONY, INDORE (M.P.) Ph.: 0731 - 4236844 / 2566799 Visit us : www.jmaindore.com

14. Two cubes of sides 6 cm each are kept side by side to from a rectangular parallelepiped. What is the surface area of the rectangular parrallelepiped ?
- (A) 180 (B) 360 (C) 432 (D) 288
15. The sum of the length of all the edges of a rectangular box is 12 cm. If the total surface area of the box is 5 sq. cm., then length of its diagonal is
- 2 cm
 - 13 cm
 - Not enough information to find the diagonal
 - 3 cm

Booklet Code - B

16. Three persons A, B and C wore shirts of black, blue and orange colours (not necessarily in that order) and pants of green, yellow and orange colours (not necessarily in that order). No persons wear shirts and pants of the same colour. Further it is given that :
- A did not wear shirt of black colour.
 - B did not wear shirt of blue colour.
 - C did not wear shirt of orange colour.
 - A did not wear pant of green colour.
 - B wore a pant of orange colour.
- What were the colours of the pants and shirts wore by C respectively ?
- Yellow and Blue
 - Orange and Black
 - Yellow and Black
 - Green and Blue

17. If South – East becomes North, North-East becomes West and son on, what will West become ?
- South – East
 - North – East
 - South – West
 - North – West

JITENDRA MISHRA ACADEMY (JMA), INDORE (India's No. 1 Institute for All India MCA Entrance Training)

JMA HOUSE - 7, CHANDRALOK COLONY, INDORE (M.P.) Ph.: 0731 - 4236844 / 2566799 Visit us : www.jmaindore.com

18. In the following number line, how many times do the two consecutive numbers have a difference of 2 ?
7 5 9 5 2 3 5 9 4 8 5 9 5 4 5 9 3 5 5 9 5 3 5 9 4 5 2 5 3 6 5 9
- 7
 - 9
 - 8
 - 5
19. The letter L, M, N, O, P, Q, R, S and T in their order are substituted by nine integers 1 to 9 but not in that order. 4 is assigned to P. The difference between P and T is 5. The difference between N and T is 3. What is the integer assigned to N ?
- 5
 - 7
 - 6
 - 4
20. In five flats, one above the other, live five professionals. The professor has to go up the met his IAS officers friend. The doctor is equally friendly to all, and has to go up as frequently as go down. The engineer hat to go up to meet his Lawyer friend above whose flat lives the professors friends. From the ground floor to the top floor, in what order do the five professionals live ?
- Professors, Engineer, Doctor, IAS officer, Lawyer
 - Engineer, Professors, Doctor, IAS officer, Lawyer
 - Professors, Engineer, Doctor, Lawyer, IAS officers
 - IAS officer, Engineer, Doctor, Professors, Lawyer

JITENDRA MISHRA ACADEMY (JMA), INDORE (India's No. 1 Institute for All India MCA Entrance Training)

JMA HOUSE - 7, CHANDRALOK COLONY, INDORE (M.P.) Ph.: 0731 - 4236844 / 2566799 Visit us : www.jmaindore.com

21. If '245' means 'Art and Talent' in a certain code language. '316' means 'Callons to Generous', '147' means 'Callous and Polite' then what is the code used for 'to' ?
- 3 or 6
 - Only 3
 - Only 6
 - Only 1
22. In a 100 m race, Bipin, Chandan, Danny and Feroz won the first four prizes, not necessarily in the same order. Each of there four sprinters hails from a different city among Delhi, Mumbai, Kolkata and Chennai. Given the following conditions, who got the first prize ?
- Bipin is not from Kolkata and didn't get the first prize. The person from Chennai is the winner. Danny is from Mumbai
 - Feroz is either from Delhi or from Chennai.
- Chandan
 - Danny
 - Can't be determined
 - Feroz
23. If $\log 2 = 0.3010$ and $\log 3 = 0.4771$, then value of $\log_5 512$ is
- 3.876
 - 2.870
 - 3.912
 - 2.867

Booklet Code - B

24. In a class of 45 students a boys is ranked 20th. When two boys joined his rank dropped by one. What is his new rank from the end ?
 (A) 26th (B) 25th (C) 28th (D) 27th
25. The least perfect square which is divisible by 21.36 and 66 is
 (A) 214344 (B) 213444 (C) 221444 (D) 214431

PART B

26. The sum of the series $1.2.3 + 2.3.4 + 3.4.5 + \dots + 18.19.20$ is
 (A) 45450 (B) 24560 (C) 35910 (D) 44820
27. In a triangle, three angles are in Arithmetic Progression and ratios of smallest to largest angle is 1 : 5. What is the smallest angle ?
 (A) $\frac{\pi}{10}$ (B) $\frac{\pi}{8}$ (C) $\frac{\pi}{6}$ (D) $\frac{\pi}{9}$

JITENDRA MISHRA ACADEMY (JMA), INDORE (India's No. 1 Institute for All India MCA Entrance Training)

JMA HOUSE - 7, CHANDRALOK COLONY, INDORE (M.P.) Ph.: 0731 - 4236844 / 2566799 Visit us : www.jmaindore.com

28. 2's complement of 010101.0101 is
 (A) 101010.1010 (B) 101011.1010 (C) 101011.0010 (D) 101010.1011
29. A logic circuit has three inputs bits; x_0 , x_1 and x_2 , where x_0 is the least significant bit and x_2 is the most significant bit. The output from the circuit is 1 when its inputs is any of the 3-bit numbers, 1, 4, 5 or 6; otherwise the output is 0. Which of the following expression represents the output from this circuit ? (x_2 is complement of x_2)
 (A) $x_2 x_0 + x_2 x_1$ (B) $x_2 + x_1 + x_0$ (C) $x_2 x_0 + x_1 x_0$ (D) $x_1 x_0 + x_2 x_0$

JITENDRA MISHRA ACADEMY (JMA), INDORE (India's No. 1 Institute for All India MCA Entrance Training)

JMA HOUSE - 7, CHANDRALOK COLONY, INDORE (M.P.) Ph.: 0731 - 4236844 / 2566799 Visit us : www.jmaindore.com

30. Let A be a 4×4 matrix. Suppose that a matrix B is obtained from A by using following row operations :
 • $R_1 \leftrightarrow R_2$
 • $R_3 \leftrightarrow 4 \times R_3$
 • $R_4 \leftrightarrow R_2$
 • $R_2 \leftrightarrow R_2 + 3 \times R_3$
 If $|A| = 20$, then what is $|B|$?
 (A) 5 (B) -80 (C) 80 (D) -5
31. Two vectors a and b are perpendicular to each other such that length of a is 8 and length of b is 3. What is $|a - 2b|$?
 (A) 2 (B) 10 (C) 12 (D) 6




Figure 1 : Vector a, b and c

Booklet Code - B

32. Express Vector n in terms of vectors a , b and c shown in Fig. 1.
 (A) $-a + b + c$ (B) $a - b - c$ (C) $-a - b - c$ (D) $a + b + c$
33. Logic X-OR operations of Hexadecimal number (4AC0) and (B53F) is
 (A) FFFF (B) AACB (C) ABCD (D) 0000
34. Find the sum of the series $1.1! + 2.2! + 3.3! + \dots + 10.10!$ is
 (A) 39916800 (B) 39916801 (C) 29999999 (D) 39916799

JITENDRA MISHRA ACADEMY (JMA), INDORE (India's No. 1 Institute for All India MCA Entrance Training)

JMA HOUSE - 7, CHANDRALOK COLONY, INDORE (M.P.) Ph.: 0731 - 4236844 / 2566799 Visit us : www.jmaindore.com

35. A circle $x^2 + y^2 - x - 3y - \frac{25}{2} = 0$ has a chord $x - y - 1 = 0$ which forms the diameter of another circle. What is the equation representing this second circle ?
 (A) $x^2 + y^2 - 3y - x - \frac{21}{2} = 0$
 (B) $x^2 + y^2 - 3y - y - \frac{21}{2} = 0$
 (C) $x^2 + y^2 - 3y - 3y - \frac{21}{2} = 0$
 (D) $x^2 + y^2 - 3y - y - \frac{21}{2} = 0$
36. The equation of the locus of a point, which moves so that the sum of its distances from two given points $(ae, 0)$ and $(-ae, 0)$ is equal to $2a$, is
 (A) $\frac{x^2}{b^2} + \frac{y^2}{a^2} = 1$ where $b^2 = a^2(1 - e^2)$
 (B) $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ where $b^2 = a^2(1 - e^2)$
 (C) $(bx)^2 + (ay)^2 =$ where $b^2 = a^2(1 - e^2)$
 (D) $(ax)^2 + (by)^2 =$ where $b^2 = a^2(1 - e^2)$

JITENDRA MISHRA ACADEMY (JMA), INDORE (India's No. 1 Institute for All India MCA Entrance Training)

JMA HOUSE - 7, CHANDRALOK COLONY, INDORE (M.P.) Ph.: 0731 - 4236844 / 2566799 Visit us : www.jmaindore.com

37. All values of α for which the point (α, α^2) lies inside the triangle formed by the lines $2x + 3y - 1 = 0$, $x + 2y - 3 = 0$ and $5x - 6y - 1 = 0$ are
 (A) $\alpha \in \left(\frac{-3}{2}, \frac{-3}{4}\right) \cup \left(\frac{1}{2}, 1\right)$ (B) $\alpha \in \left(\frac{-3}{2}, -1\right) \cup \left(\frac{1}{2}, 1\right)$
 (C) $\alpha \in \left(\frac{-5}{2}, -1\right) \cup \left(\frac{1}{2}, 1\right)$ (D) $\alpha \in \left(-1, \frac{-3}{4}\right) \cup \left(\frac{1}{2}, 1\right)$
38. Which is the equation of a line whose slope is undefined ?
 (A) $X = -5$ (B) $Y = 7$ (C) $X + Y = 0$ (D) $X = Y$
39. $y = e^{-3x}$ is solution of the differential equation
 (A) $\frac{dy}{dx} + y^3 = 0$ (B) $3\frac{dy}{dx} + y = 0$ (C) $\frac{dy}{dx} + 3y^2 = 0$ (D) $\frac{dy}{dx} + 3y = 0$
40. The area of the region bounded by the curve $y^2 = 4x$, y - axis and the line $y = 3$ is
 (A) $28/3$ sq. units (B) $9/4$ sq. units (C) $9/16$ sq. units (D) $3/4$ sq. units

Booklet Code - B

41. Let c_1 and c_2 be constants. The general solution of the differential equation $\frac{d^2y}{dx^2} - y = 0$ is
- (A) $y = c_1e^x$ (B) $y = c_1e^x + c_2e^{-x}$
 (C) $y = c_1e^{x^2} + c_2e^{-x^2}$ (D) $y = c_2e^{-x}$
42. Let the set S contains all strings w of 0s and 1s which have the following property; w contains an equal number of occurrences of 01 and 10. Which of the following words DOES NOT be long to S ?
 (A) 101010 (B) 1010101 (C) 0101010 (D) All of the above
43. Let N_2^+ denote the natural numbers greater than or equal to 2. Let m be a relation with n , mRn if $\gcd(m, n) > 1$. The binary relation R on N_2 is
 (A) Reflexive, Symmetric, Transitive
 (B) Reflexive, Not Symmetric, Transitive
 (C) Reflexive, Not Symmetric, Not Transitive
 (D) Reflexive, Symmetric, Not Transitive

JITENDRA MISHRA ACADEMY (JMA), INDORE (India's No. 1 Institute for All India MCA Entrance Training)

JMA HOUSE - 7, CHANDRALOK COLONY, INDORE (M.P.) Ph.: 0731 - 4236844 / 2566799 Visit us : www.jmaindore.com

44. If two sets A and B have 500 elements in common, then the number of common element in the sets $A \times B$ and $B \times A$ is :
 (A) 500 (B) 500^2 (C) 2^{500} (D) 1000
45. What is the number of points of intersection between the graphs given by the functions $f(x) = x^3 + 2$ and $g(x) = x^2$, where $x \in R$?
 (A) 3 (B) 2 (C) 1 (D) 0
46. A ray of light is sent along the line $x - 2y - 3 = 0$. On reaching the line $3x - 2y - 5 = 0$ the ray is reflected from it. Find the equation of the line containing the reflected ray.
 (A) $29x - 2y - 13 = 0$ (B) $2x - 29y - 31 = 0$ (C) $29x - 2y - 31 = 0$ (D) $2x - 29y - 13 = 0$

JITENDRA MISHRA ACADEMY (JMA), INDORE (India's No. 1 Institute for All India MCA Entrance Training)

JMA HOUSE - 7, CHANDRALOK COLONY, INDORE (M.P.) Ph.: 0731 - 4236844 / 2566799 Visit us : www.jmaindore.com

47. Let $f(x) = 4x(1 - x)$ be defined for all $x \in R$. Then, the number of real numbers which satisfy the equation, $f(x) = x$ is
 (A) 1 (B) 0 (C) 3 (D) 2
48. The correlation coefficient between X and Y where $Y = X^3$ and X follows normal distribution with mean 0 and variance 1 is
 (A) $\frac{1}{\sqrt{15}}$ (B) 0 (C) $\frac{3}{\sqrt{15}}$ (D) $\frac{2}{\sqrt{15}}$
49. There are 13 players and the deck of cards are distributed equally amongst all the players at random. What is the probability of the player getting all the queens ?
 (A) $\frac{1}{3^4 \times 4 \times 47}$ (B) $\frac{1}{3 \times 4^3 \times 47}$ (C) $\frac{1}{3^3 \times 4^3 \times 47}$ (D) $\frac{1}{3^3 \times 4^4 \times 47}$
50. The coefficient variation for the data $X = [16 \ 98 \ 96 \ 49 \ 81 \ 15 \ 43 \ 92 \ 80 \ 96]$ is :
 (A) 49.63 (B) 33.05 (C) 99.93 (D) 66.6
51. Which of the following is not a linear mapping given that $f : R^2 \rightarrow R^2$?
 (A) $(x_1, x_2) \rightarrow (x_1 + x_2, x_2)$ (B) $(x_1, x_2) \rightarrow (x_2, x_1)$
 (C) $(x_1, x_2) \rightarrow (10, 0)$ (D) $(x_1, x_2) \rightarrow (x_1 + 1, x_2)$

52. The function f is twice differentiable and the graph of f has no points of inflection. If $f(6) = 3$, $f'(6) = -\frac{1}{2}$ and $f''(6) = -2$ which of the following could be the value of $f(7)$?
 (A) 2.5 (B) 2 (C) 3 (D) 2.9
53. A spherical balloon is being inflated at a rate of $1\text{m}^3/\text{s}$. How is the diameter of the balloon changing at the instant when the radius of the balloon is 2m ?
 (A) $\frac{1}{16\pi}\text{m/s}$ (B) $\frac{1}{4\pi}\text{m/s}$ (C) $\frac{1}{8\pi}\text{m/s}$ (D) None of the above
54. $\int_1^e \frac{\ln(x^2)}{x} dx =$
 (A) e (B) 0 (C) 1 (D) $e - 1$

JITENDRA MISHRA ACADEMY (JMA), INDORE (India's No. 1 Institute for All India MCA Entrance Training)

JMA HOUSE - 7, CHANDRALOK COLONY, INDORE (M.P.) Ph.: 0731 - 4236844 / 2566799 Visit us : www.jmaindore.com

55. Find the arc-length of the curve given by the parametric equation $x = \frac{t^2}{\sqrt{2}}$ and $y = \frac{t^3}{3}$, where $0 \leq t \leq 1$.
 (A) $\sqrt{2} + \sqrt{3}$ (B) $\sqrt{3} + \frac{2\sqrt{2}}{3}$ (C) $\sqrt{3} - \frac{2\sqrt{2}}{3}$ (D) $\sqrt{3} - \sqrt{2}$
56. Let $w = \log(u^2 + v^2)$ where $u = e^{(x^2+y)}$ and $v = e^{(x+y^2)}$. Then $\left. \frac{\partial w}{\partial x} \right|_{x=0, y=0}$
 (A) 1 (B) 0 (C) 4 (D) 2
57. The differential equation of all the ellipse centred at the origin is :
 (A) $xyy'' + x(y')^2 - yy' = 0$ (B) $y^2 + x(y')^2 - yy' = 0$
 (C) $yy'' + x(y')^2 - xy' = 0$ (D) None of the above

JITENDRA MISHRA ACADEMY (JMA), INDORE (India's No. 1 Institute for All India MCA Entrance Training)

JMA HOUSE - 7, CHANDRALOK COLONY, INDORE (M.P.) Ph.: 0731 - 4236844 / 2566799 Visit us : www.jmaindore.com

58. Let X be a binomial random variable with $n = 5$ and $p = 0.5$ and $Y = X \bmod 2$. Probability of $Y = 0$ is
 (A) $\frac{1}{2^3}$ (B) $\frac{1}{2^5}$ (C) 1 (D) $\frac{1}{2}$
59. A 120 gallon tank initially contains 90 lb of salt dissolved in 90 gallons of water. Solution containing 2 lb / gallon of salt flows into the tank at the rate of 4 gallons / min and the well - stirred mix - true flow out of the tank at the rate of 3 gallons / min. A differential equation for the amount x (in lb) of salt in the tank is :
 (A) $\frac{dx}{dt} + \frac{3x}{90+2t} = 4$ (B) $\frac{dx}{dt} + \frac{x}{30} = 8$
 (C) $\frac{dx}{dt} + 90x = 2$ (D) $\frac{dx}{dt} + \frac{3x}{90+t} = 8$

Questions 60 – 62 are based on the flowchart shown in Fig. 2.

60. If the following are the elements in K , 3 7 9 16 73 97 683 923 3883 9003 then the number of iterations to determine whether 7 is in K is :
 (A) 2 (B) 3 (C) 4 (D) 1




Figure 2 : Flow Chart

61. How many total iterations are needed to find if an element is not in K given that there are N element in K ?
 (A) N^2 (B) N (C) $N - 1$ (D) $\log N$
62. If the following are the elements in K, 3 5 7 12 17 28 49 67 89 101 121 138 147 164 189 201 220 345 567 879 903 1001, what is the number pointed to by M ?
 (A) 345 (B) 567 (C) 903 (D) 879

JITENDRA MISHRA ACADEMY (JMA), INDORE (India's No. 1 Institute for All India MCA Entrance Training)

JMA HOUSE - 7, CHANDRALOK COLONY, INDORE (M.P.) Ph.: 0731 - 4236844 / 2566799 Visit us : www.jmaindore.com

63. A teacher tells her students that although she always use an average to calculate their course grades, she gives more weight to the final exam grade. She assures students that if they can perform well on final, then even if they performed poorly on the other exams, they must have learned the material. For three semesters she kept track of how students did on the final and how they did in the course which is given below. What is the probability that a student, taken at random from the class, would have passed her course ?

	Course	
Final	Pass	Fail
Pass	142	34
Fail	89	56

- (A) 0.52 (B) 0.62 (C) 0.42 (D) 0.72

JITENDRA MISHRA ACADEMY (JMA), INDORE (India's No. 1 Institute for All India MCA Entrance Training)

JMA HOUSE - 7, CHANDRALOK COLONY, INDORE (M.P.) Ph.: 0731 - 4236844 / 2566799 Visit us : www.jmaindore.com

64. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be the function defined by $f(x) = \frac{\sin x}{|x| + \cos x}$, Then
- (A) f is not differentiable at $x = 0$
 (B) f is differentiable at all $x \in \mathbb{R}$
 (C) f is differentiable at $x = 0$ but f' is not continuous at $x = 0$
 (D) f is not differentiable at $x = \frac{\pi}{2}$

Booklet Code - B

65. If f is a continuous function on the closed interval $[a, b]$. which of the following MUST be true ?
 (A) There is a number c in the open interval (a, b) such that $f(a) < f(c) < f(b)$.
 (B) There is a number c in the open interval (a, b) such that $f(c) = 0$
 (C) There is a number c in the open interval (a, b) such that $f'(c) = 0$.
 (D) There is a number c in the closed interval $[a, b]$ such that $f(c) \geq f(x)$ for all $x \in [a, b]$
66. The differential equation $\frac{dy}{dx} = (x + y + 1)^2$ is
 (A) linear (B) separable (C) homogeneous (D) None of the above
67. If $\int_0^1 f(x)dx = 2$ and $\int_0^1 f(x)dx = -3$ then, $\int_1^4 (3f(x) + 2)dx =$
 (A) -9 (B) -13 (C) 21 (D) 3

JITENDRA MISHRA ACADEMY (JMA), INDORE (India's No. 1 Institute for All India MCA Entrance Training)

JMA HOUSE - 7, CHANDRALOK COLONY, INDORE (M.P.) Ph.: 0731 - 4236844 / 2566799 Visit us : www.jmaindore.com

68. In the interval $(-2\pi, 0)$, the function $f(x) = \sin\left(\frac{1}{x^3}\right)$ changes sign :
 (A) One
 (B) Never
 (C) Infinite number of times
 (D) More than once but finite number of times
69. A set $S = \left\{s \mid s = \frac{3m+5}{m+2}, m \geq 0\right\}$ has
 (A) neither a minimum element nor a maximum element in it.
 (B) both maximum element and minimum element in it.
 (C) a maximum element but not a minimum element in it
 (D) a minimum element but not a maximum element in it.

JITENDRA MISHRA ACADEMY (JMA), INDORE (India's No. 1 Institute for All India MCA Entrance Training)

JMA HOUSE - 7, CHANDRALOK COLONY, INDORE (M.P.) Ph.: 0731 - 4236844 / 2566799 Visit us : www.jmaindore.com

70. Let function $f : X \rightarrow X$ and g is its inverse function i.e.,
 $g \circ f(x) = x, \forall x \in X$
 then
 (A) f must be 1-to-1 but not always onto
 (B) f is always 1-to-1 and onto
 (C) f must be onto but not always 1-to-1
 (D) None of the above is always true
71. Centers of two circles are separated by 30 cms. Radius of first circle is 40 cm and radius of second circle is 50 cm. What is the length of the common chord ?
 (A) 48 (B) 24 (C) 56 (D) 40
72. Which of the following are true for all real values of x ? All arguments are given in radians.
 I. $\sin\left(\frac{\pi}{2} + x\right) = \cos\left(\frac{\pi}{2} - x\right)$
 II. $2 + 2 \sin(x) - \cos^2(x) \geq 0$
 III. $\sin\left(x + \frac{3\pi}{2}\right) = \cos(\pi - x)$
 IV. $\sin(x) \cos(x) \leq \frac{1}{4}$
 (A) I and III (B) I and II (C) II and III (D) III and IV

Booklet Code - B

73. For all sets A, B and C, which of the following statement is TRUE ?

(A) $(A - B) \cap (C - B) = (A \cap C) - B$

(B) $A - (B - C) = (A - B) - C$

(C) If $A \cap C = B \cap C$ then $A = B$

(D) $(A - B) \cap (C - B) = A - (B \cup C)$

74. Which among the following functions is not periodic :

(A) $e^{\cos(x)}$

(B) $\log_e(\sin(x))$

(C) $\sin(e^x)$

(D) $\frac{10}{100 + \sin(x) + \cos(x)}$

75. The number of roots of the equation $X^2 + \sin^2(X) = 1$ in the closed interval $\left[0, \frac{\pi}{2}\right]$ is

(A) 1

(B) 0

(C) Infinite

(D) Finite and greater than 1