

# Mathematics Notes for Class 12 chapter 1.

## Relations and Functions

### Relation

If A and B are two non-empty sets, then a relation R from A to B is a subset of  $A \times B$ .

If  $R \subseteq A \times B$  and  $(a, b) \in R$ , then we say that a is related to b by the relation R, written as  $aRb$ .

### Domain and Range of a Relation

Let R be a relation from a set A to set B. Then, set of all first components or coordinates of the ordered pairs belonging to R is called : the domain of R, while the set of all second components or coordinates = of the ordered pairs belonging to R is called the range of R.

Thus, domain of  $R = \{a : (a, b) \in R\}$  and range of  $R = \{b : (a, b) \in R\}$

### Types of Relations

**(i) Void Relation** As  $\Phi \subset A \times A$ , for any set A, so  $\Phi$  is a relation on A, called the empty or void relation.

**(ii) Universal Relation** Since,  $A \times A \subseteq A \times A$ , so  $A \times A$  is a relation on A, called the universal relation.

**(iii) Identity Relation** The relation  $I_A = \{(a, a) : a \in A\}$  is called the identity relation on A.

**(iv) Reflexive Relation** A relation R is said to be reflexive relation, if every element of A is related to itself.

Thus,  $(a, a) \in R, \forall a \in A = R$  is reflexive.

**(v) Symmetric Relation** A relation R is said to be symmetric relation, iff

$$(a, b) \in R \implies (b, a) \in R, \forall a, b \in A$$

$$\text{i.e., } a R b \implies b R a, \forall a, b \in A$$

$\implies R$  is symmetric.

**(vi) Anti-Symmetric Relation** A relation R is said to be anti-symmetric relation, iff

$$(a, b) \in R \text{ and } (b, a) \in R \implies a = b, \forall a, b \in A$$

**(vii) Transitive Relation** A relation  $R$  is said to be transitive relation, iff  $(a, b) \in R$  and  $(b, c) \in R$

$$\Rightarrow (a, c) \in R, \forall a, b, c \in A$$

**(viii) Equivalence Relation** A relation  $R$  is said to be an equivalence relation, if it is simultaneously reflexive, symmetric and transitive on  $A$ .

**(ix) Partial Order Relation** A relation  $R$  is said to be a partial order relation, if it is simultaneously reflexive, symmetric and anti-symmetric on  $A$ .

**(x) Total Order Relation** A relation  $R$  on a set  $A$  is said to be a total order relation on  $A$ , if  $R$  is a partial order relation on  $A$ .

### Inverse Relation

If  $A$  and  $B$  are two non-empty sets and  $R$  be a relation from  $A$  to  $B$ , such that  $R = \{(a, b) : a \in A, b \in B\}$ , then the inverse of  $R$ , denoted by  $R^{-1}$ , is a relation from  $B$  to  $A$  and is defined by

$$R^{-1} = \{(b, a) : (a, b) \in R\}$$

### Equivalence Classes of an Equivalence Relation

Let  $R$  be equivalence relation in  $A$  ( $\neq \Phi$ ). Let  $a \in A$ .

Then, the equivalence class of  $a$  denoted by  $[a]$  or  $\{a\}$  is defined as the set of all those points of  $A$  which are related to  $a$  under the relation  $R$ .

### Composition of Relation

Let  $R$  and  $S$  be two relations from sets  $A$  to  $B$  and  $B$  to  $C$  respectively, then we can define relation  $SoR$  from  $A$  to  $C$  such that  $(a, c) \in SoR \Leftrightarrow \exists b \in B$  such that  $(a, b) \in R$  and  $(b, c) \in S$ .

This relation  $SoR$  is called the composition of  $R$  and  $S$ .

- (i)  $RoS \neq SoR$
- (ii)  $(SoR)^{-1} = R^{-1}oS^{-1}$

known as **reversal rule**.

### Congruence Modulo $m$

Let  $m$  be an arbitrary but fixed integer. Two integers  $a$  and  $b$  are said to be congruence modulo  $m$ , if  $a - b$  is divisible by  $m$  and we write  $a \equiv b \pmod{m}$ .

i.e.,  $a \equiv b \pmod{m} \Leftrightarrow a - b$  is divisible by  $m$ .

## Important Results on Relation

- If  $R$  and  $S$  are two equivalence relations on a set  $A$ , then  $R \cap S$  is also an equivalence relation on  $A$ .
- The union of two equivalence relations on a set is not necessarily an equivalence relation on the set.
- If  $R$  is an equivalence relation on a set  $A$ , then  $R^{-1}$  is also an equivalence relation on  $A$ .
- If a set  $A$  has  $n$  elements, then number of reflexive relations from  $A$  to  $A$  is  $2^{n^2-2}$ .
- Let  $A$  and  $B$  be two non-empty finite sets consisting of  $m$  and  $n$  elements, respectively. Then,  $A \times B$  consists of  $mn$  ordered pairs. So, total number of relations from  $A$  to  $B$  is  $2^{nm}$ .

## Binary Operations

### Closure Property

An operation  $*$  on a non-empty set  $S$  is said to satisfy the closure property, if

$$a \in S, b \in S \Rightarrow a * b \in S, \forall a, b \in S$$

Also, in this case we say that  $S$  is closed for  $*$ .

An operation  $*$  on a non-empty set  $S$ , satisfying the closure property is known as a binary operation.

or

Let  $S$  be a non-empty set. A function  $f$  from  $S \times S$  to  $S$  is called a binary operation on  $S$  i.e.,  $f : S \times S \rightarrow S$  is a binary operation on set  $S$ .

### Properties

- Generally binary operations are represented by the symbols  $*$ ,  $+$ , ... etc., instead of letters figure etc.
- Addition is a binary operation on each one of the sets  $N$ ,  $Z$ ,  $Q$ ,  $R$  and  $C$  of natural numbers, integers, rationals, real and complex numbers, respectively. While addition on the set  $S$  of all irrationals is not a binary operation.
- Multiplication is a binary operation on each one of the sets  $N$ ,  $Z$ ,  $Q$ ,  $R$  and  $C$  of natural numbers, integers, rationals, real and complex numbers, respectively. While multiplication on the set  $S$  of all irrationals is not a binary operation.
- Subtraction is a binary operation on each one of the sets  $Z$ ,  $Q$ ,  $R$  and  $C$  of integers, rationals, real and complex numbers, respectively. While subtraction on the set of natural numbers is not a binary operation.
- Let  $S$  be a non-empty set and  $P(S)$  be its power set. Then, the union and intersection on  $P(S)$  is a binary operation.

- Division is not a binary operation on any of the sets  $N$ ,  $Z$ ,  $Q$ ,  $R$  and  $C$ . However, it is not a binary operation on the sets of all non-zero rational (real or complex) numbers.
- Exponential operation  $(a, b) \rightarrow a^b$  is a binary operation on set  $N$  of natural numbers while it is not a binary operation on set  $Z$  of integers.

### Types of Binary Operations

**(i) Associative Law** A binary operation  $*$  on a non-empty set  $S$  is said to be associative, if  $(a * b) * c = a * (b * c)$ ,  $\forall a, b, c \in S$ .

Let  $R$  be the set of real numbers, then addition and multiplication on  $R$  satisfies the associative law.

**(ii) Commutative Law** A binary operation  $*$  on a non-empty set  $S$  is said to be commutative, if

$$a * b = b * a, \forall a, b \in S.$$

Addition and multiplication are commutative binary operations on  $Z$  but subtraction not a commutative binary operation, since

$$2 - 3 \neq 3 - 2.$$

Union and intersection are commutative binary operations on the power  $P(S)$  of all subsets of set  $S$ . But difference of sets is not a commutative binary operation on  $P(S)$ .

**(iii) Distributive Law** Let  $*$  and  $\circ$  be two binary operations on a non-empty sets. We say that  $*$  is distributed over  $\circ$ , if

$$a * (b \circ c) = (a * b) \circ (a * c), \forall a, b, c \in S \text{ also called (left distribution) and } (b \circ c) * a = (b * a) \circ (c * a), \forall a, b, c \in S \text{ also called (right distribution).}$$

Let  $R$  be the set of all real numbers, then multiplication distributes addition on  $R$ .

$$\text{Since, } a \cdot (b + c) = a \cdot b + a \cdot c, \forall a, b, c \in R.$$

**(iv) Identity Element** Let  $*$  be a binary operation on a non-empty set  $S$ . An element  $e \in S$ , if it exist such that

$$a * e = e * a = a, \forall a \in S.$$

is called an identity elements of  $S$ , with respect to  $*$ .

For addition on  $R$ , zero is the identity elements in  $R$ .

$$\text{Since, } a + 0 = 0 + a = a, \forall a \in R$$

For multiplication on  $\mathbb{R}$ , 1 is the identity element in  $\mathbb{R}$ .

Since,  $a \times 1 = 1 \times a = a, \forall a \in \mathbb{R}$

Let  $P(S)$  be the power set of a non-empty set  $S$ . Then,  $\Phi$  is the identity element for union on  $P(S)$  as

$A \cup \Phi = \Phi \cup A = A, \forall A \in P(S)$

Also,  $S$  is the identity element for intersection on  $P(S)$ .

Since,  $A \cap S = A \cap S = A, \forall A \in P(S)$ .

For addition on  $\mathbb{N}$  the identity element does not exist. But for multiplication on  $\mathbb{N}$  the identity element is 1.

(v) Inverse of an Element Let  $*$  be a binary operation on a non-empty set ' $S$ ' and let ' $e$ ' be the identity element.

Let  $a \in S$ . we say that  $a^{-1}$  is invertible, if there exists an element  $b \in S$  such that  $a * b = b * a = e$

Also, in this case,  $b$  is called the inverse of  $a$  and we write,  $a^{-1} = b$

Addition on  $\mathbb{N}$  has no identity element and accordingly  $\mathbb{N}$  has no invertible element.

Multiplication on  $\mathbb{N}$  has 1 as the identity element and no element other than 1 is invertible.

Let  $S$  be a finite set containing  $n$  elements. Then, the total number of binary operations on  $S$  is  $n^{n^2}$

Let  $S$  be a finite set containing  $n$  elements. Then, the total number of commutative binary operation on  $S$  is  $n \cdot [n(n+1)/2]$ .