

Study Material Downloaded from Vedantu

About **Vedantu**

Vedantu is India's largest **LIVE online teaching platform** with best teachers from across the country.

Vedantu offers Live Interactive Classes for **JEE**, **NEET**, KVPY, NTSE, Olympiads, **CBSE**, **ICSE**, IGCSE, IB & State Boards for Students Studying in **6-12th Grades** and Droppers.

HARSTER CLASSES REF Webinars by Expert Teachers The second seco

FREE LIVE ONLINE

Awesome Master Teachers



Anand Prakash B.Tech, IIT Roorkee Co-Founder, Vedantu



Pulkit Jain B.Tech, IIT Roorkee Co-Founder, Vedantu



Vamsi Krishna B.Tech, IIT Bombay Co-Founder, Vedantu



My mentor is approachable and **guides me** in my future aspirations as well.

Student **- Ayushi**



My son loves the sessions and **I can** already see the change. Parent - Sreelatha









95% Students of Regular Tuitions on Vedantu scored above **90%** in exams!

Vedantii FREE MASTER CLASS SERIES

- ⊘ For Grades 6-12th targeting JEE, CBSE, ICSE & much more
- Free 60 Minutes Live Interactive classes everyday
- Learn from the Master Teachers India's best

Register for **FREE**

Limited Seats!



Download Vedantu's App & Get



DOWNLOAD THE APP

EXERCISE NO: 4.1



Question 1:

Check whether the following are quadratic equations:

(i)
$$(x + 1)^2 = 2(x - 3)$$

(ii) $x^2 - 2x = (-2)(3 - x)$
(iii) $(x - 2)(x + 1) = (x - 1)(x + 3)$
(iv) $(x - 3)(2x + 1) = x(x + 5)$
(v) $(2x - 1)(x - 3) = (x + 5)(x - 1)$
(vi) $x^2 + 3x + 1 = (x - 2)^2$
(vii) $(x + 2)^3 = 2x(x^2 - 1)$
(viii) $x^3 - 4x^2 - x + 1 = (x - 2)^3$

Solution 1: (i) $(x+1)^2 = 2(x-3)$ $\Rightarrow x^2 + 2x + 1 = 2x - 6$ $\Rightarrow x^2 + 7 = 0$ It is of the form $ax^2 + bx + c = 0$. Hence, the given equation is a quadratic equation.

(ii)
$$x^2 - 2x = (-2)(3 - x)$$

 $\Rightarrow x^2 - 2x = -6 + 2x$
 $\Rightarrow x^2 - 4x + 6 = 0$
It is of the form $ax^2 + bx + a = 0$

It is of the form $ax^2 + bx + c = 0$. Hence, the given equation is a quadratic equation.

(iii)
$$(x-2)(x+1) = (x-1)(x+3)$$

$$\Rightarrow x^2 - x - 2 = x^2 + 2x - 3$$

$$\Rightarrow 3x - 1 = 0$$

It is not of the form $ax^2 + bx + c = 0$. Hence, the given equation is not a quadratic equation.



NCERT Solutions for Class 6 to 12 (Math & Science) Revision Notes for Class 6 to 12 (Math & Science) **RD Sharma Solutions for Class 6 to 12 Mathematics** RS Aggarwal Solutions for Class 6, 7 & 10 Mathematics Important Questions for Class 6 to 12 (Math & Science) CBSE Sample Papers for Class 9, 10 & 12 (Math & Science) Important Formula for Class 6 to 12 Math **CBSE Syllabus for Class 6 to 12** Lakhmir Singh Solutions for Class 9 & 10 **Previous Year Question Paper CBSE Class 12 Previous Year Question Paper CBSE Class 10 Previous Year Question Paper** JEE Main & Advanced Question Paper **NEET Previous Year Question Paper**

> Vedantu Innovations Pvt. Ltd. Score high with a personal teacher, Learn LIVE Online! www.vedantu.com



(iv) (x-3)(2x+1) = x (x+5) $\Rightarrow 2x^2 - 5x - 3 = x^2 + 5x$ $\Rightarrow x^2 - 10x - 3 = 0$

It is of the form $ax^2 + bx + c = 0$. Hence, the given equation is a quadratic equation.

(v) (2x-1)(x-3) = (x+5)(x-1) $\Rightarrow 2x^{2} - 7x + 3 = x^{2} + 4x - 5$ $\Rightarrow x^{2} - 11x + 8 = 0$ It is of the form $ax^{2} + bx + c = 0$. Hence, the given equation is a quadratic equation.

(vi) $x^2 + 3x + 1 = (x - 2)^2$ $\Rightarrow x^2 + 3x + 1 = x^2 + 4 - 4x$ $\Rightarrow 7x - 3 = 0$ It is not of the form $ax^2 + bx + c = 0$. Hence, the given equation is not a quadratic equation.

$$(vii) (x+2)^{3} = 2x(x^{2}-1)$$

⇒ x³+8+6x²+12x = 2x³ - 2x
⇒ x³-14x - 6x² - 8 = 0

It is not of the form $ax^2 + bx + c = 0$. Hence, the given equation is not a quadratic equation.

(viii) $x^{3} - 4x^{2} - x + 1 = (x - 2)^{3}$ $\Rightarrow x^{3} - 4x^{2} - x + 1 = x^{3} - 8 - 6x^{2} + 12x$ $\Rightarrow 2x^{2} - 13x + 9 = 0$

It is of the form $ax^2 + bx + c = 0$. Hence, the given equation is a quadratic equation.



Question 2:

Represent the following situations in the form of quadratic equations.

(i) The area of a rectangular plot is 528 m2. The length of the plot (in metres) is one more than twice its breadth. We need to find the length and breadth of the plot.

(ii) The product of two consecutive positive integers is 306. We need to find the integers.

(iii) Rohan's mother is 26 years older than him. The product of their ages (in years) 3 years from now will be 360. We would like to find Rohan's present age.

(iv) A train travels a distance of 480 km at a uniform speed. If the speed had been 8 km/h less, then it would have taken 3 hours more to cover the same distance. We need to find the speed of the train.

Solution 2:

(i) Let the breadth of the plot be x m. Hence, the length of the plot is (2x + 1) m. Area of a rectangle = Length × Breadth $\therefore 528 = x (2x + 1)$ $\Rightarrow 2x^2 + x - 528 = 0$

(ii) Let the consecutive integers be x and x + 1. It is given that their product is 306. $x + (x + 1) = 206 \Rightarrow x^2 + x = 206 = 0$

 $\therefore x(x+1) = 306 \Longrightarrow x^2 + x - 306 = 0$

(iii) Let Rohan's age be x. Hence, his mother's age = x + 263 years hence, Rohan's age = x + 3Mother's age = x + 26 + 3 = x + 29It is given that the product of their ages after 3 years is 360.



$$\therefore (x+3)(x+29) = 360$$
$$\Rightarrow x^2 + 32x - 273 = 0$$

(iv) Let the speed of train be x km/h. Time taken to travel 480 km = $\frac{480}{x}$ hrs

In second condition, let the speed of train = (x - 8) km/h It is also given that the train will take 3 hours to cover the same distance.

Therefore, time taken to travel 480 km = $\left(\frac{480}{x} + 3\right)$ hrs

Speed × Time = Distance

$$(x-8)\left(\frac{480}{x}+3\right) = 480$$

$$\Rightarrow 480+3x-\frac{3840}{x}-24 = 480$$

$$\Rightarrow 3x-\frac{3840}{x} = 24$$

$$\Rightarrow 3x^2-24x-3840 = 0$$

$$\Rightarrow x^2-8x-1280 = 0$$

EXERCISE NO: 4.2



Question 1:

Find the roots of the following quadratic equations by factorisation:

(i) $x^{2} - 3x - 10 = 0$ (ii) $2x^{2} + x - 6 = 0$ (iii) $\sqrt{2}x^{2} + 7x + 5\sqrt{2} = 0$ (iv) $2x^{2} - x + \frac{1}{8} = 0$ (v) $100x^{2} - 20x + 1 = 0$

Solution 1:

(i) $x^{2} - 3x - 10 = 0$ = $x^{2} - 5x + 2x - 10$ = x(x - 5) + 2(x - 5)= (x - 5)(x + 2)

Roots of this equation are the values for which (x - 5) (x + 2) = 0 $\therefore x - 5 = 0$ or x + 2 = 0i.e., x = 5 or x = -2

(ii) $2x^2 + x - 6 = 0$ = $2x^2 + 4x - 3x - 6$ = 2x(x+2) - 3(x+2)= (x+2)(2x-3)Roots of this equation are the values for which (x+2)(2x-3) = 0 $\therefore x + 2 = 0$ or 2x - 3 = 0i.e., x = -2 or $x = \frac{3}{2}$

(iii)
$$\sqrt{2x^2 + 7x + 5\sqrt{2}} = 0$$

= $\sqrt{2x^2 + 5x + 2x + 5\sqrt{2}}$
= $x(\sqrt{2x} + 5) + \sqrt{2}(\sqrt{2x} + 5)$
= $(\sqrt{2x} + 5)(x + \sqrt{2})$



Roots of this equation are the values for which $(\sqrt{2}x + 5)(x + \sqrt{2}) = 0$

$$\therefore \sqrt{2x + 5} = 0 \text{ or } x + \sqrt{2} = 0$$

i.e., $x = \frac{-5}{\sqrt{2}} \text{ or } x = -\sqrt{2}$
(iv) $2x^2 - x + \frac{1}{8} = 0$
 $= \frac{1}{8} (16x^2 - 8x + 1)$
 $= \frac{1}{8} (16x^2 - 4x - 4x + 1)$
 $= \frac{1}{8} (4x(4x - 1) - 1(4x - 1)))$
 $= \frac{1}{8} (4x - 1)^2$

Roots of this equation are the values for which $\frac{1}{8}(4x-1)^2 = 0$

Therefore
$$(4x - 1) = 0$$
 or $(4x - 1) = 0$,
i.e., $x = \frac{1}{4}$ or $x = \frac{1}{4}$
 $(v)100x^2 - 20x + 1 = 0$
 $= 100x^2 - 10x - 10x + 1$
 $= 10x(10x - 1) - 1(10x - 1)$
 $= (10x - 1)^2$

Roots of this equation are the values for which $(10x - 1)^2 = 0$ Therefore, (10x - 1) = 0 or (10x - 1) = 0i.e., $x = \frac{1}{10}$ or $x = \frac{1}{10}$

Question 2:

(i) John and Jivanti together have 45 marbles. Both of them lost 5



marbles each, and the product of the number of marbles they now have is 124. Find out how many marbles they had to start with.

(ii) A cottage industry produces a certain number of toys in a day. The cost of production of each toy (in rupees) was found to be 55 minus the number of toys produced in a day. On a particular day, the total cost of production was Rs 750. Find out the number of toys produced on that day.

Solution 2:

(i) Let the number of John's marbles be x. Therefore, number of Jivanti's marble = 45 - xAfter losing 5 marbles, Number of John's marbles = x - 5Number of Jivanti's marbles = 45 - x - 5 = 40 - xIt is given that the product of their marbles is 124. $\therefore (x - 5)(40 - x) = 124$ $\Rightarrow x^2 - 45x + 324 = 0$ $\Rightarrow x^2 - 36x - 9x + 324 = 0$ $\Rightarrow x(x - 36) - 9(x - 36) = 0$ $\Rightarrow (x - 36)(x - 9) = 0$

Either x - 36 = 0 or x - 9 = 0i.e., x = 36 or x = 9If the number of John's marbles = 36, Then, number of Jivanti's marbles = 45 - 36 = 9If number of John's marbles = 9, Then, number of Jivanti's marbles = 45 - 9 = 36

(ii) Let the number of toys produced be x. \therefore Cost of production of each toy = Rs (55 - x) It is given that, total production of the toys = Rs 750 $\therefore (55 - x)x = 750$ $\Rightarrow x^2 - 55x + 750 = 0$ $\Rightarrow x^2 - 25x - 30x + 750 = 0$



 $\Rightarrow x(x-25)-30(x-25)=0$ $\Rightarrow (x-25)(x-30)=0$ Either x - 25 = 0 or x - 30 = 0 i.e., x = 25 or x = 30 Hence, the number of toys will be either 25 or 30.

Question 3:

Find two numbers whose sum is 27 and product is 182.

Solution 3:

Let the first number be x and the second number is 27 - x. Therefore, their product = x (27 - x)It is given that the product of these numbers is 182. Therefore, x(27 - x) = 182 $\Rightarrow x^2 - 27x + 182 = 0$ $\Rightarrow x^2 - 13x - 14x + 182 = 0$ $\Rightarrow x (x - 13) - 14(x - 13) = 0$ $\Rightarrow (x - 13)(x - 14) = 0$

```
Either x - 13 = 0 or x - 14 = 0
i.e., x = 13 or x = 14
If first number = 13, then
Other number = 27 - 13 = 14
If first number = 14, then
Other number = 27 - 14 = 13
Therefore, the numbers are 13 and 14.
```

Question 4:

Find two consecutive positive integers, sum of whose squares is 365.

Solution 4:

Let the consecutive positive integers be x and x + 1. Given that $x^2 + (x + 1)^2 = 365$



 $\Rightarrow x^{2} + x^{2} + 1 + 2x = 365$ $\Rightarrow 2x^{2} + 2x - 364 = 0$ $\Rightarrow x^{2} + x - 182 = 0$ $\Rightarrow x^{2} + 14x - 13x - 182 = 0$ $\Rightarrow x(x + 14) - 13(x + 14) = 0$ $\Rightarrow (x + 14)(x - 13) = 0$ Either x + 14 = 0 or x - 13 = 0, i.e., x = -14 or x = 13Since the integers are positive, x can only be 13. $\therefore x + 1 = 13 + 1 = 14$ Therefore, two consecutive positive integers will be 13 and 14.

Question 5:

The altitude of a right triangle is 7 cm less than its base. If the hypotenuse is 13 cm, find the other two sides.

Solution 5:

Let the base of the right triangle be x cm. Its altitude = (x - 7) cm From Pythagoras theorem Base² + altitude² = Hypotenuse² $\therefore x^{2} + (x - 7)^{2} = 13^{2}$ $\Rightarrow x^{2} + x^{2} + 49 - 14x = 169$ $\Rightarrow 2x^{2} - 14x - 120 = 0$ $\Rightarrow x^{2} - 7x - 60 = 0$ $\Rightarrow x^{2} - 12x + 5x - 60 = 0$ $\Rightarrow x(x - 12) + 5(x - 12) = 0$ $\Rightarrow (x - 12)(x + 5) = 0$ Either x - 12 = 0 or x + 5 = 0, i.e., x = 12 or x = -5Since sides are positive, x can only be 12. Therefore, the base of the given triangle is 12 cm and the altitude of

this triangle will be (12 - 7) cm = 5 cm.



Question 6:

A cottage industry produces a certain number of pottery articles in a day. It was observed on a particular day that the cost of production of each article (in rupees) was 3 more than twice the number of articles produced on that day. If the total cost of production on that day was Rs 90, find the number of articles produced and the cost of each article.

Solution 6:

Let the number of articles produced be x. Therefore, cost of production of each article = Rs (2x + 3)It is given that the total production is Rs 90. $\therefore x(2x+3) = 90$ $\Rightarrow 2x^2 + 3x - 90 = 0$ $\Rightarrow 2x^2 + 15x - 12x - 90 = 0$ $\Rightarrow x(2x+15) - 6(2x+15) = 0$ $\Rightarrow (2x+15)(x-6) = 0$ Either 2x + 15 = 0 or x - 6 = 0, i.e., $x = \frac{-15}{2}$ or x = 6

As the number of articles produced can only be a positive integer, therefore, x can only be 6. Hence, number of articles produced = 6

Cost of each article = $2 \times 6 + 3 = \text{Rs } 15$

EXERCISE NO: 4.3



Question 1:

Find the roots of the following quadratic equations, if they exist, by the method of completing the square:

(i) $2x^{2} - 7x + 3 = 0$ (ii) $2x^{2} + x - 4 = 0$ (iii) $4x^{2} + 4\sqrt{3}x + 3 = 0$ (iv) $2x^{2} + x + 4 = 0$

Solution 1:

(i) $2x^2 - 7x + 3 = 0$ $\Rightarrow 2x^2 - 7x + 3 = 0$ On dividing both sides of the equation by 2, we obtain $\Rightarrow x^2 - \frac{7}{2}x = -\frac{3}{2}$ $\Rightarrow x^2 - 2 \times x \times \frac{7}{4} = -\frac{3}{2}$ On adding $\left(\frac{7}{4}\right)^2$ to both sides of equation, we obtain $\Rightarrow (x)^2 - 2 \times x \times \frac{7}{4} + \left(\frac{7}{4}\right)^2 = \left(\frac{7}{4}\right)^2 - \frac{3}{2}$ $\Rightarrow \left(x - \frac{7}{4}\right)^2 = \frac{49}{16} - \frac{3}{2}$ $\Rightarrow \left(x - \frac{7}{4}\right)^2 = \frac{25}{16}$ $\Rightarrow \left(x - \frac{7}{4}\right)^2 = \pm \frac{5}{4}$

$$\Rightarrow x = \frac{7}{4} \pm \frac{3}{4}$$

$$\Rightarrow x = \frac{7}{4} + \frac{5}{4} \text{ or } x = \frac{7}{4} - \frac{5}{4}$$
$$\Rightarrow x = \frac{12}{4} \text{ or } x = \frac{2}{4}$$



$$\Rightarrow$$
 x = 3 or $\frac{1}{2}$

(ii) $2x^2 + x - 4 = 0$ $\Rightarrow 2x^2 + x = 4$ On dividing both sides of the equation by 2, we obtain \Rightarrow x² + $\frac{1}{2}$ x = 2 On adding $\left(\frac{1}{4}\right)^2$ to both sides of the equation, we obtain $\Rightarrow (x)^{2} + 2 \times x \times \frac{1}{4} + \left(\frac{1}{4}\right)^{2} = 2 + \left(\frac{1}{4}\right)^{2}$ $\Rightarrow \left(x + \frac{1}{4}\right)^2 = \frac{33}{16}$ \Rightarrow x + $\frac{1}{4} = \pm \frac{\sqrt{33}}{4}$ \Rightarrow x = $\pm \frac{\sqrt{33}}{4} - \frac{1}{4}$ \Rightarrow x = $\frac{\pm\sqrt{33}-1}{4}$ \Rightarrow x = $\frac{\sqrt{33}-1}{4}$ or $\frac{-\sqrt{33}-1}{4}$ (iii) $4x^2 + 4\sqrt{3}x + 3 = 0$ $\Rightarrow (2x)^2 + 2 \times 2x \times \sqrt{3} + (\sqrt{3})^2 = 0$ $\Rightarrow \left(2x + \sqrt{3}\right)^2 = 0$ $\Rightarrow (2x + \sqrt{3}) = 0$ and $(2x + \sqrt{3}) = 0$ \Rightarrow x = $\frac{-\sqrt{3}}{2}$ and x = $\frac{-\sqrt{3}}{2}$



(iv) $2x^2 + x + 4 = 0$ $\Rightarrow 2x^2 + x = -4$ On dividing both sides of the equation by 2, we obtain $\Rightarrow x^2 + \frac{1}{2}x = -2$ $\Rightarrow x^2 + 2 \times x \times \frac{1}{4} = -2$ On adding $\left(\frac{1}{4}\right)^2$ to both sides of the equation, we obtain $\Rightarrow (x)^2 + 2 \times x \times \frac{1}{4} + \left(\frac{1}{4}\right)^2 = \left(\frac{1}{4}\right)^2 - 2$ $\Rightarrow \left(x + \frac{1}{4}\right)^2 = \frac{1}{16} - 2$ $\Rightarrow \left(x + \frac{1}{4}\right)^2 = -\frac{31}{16}$

However, the square of a number cannot be negative Therefore, there is no real root for the given equation.

Question 2:

Find the roots of the quadratic equations given in Q.1 above by applying the quadratic formula.

Solution 2:

(i) $2x^2 - 7x + 3 = 0$

On comparing this equation with $ax^2 + bx + c = 0$, we obtain a = 2, b = -7, c = 3

By using quadratic formula, we obtain

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$
$$\Rightarrow x = \frac{7 \pm \sqrt{49 - 24}}{4}$$
$$\Rightarrow x = \frac{7 \pm 5}{4}$$



$$\Rightarrow x = \frac{7 \pm 5}{4} \text{ or } \frac{7 - 5}{4}$$
$$\Rightarrow x = \frac{12}{4} \text{ or } \frac{2}{4}$$
$$\therefore x = 3 \text{ or } \frac{1}{2}$$

(ii) $2x^2 + x - 4 = 0$

On comparing this equation with $ax^2 + bx + c = 0$, we obtain a = 2, b = 1, c = -4By using quadratic formula, we obtain

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$\Rightarrow x = \frac{-1 \pm \sqrt{1 - 32}}{4}$$

$$\Rightarrow x = \frac{-1 \pm \sqrt{33}}{4}$$

$$\therefore x = \frac{-1 \pm \sqrt{33}}{4} \text{ or } \frac{-1 - \sqrt{33}}{4}$$

(iii)
$$4x^2 + 4\sqrt{3}x + 3 = 0$$

On comparing this equation with $ax^2 + bx + c = 0$, we obtain $a = 4, b = 4\sqrt{3}, c = 3$ By using quadratic formula, we obtain $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ $\Rightarrow x = \frac{-4\sqrt{3} \pm \sqrt{48 - 48}}{8}$ $\Rightarrow x = \frac{-4\sqrt{3} \pm \sqrt{48 - 48}}{8}$ $\Rightarrow x = \frac{-4\sqrt{3} \pm 0}{8}$ $\therefore x = \frac{-\sqrt{3}}{2}$ or $\frac{-\sqrt{3}}{2}$



(iv) $2x^2 + x + 4 = 0$

On comparing this equation with $ax^2 + bx + c = 0$, we obtain a = 2, b = 1, c = 4 By using quadratic formula, we obtain

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$
$$\Rightarrow x = \frac{-1 \pm \sqrt{1 - 32}}{4}$$
$$\Rightarrow x = \frac{-1 \pm \sqrt{-31}}{4}$$

However, the square of a number cannot be negative therefore; there is no real root for the given equation.

Question 3:

Find the roots of the following equations:

(i)
$$x - \frac{1}{x} = 3, x \neq 0$$
 (ii) $\frac{1}{x+4} - \frac{1}{x-7} = \frac{11}{30}, x \neq -4, 7$

Solution 3:

(i)
$$x - \frac{1}{x} = 3$$
, $x^2 - 3x - 1 = 0$

On comparing this equation with $ax^2 + bx + c = 0$, we obtain a = 1, b = -3, c = -1

By using quadratic formula, we obtain

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$\Rightarrow x = \frac{3 \pm \sqrt{9 + 4}}{2}$$

$$\Rightarrow x = \frac{3 \pm \sqrt{13}}{2}$$

Therefore, $x = \frac{3 \pm \sqrt{13}}{2}$ or $\frac{3 - \sqrt{13}}{2}$



(ii)
$$\frac{1}{x+4} - \frac{1}{x-7} = \frac{11}{30}$$

 $\Rightarrow \frac{x-7-x-4}{(x+4)(x-7)} = \frac{11}{30}$
 $\Rightarrow \frac{-11}{(x+4)(x-7)} = \frac{11}{30}$
 $\Rightarrow (x+4)(x-7) = -30$
 $\Rightarrow x^2 - 3x - 28 = -30$
 $\Rightarrow x^2 - 3x + 2 = 0$
 $\Rightarrow x^2 - 2x - x + 2 = 0$
 $\Rightarrow x(x-2) - 1(x-2) = 0$
 $\Rightarrow (x-2)(x-1) = 0$
 $\Rightarrow x = 1 \text{ or } 2$

Question 4:

The sum of the reciprocals of Rehman's ages, (in years) 3 years ago and 5 years from now is $\frac{1}{3}$. Find his present age.

Solution 4:

Let the present age of Rehman be x years. Three years ago, his age was (x - 3) years. Five years hence, his age will be (x + 5) years. It is given that the sum of the reciprocals of Rehman's ages 3 years

ago and 5 years from now is $\frac{1}{3}$

. 1 1	_1
$\frac{1}{x-3}$ $\frac{1}{x+5}$	$5\overline{3}$
x + 5 + x - 3	_1
$\overline{(x-3)(x+5)}$	$-\frac{1}{3}$
2x + 2	_ 1
$\overline{(x-3)(x+5)}$	$-\frac{1}{3}$



 $\Rightarrow 3(2x+2) = (x-3)(x+5)$ $\Rightarrow 6x + 6 = x^{2} + 2x - 15$ $\Rightarrow x^{2} - 4x - 21 = 0$ $\Rightarrow x^{2} - 7x + 3x - 21 = 0$ $\Rightarrow x(x-7) + 3(x-7) = 0$ $\Rightarrow (x-7)(x+3) = 0$ $\Rightarrow x = 7, -3$

However, age cannot be negative. Therefore, Rehman's present age is 7 years.

Question 5:

In a class test, the sum of Shefali's marks in Mathematics and English is 30. Had she got 2 marks more in Mathematics and 3 marks less in English, the product of their marks would have been 210. Find her marks in the two subjects.

Solution 5:

Let the marks in Maths be x. Then, the marks in English will be 30 - x. According to the given question, (x + 2)(30 - x - 3) = 210 (x + 2)(27 - x) = 210 $\Rightarrow -x^2 + 25x + 54 = 210$ $\Rightarrow x^2 - 25x + 156 = 0$ $\Rightarrow x^2 - 12x - 13x + 156 = 0$ $\Rightarrow x(x - 12) - 13(x - 12) = 0$ $\Rightarrow (x - 12)(x - 13) = 0$ $\Rightarrow x = 12, 13$ If the marks in mathematics are 12, then marks in English will be 30 - 12 = 18

If the marks in mathematics are 13, then marks in English will be 30 - 13 = 17



Question 6:

The diagonal of a rectangular field is 60 metres more than the shorter side. If the longer side is 30 metres more than the shorter side, find the sides of the field.

Solution 6:

Let the shorter side of the rectangle be x m. Then, larger side of the rectangle = (x + 30) m

Diagonal of the rectangle = $\sqrt{x^2 + (x + 30)^2}$

It is given that the diagonal of the rectangle is 60 m more than the shorter side.

$$\therefore \sqrt{x^{2} + (x + 30)^{2}} = x + 60$$

$$\Rightarrow x^{2} + (x + 30)^{2} = (x + 60)^{2}$$

$$\Rightarrow x^{2} + x^{2} + 900 + 60x = x^{2} + 3600 + 120x$$

$$\Rightarrow x^{2} - 60x - 2700 = 0$$

$$\Rightarrow x^{2} - 90x + 30x - 2700 = 0$$

$$\Rightarrow x(x - 90) + 30(x - 90)$$

$$\Rightarrow (x - 90)(x + 30) = 0$$

$$\Rightarrow x = 90, -30$$

However, side cannot be negative. Therefore, the length of the shorter side will be

90 m.

Hence, length of the larger side will be (90 + 30) m = 120 m

Question 7:

The difference of squares of two numbers is 180. The square of the smaller number is 8 times the larger number. Find the two numbers.

Solution 7:

Let the larger and smaller number be x and y respectively. According to the given question, $x^2 - y^2 = 180$ and $y^2 = 8x$

$$\Rightarrow x^2 - 8x = 180$$



 $\Rightarrow x^{2} - 8x - 180 = 0$ $\Rightarrow x^{2} - 18x + 10x - 180 = 0$ $\Rightarrow x (x - 18) + 10 (x - 18) = 0$ $\Rightarrow (x - 18) (x + 10) = 0$ $\Rightarrow x = 18, -10$

However, the larger number cannot be negative as 8 times of the larger number will be negative and hence, the square of the smaller number will be negative which is not possible.

Therefore, the larger number will be 18 only.

x = 18 $\therefore y^2 = 8x = 8 \times 18 = 144$ $\Rightarrow y = \pm \sqrt{144} = \pm 12$

 \therefore smaller number = ± 12

Therefore, the numbers are 18 and 12 or 18 and -12.

Question 8:

A train travels 360 km at a uniform speed. If the speed had been 5 km/h more, it would have taken 1 hour less for the same journey. Find the speed of the train.

Solution 8:

Let the speed of the train be x km/hr.

Time taken to cover 360 km hr = $\frac{360}{100}$ hr

According to the given question,

$$(x+5)\left(\frac{360}{x}-1\right) = 360$$
$$\Rightarrow 360 - x + \frac{1800}{x} - 5 = 360$$



 \Rightarrow x² + 5x - 1800 = 0 \Rightarrow x² + 45x - 40x - 1800 = 0 $\Rightarrow x(x+45)-40(x+45)=0$ $\Rightarrow (x+45)(x-40) = 0$ \Rightarrow x = 40, -45 However, speed cannot be negative. Therefore, the speed of train is 40 km/h

Question 9:

Two water taps together can fill a tank in $9\frac{3}{8}$ hours. The tap of larger diameter takes 10 hours less than the smaller one to fill the tank separately. Find the time in which each tap can separately fill the tank.

Solution 9:

Let the time taken by the smaller pipe to fill the tank be x hr. Time taken by the larger pipe = (x - 10) hr

Part of tank filled by smaller pipe in 1 hour = $\frac{1}{2}$ Part of tank filled by larger pipe in 1 hour = $\frac{1}{x-10}$

It is given that the tank can be filled in $9\frac{3}{8} = \frac{75}{8}$ hours by both the pipes

together. Therefore,

$$\frac{1}{x} + \frac{1}{x-10} = \frac{8}{75}$$
$$\frac{x-10+x}{x(x-10)} = \frac{8}{75}$$
$$\Rightarrow \frac{2x-10}{x(x-10)} = \frac{8}{75}$$



 $\Rightarrow 75(2x-10) = 8x^{2} - 80x$ $\Rightarrow 150x - 750 = 8x^{2} - 80x$ $\Rightarrow 8x^{2} - 230x + 750 = 0$ $\Rightarrow 8x^{2} - 200x - 30x + 750 = 0$ $\Rightarrow 8x(x - 25) - 30(x - 25) = 0$ $\Rightarrow (x - 25)(8x + 30) = 0$ i.e., x = 25, $\frac{-30}{8}$

Time taken by the smaller pipe cannot be $\frac{-30}{8} = -3.75$ hours. As in this

case, the time taken by the larger pipe will be negative, which is logically not possible.

Therefore, time taken individually by the smaller pipe and the larger pipe will be 25 and 25 - 10 = 15 hours respectively.

Question 10:

An express train takes 1 hour less than a passenger train to travel 132 km between Mysore and Bangalore (without taking into consideration the time they stop at intermediate stations). If the average speeds of the express train is 11 km/h more than that of the passenger train, find the average speed of the two trains.

Solution 10:

Let the average speed of passenger train be x km/h. Average speed of express train = (x + 11) km/h It is given that the time taken by the express train to cover 132 km is 1 hour less than the passenger train to cover the same distance.

$$\therefore \frac{132}{x} - \frac{132}{x+11} = 1$$
$$\Rightarrow 132 \left[\frac{x+11-x}{x(x+11)} \right] = 1$$
$$\Rightarrow \frac{132 \times 11}{x(x+11)} = 1$$



 $\Rightarrow 132 \times 11 = x(x+11)$ $\Rightarrow x^{2} + 11x - 1452 = 0$ $\Rightarrow x^{2} + 44x - 33x - 1452 = 0$ $\Rightarrow x(x+44) - 33(x+44) = 0$ $\Rightarrow (x+44)(x-33) = 0$ $\Rightarrow x = -44, 33$ Speed cannot be negative.

Therefore, the speed of the passenger train will be 33 km/h and thus, the speed of the express train will be 33 + 11 = 44 km/h.

Question 11:

Sum of the areas of two squares is 468 m2. If the difference of their perimeters is 24 m, find the sides of the two squares.

Solution 11:

Let the sides of the two squares be x m and y m. Therefore, their perimeter will be 4x and 4y respectively and their areas will be x^2 and y^2 respectively.

It is given that

$$4x - 4y = 24$$

$$x - y = 6$$

$$x = y + 6$$
Also, $x^2 + y^2 = 468$

$$\Rightarrow (6 + y)^2 + y^2 = 468$$

$$\Rightarrow 36 + y^2 + 12y + y^2 = 468$$

$$\Rightarrow 2y^2 + 12y - 432 = 0$$

$$\Rightarrow y^2 + 6y - 216 = 0$$

$$\Rightarrow y^2 + 18y - 12y - 216 = 0$$

$$\Rightarrow y(y + 18) - 12(y + 18) = 0$$

$$\Rightarrow (y + 18)(y - 12) = 0$$

$$\Rightarrow y = -18 \text{ or } 12$$
However, side of a square cannot be negative.

Hence, the sides of the squares are 12 m and (12 + 6) m = 18 m

EXERCISE NO: 4.4



Question 1:

Find the nature of the roots of the following quadratic equations. If the real roots exist, find them;

- I. $2x^2 3x + 5 = 0$ II. $3x^2 - 4\sqrt{3x} + 4 = 0$
- 11. $3x 4\sqrt{3x} + 4 = 0$
- III. $2x^2 6x + 3 = 0$

Solution 1:

We know that for a quadratic equation $ax^2 + bx + c = 0$

 $Discriminant = b^2 - 4ac$

(A) If $b^2 - 4ac > 0 \rightarrow two distinct real roots$ $(B) If <math>b^2 - 4ac = 0 \rightarrow two equal real roots$

(C) If $b^2 - 4ac < 0 \rightarrow$ no real roots

(I) $2x^2 - 3x + 5 = 0$

Comparing this equation with $ax^2 + bx + c = 0$, we obtain a = 2, b = -3, c = 5Discriminant : $b^2 - 4ac$ $= (-3)^2 - 4 (2) (5) = 9 - 40$ = -31As $b^2 - 4ac < 0$, Therefore, no real root is possible for the given equation.

(II) $3x^2 - 4\sqrt{3x} + 4 = 0$ Comparing this equation with $ax^2 + bx + c = 0$, we obtain $a = 3, b = -4\sqrt{3}, c = 4$ Discriminant : $b^2 - 4ac = (-4\sqrt{3})^2 - 4(3)(4)$ = 48 - 48 = 0As $b^2 - 4ac = 0$,

Therefore, real roots exist for the given equation and they are equal to each other.

And the roots will be
$$\frac{-b}{2a}$$
 and $\frac{-b}{2a}$.



$$\frac{-b}{2a} = \frac{-(-4\sqrt{3})}{2\times3} = \frac{-4\sqrt{3}}{6} = \frac{2\sqrt{3}}{3} = \frac{2}{\sqrt{3}}$$

Therefore, the roots are $\frac{2}{\sqrt{3}}$ and $\frac{2}{\sqrt{3}}$.

(III) $2x^2 - 6x + 3 = 0$ Comparing this equation with $ax^2 + bx + c = 0$, we obtain a = 2, b = -6, c = 3Discriminant $= b^2 - 4ac = (-6)^2 - 4$ (2) (3) = 36 - 24 = 12As $b^2 - 4ac > 0$, Therefore, distinct real roots exist for this equation as follows. $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ $= \frac{-(-6) \pm \sqrt{(-6)^2 - 4(2)(3)}}{2(2)}$

$$= \frac{6 \pm \sqrt{12}}{4} = \frac{6 \pm 2\sqrt{3}}{4}$$
$$= \frac{3 \pm \sqrt{3}}{2}$$
Therefore, the root are $\frac{3 + \sqrt{3}}{2}$ or $\frac{3 - \sqrt{3}}{2}$

Question 2:

Find the values of k for each of the following quadratic equations, so that they have two equal roots.

- I. $2x^2 + kx + 3 = 0$
- II. kx(x-2) + 6 = 0

Solution 2:

We know that if an equation $ax^2 + bx + c = 0$ has two equal roots, its Discriminant ($b^2 - 4ac$) will be 0.

(I) $2x^2 + kx + 3 = 0$ Comparing equation with $ax^2 + bx + c = 0$, we obtain



a = 2, b = k, c = 3 Discriminant = b² - 4ac = (k)² - 4(2) (3) = k² - 24 For equal roots, Discriminant = 0 k² - 24 = 0 k² = 24 k = $\pm \sqrt{24} = \pm 2\sqrt{6}$

(II) kx (x - 2) + 6 = 0or $kx^2 - 2kx + 6 = 0$ Comparing this equation with $ax^2 + bx + c = 0$, we obtain a = k, b = -2k, c = 6Discriminant $= b^2 - 4ac = (-2k)^2 - 4$ (k) (6) $= 4k^2 - 24k$ For equal roots, $b^2 - 4ac = 0$ $4k^2 - 24k = 0$ $4k^2 - 24k = 0$ 4k (k - 6) = 0Either 4k = 0 or k = 6 = 0k = 0 or k = 6However, if k = 0, then the equation will not have the terms 'x²' and 'x'. Therefore, if this equation has two equal roots, k should be 6 only.

Question 3:

Is it possible to design a rectangular mango grove whose length is twice its breadth, and the area is 800 m^2 ? If so, find its length and breadth.

Solution 3:

Let the breadth of mango grove be *l*. Length of mango grove will be 2*l*. Area of mango grove = $(2l) (l) = 2l^2$ $2l^2 = 800$

$$1^2 = \frac{800}{2} = 400$$
$$1^2 = 400 = 0$$



Comparing this equation with $al^2 + bl + c = 0$, we obtain a = 1 b = 0, c = 400Discriminant $= b^2 - 4ac = (0)^2 - 4 \times (1) \times (-400) = 1600$ Here, $b^2 - 4ac > 0$ Therefore, the equation will have real roots. And hence, the desired rectangular mango grove can be designed. $l = \pm 20$ However, length cannot be negative. Therefore, breadth of mango grove = 20 m Length of mango grove $= 2 \times 20 = 40$ m

Question 4:

Is the following situation possible? If so, determine their present ages. The sum of the ages of two friends is 20 years. Four years ago, the product of their ages in years was 48.

Solution 4:

Let the age of one friend be x years. Age of the other friend will be (20 - x) years. 4 years ago, age of 1st friend = (x - 4) years And, age of 2^{nd} friend = (20 - x - 4)=(16 - x) years Given that, (x-4)(16-x) = 48 $16x - 64 - x^2 + 4x = 48$ $-x^{2} + 20x - 112 = 0$ $x^2 - 20x + 112 = 0$ Comparing this equation with $ax^2 + bx + c = 0$, we obtain a = 1, b = -20, c = 112Discriminant = $b^2 - 4ac = (-20)^2 - 4(1)(112)$ =400 - 448 = -48As $b^2 - 4ac < 0$. Therefore, no real root is possible for this equation and hence, this situation is not possible.

Question 5:

Is it possible to design a rectangular park of perimeter 80 and area 400 m²? If so find its length and breadth.



Solution 5:

Let the length and breadth of the park be 1 and b. Perimeter = 2(1 + b) = 80 1 + b = 40Or, b = 40 - 1Area = $1 \times b = 1(40 - 1) = 401 - 1^2$ $401 - 1^2 = 400$ $1^2 - 401 + 400 = 0$ Comparing this equation with $al^2 + bl + c = 0$, we obtain a = 1, b = -40, c = 400Discriminate = $b^2 - 4ac = (-40)^2 - 4(1)(400)$ = 1600 - 1600 = 0As $b^2 - 4ac = 0$, Therefore, this equation has equal real roots. And hence, this situation is possible.

Root of this equation,

$$l = -\frac{b}{2a}$$

$$l = -\frac{(-40)}{2(1)} = \frac{40}{2} = 20$$

Therefore, length of park, l = 20 mAnd breadth of park, b = 40 - 1 = 40 - 20 = 20 m



Thank YOU for downloading the PDF

FREE LIVE ONLINE

MASTER CLASSES FREE Webinars by Expert Teachers



Vedantii FREE MASTER CLASS SERIES

- ⊘ For Grades 6-12th targeting JEE, CBSE, ICSE & much more
- Since Free 60 Minutes Live Interactive classes everyday
- ⊘ Learn from the Master Teachers India's best

Register for **FREE**

Limited Seats!