

NCERT SOLUTIONS
CLASS-XI CHEMISTRY
CHAPTER-4
CHEMICAL BONDING & MOLECULAR STRUCTURE

Q-1) Explain Chemical bond formation process.

Ans.)

"Chemical bond is an attractive force that bounds the constituents of a chemical species together."

So many theories are suggested for chemical bond formation such as valence shell electron pair repulsion theory, electronic theory, molecular orbital theory, and valence bond theory.

Formation of chemical bond is credited to the tendency of system to achieve stability. It was noticed that inertness of noble gases is direct result of their completely filled outermost orbitals. Consequently, it was proposed that the elements having deficiency of electrons in outermost shells are unstable. Thus, atoms combine with one another and finish their separate octets or duplets to achieve the stable configuration of the closest inert gases. So, this combination may occur either by sharing of electrons. The formed chemical bond is a result of sharing of electrons among atoms is known as a covalent bond. Also a formed ionic bond is a result of sharing of electrons among atoms.

Q-2) Give the Lewis dot symbols for the elements given below:

a) Na

b) Mg

c) O

d) N

e) B

f) Br

Ans.)

a) Na

Sodium atom contains only 1 valence electron. Thus, the Lewis dot symbols for Na is $\text{Na}\cdot$.

b) Mg

Magnesium atom contains only 2 valence electrons. Thus, the Lewis dot symbols for Mg is



c) O

Oxygen atom contains only 6 valence electrons. Thus, the Lewis dot symbols for O is



d) N

Nitrogen atom contains only 5 valence electrons. Thus, the lewis dot symbols for N is



e) B

Boron atom contains only 3 valence electrons. Thus, the lewis dot symbols for B is



f) Br

Bromine atom contains only 7 valence electrons. Thus, the lewis dot symbols for Br is



Q-3) Give the lewis dot symbols for the elements given below:

(a) Al and Al^{3+}

(b) H and H^-

(c) S and S^{2-}

Ans.)

(a) Al and Al^{3+}

Aluminium atom contains only 3 valence electrons. Thus, the lewis dot symbols for Al is



The tri-positive charge on aluminium indicates that it has donated 3 electrons.

Thus lewis dot symbol is




(b) H and H^-

Hydrogen atom contains only 1 valence electrons. Thus, the lewis dot symbols for H is



The single negative charge on hydrogen indicates that it has gained 1 electron. So, one valance electron plus one gained electron.

Thus lewis dot symbol is




(c) S and S^{2-}

Sulphur atom contains only 6 valence electrons. Thus, the lewis dot symbols for S is



The bi-negative charge on sulphur indicates that it has gained 2 electron. So, six valance electron plus two gained electron.

Thus lewis dot symbol is



Q-4) Give the lewis structure for the ions and molecules given below:

(a) SiCl_4

(b) H_2S


(c) HCOOH

(d) BeF_2


(e) CO_3^{2-}

Ans.)


(a) SiCl_4




(b) H_2S




(c) HCOOH



(d) BeF_2




(e) CO_3^{2-}



Q-5) Write the definition of octet rule. State the significance and limitation of octet rule.

Ans.)

Octet rule says, "atoms can combine either by transfer of valence electrons from one atom to another or by sharing their valence electrons in order to achieve the nearest inert gas configuration by having an octet in their valence shell."



Octet rule explains chemical bond formation depending upon nature of element.

Limitations:


(a) Octet rule fails to predict the relative stability and shape of the molecules.

(b) It is based on inert nature of noble gases. But, some inert gases say, krypton(Kr) and xenon(Xe) form compounds like KrF_2 , XeF_2 etc.


(c) For elements beyond 3rd period the octet rule cannot be applied. Elements present beyond 3rd period have more than 8 valence electrons surrounding central atom. E.g. SF_6 , PF_6 etc.



(d) For atoms in a molecule having odd number of electrons octet rule is not applied. E.g. For NO_2 and NO octet rule is not applicable.



(e) If an compound is having less than 8 electrons surrounding the central atom then octet rule cannot be applied to that compound. E.g. BeH_2 , $AlCl_3$, $LiCl$ etc. is not obeying octet rule.



Q-6) State the factors that are favourable for ionic bond formation.

Ans.)

Formation of ionic bond takes place by transfer of 1 or more electrons from one atom to another. Thus, ionic bond formation depends on the flexibility of neutral atoms to lose or gain electrons. Formation of ionic bond also depends on the lattice energy of compound which is formed.

The factors that are favourable for ionic bond formation:

- (a) High electron affinity of atoms of non- metal.
- (b) High lattice energy of compound which is formed.
- (c) Low ionization enthalpy of atom of metal.


Q-7) Comment about the shape of the molecules given below by using VSEPR model:

- (a) BCl_3
- (b) SiCl_4
- (c) AsF_5
- (d) PH_3
- (e) BeCl_2
- (f) H_2S

Ans.)


- (a) BCl_3

Central atom is not having any lone pair but have 3 bond pairs. Thus, its shape is AB_3 . i.e..Trigonal planar.



- (b) SiCl_4

Central atom is not having any lone pair but have 4 bond pairs. Thus, its shape is AB_4 . i.e. Tetrahedral.



- (c) AsF_5

Central atom is not having any lone pair but have 5 bond pairs. Thus, its shape is

AB₅. i.e. Trigonalbipyramidal.




(d) PH₃

Central atom is having 1 lone pair and is having 3 bond pairs. Thus, its shape is AB₃E. i.e. Trigonalbipyramidal.



(e) BeCl₂

Central atom is not having any lone pair but have 2 bond pairs. Thus, its shape is AB₂. i.e. Linear shape.



(f) H₂S


Central atom is having 1 lone pair and is having 2 bond pairs. Thus, its shape is AB₂E. i.e. Bent shape.



Q-8) Explain why the bond angle in H₂O is less than NH₃, even though they are having distorted tetrahedral structure.

Ans.)

The geometry of H₂O and NH₃:



Central atom(N) in ammonia is having 1 lone pair and is having 3 bond pairs.

Central atom(O) in water is having 2 lone pair and is having 2 bond pairs.

Thus, these 2 lone pairs on O- atom in water molecule repels the 2 bond pairs. And this repulsion is between lone pair and bond pair on O- atom of H₂O is stronger than the repulsion is between lone pair and bond pair on N-atom of NH₃.

Thus, the bond angle in H₂O is less than NH₃, even though they are having distorted tetrahedral structure.

Q-9) Express, 'bond strength in terms of bond order'.

Ans.)

The extent of bonding which occurs between two atoms while forming a molecule is represented by bond strength. As the bond strength increases the bond becomes stronger and the bond order increases.

Q-10) Define: Bond length.

Ans.)

"Bond length is defined as the equilibrium distance between the nuclei of 2 bonded atoms in a molecule."


Q-11) Explain important aspects of the resonance with reference to CO_3^{2-} ion.

Ans.)

Experimental results shows that, all the C-O bond in CO_3^{2-} are equivalent.

Thus, it is inefficient to represent CO_3^{2-} ion by single lewis structure which is having 1 double bond and 2 single bond.


Thus, the resonance structures of CO_3^{2-} is :



Q-12) H_3PO_3 can be represented by following structures.

Can we take these 2 structures as canonical form of resonance hybrid which is representing H_3PO_3 ? Explain.

Ans.)



In the given structures the position of atoms are changed, so we cannot take the 2 given structures as canonical form of resonance hybrid which is representing H_3PO_3 .

Q-13) Give the resonance structures of the following ions


(a) NO_3^-

(b) NO_2


(c) SO_3

Ans.)


(a) NO_3^-



(b) NO_2



(c) SO_3



Q-14) Show transfer of electrons between the atoms given below to form anions and cation.

(i) O and Ca

(ii) N and Al

(iii) S and K.

Ans.)


(i) O and Ca

Electronic configurations of O and Ca are:

O: 2, 6

Ca: 2, 8, 8, 2

Here, it is clear that Ca has 2 more electrons than nearest inert gas, i.e. Ar, whereas O needs 2 electrons to complete its octet. Thus, the transfer of electrons takes place in the following way,




(ii) N and Al

Electronic configurations of N and Al are:

N: 2, 5

Al: 2, 8, 3

Here, it clear that Al has 3 more electrons then nearest inert gas. i.e. Ne, whereas N needs 3 electrons to complete its octet. Thus, the transfer of electrons takes place in following way,




(iii) S and K

Electronic configurations of S and K are:

S: 2, 8, 6

K: 2, 8, 8, 1

Here, it clear that K has 1 more electrons then nearest inert gas. i.e. Ne, whereas S needs 2 electrons to complete its octet. Thus, the transfer of electrons takes place in following way,




Q-15) H_2O and CO_2 are tri-atomic molecules, then also the shape of H_2O is bent and that of shape of CO_2 is linear. Why?

Explain with the help of dipole moment.


Ans.)

Experimental results show that the dipole moment of CO_2 is 0. And it is possible only if the shape of molecule is linear as dipole moments of bond between C-O is equal and opposite so, it nullifies each other.



$\therefore$ Resultant, $\mu = 0$

H_2O has 1.84D dipole moment. The value of dipole moments indicates that structure of water molecule is bent as dipole moments of bond between O-H is unequal.



Q-16) Give the significance of dipole moment.

Ans.)

In heteronuclear molecule, there is difference in electro-negativities of constituents of atom, due to which polarisation arises in it. Thus, one end of acquires positive charge and other end acquires negative charge. So, molecules are said to have a dipole moment.


The product of distance between centres of negative and positive charges and magnitude of charge is known as dipole moment (μ). Dipole moment is a vector quantity and it is represented by an arrow in such a way that head of a tail is pointing towards negative centre and tail of arrow is pointing towards positive centre.

μ = distance between the centres (r) * charge (Q)

SI unit of dipole moment is 'esu'.

$$1 \text{ esu} = 3.335 \times 10^{-30} \text{ cm}$$

It is the measure of polarity of a bond. It differentiates the non-polar and polar bonds as all the non-polar molecules have dipole moment equal to 0. It also calculates the ionic character of a molecule.



Q-17) Give definition of electronegativity. State difference between electronegativity and electron affinity.

Ans.)

"Electronegativity is the ability of an atom in a chemical compound to attract a bond pair of electrons towards itself".

Sr. No	Electronegativity	Electron affinity
1	Tendency to attract the shared pairs of electrons for an atom which is in chemical compound is its electronegativity.	Tendency to gain electrons for an isolated gaseous atom is its electron gain enthalpy.
2	It varies according to element to which it is bounded with.	It does not vary according to element to which it is bounded with.
3	It is not constant for any element.	It is constant for an element.
4	It is not a measurable quantity.	It is a measurable quantity.

Q-18) Explain polar covalent bond with suitable example.


Ans.)

When two unique atoms having distinct electronegativities join to form a covalent bond, the bond pair of electrons is not shared equally. The nucleus of an atom having greater electronegativity attracts the bond pair. So, the electron distribution gets distorted and an electronegativity atom attracts the electron cloud.

Thus, the electronegative element gets slightly negatively charged and on the

Thus, the electronegative element gets slightly negatively charged and on the other hand, the other atom gets slightly positively charged. As a result of this, two opposite poles are developed in a molecule and this type of bond formed is termed as 'polar covalent bond'.

E.g. HCl is having a polar covalent bond. In HCl, Cl- atom is having more electronegativity than H- atom. Thus, bond pair shifts towards Cl- atom and because of that it acquires positive charge.



Q-19) For molecules given below, arrange the bonds in increasing order of their ionic characteristic.

K₂O, N₂, ClF₃, LiF, SO₂

Ans.)


Ionic characteristic of a molecule depends on the difference in electronegativity between constituents atoms. So, higher the difference, the ionic characteristic of a molecule will be higher.

So, the required order of ionic characteristic of the given molecules is


$\text{N}_2 < \text{SO}_2 < \text{ClF}_3 < \text{K}_2\text{O} < \text{LiF}$.

Q-20) Some of the bonds are incorrectly shown in the skeletal structure of CH₃COOH. Give the correct Lewis structure of CH₃COOH.

Ans.)




Correct Lewis structure of CH₃COOH is given below:




Q-21) CH₄ can also have square planar geometry other than tetrahedral geometry, and in that geometry all the 4 H- atoms are at the corner of the square and C- atom is at the centre. Give reason why Methane's geometry is not square planar.

Ans.)


Electronic configuration of C- atom:



Orbital picture of C- atom in excited state is:



Thus, C- atom undergoes sp^3 hybridization in methane molecule and forms tetrahedral structure.



For square planer geometry, C-atom should have dsp^2 hybridization. But as C-atom is not having d- orbital so it cannot undergo dsp^2 hybridization. Thus, Methane's geometry cannot be square planer.

Also in square planar geometry the bond angle is 90° so the stability is not there because of repulsion between bond pairs. So as per VSEPR theory methane's tetrahedral structure is perfect.


Q-22) Even though Be- H bonds are polar In BeH_2 , the dipole moment of BeH_2 is 0. Why is it so?

Ans.)

Lewis structure of BeH_2 is:



Central atom is not having any lone pair but have 2 bond pairs. Thus, its shape is AB_2 . i.e. Linear shape.



Thus, the dipole moment of Be- H bond is equal and opposite in direction so it nullifies one another. Thus, the dipole moment of BeH_2 is 0.

Q-23) Which is having higher dipole moment out of NF_3 and NH_3 ?

Ans.)


N- atom is the central atom of NF_3 and NH_3

N-atom is the central atom of NF_3 and NH_3 .

Central atom is having 1 lone pair and is having 3 bond pairs. Thus, for both the shape is AB_3E . i.e. Pyramidal.

As, F-atom is more electronegativity than H-atom, NF_3 should have higher dipole moment than NH_3 . But the dipole moment of NH_3 is 1.46D which is higher than dipole moment of NF_3 which is 0.24D.

It gets clear from the directions of dipole moments of individual bond in NF_3 and NH_3 .



As, both the N-H bond are in same direction it adds to the bond moment of the lone pair, while N-F bond are in opposite direction so they partly cut the bond moment of lone pair.

Thus, dipole moment of NH_3 is higher than that of NF_3 .

Q-24) Explain 'hybridisation of atomic orbitals'. Draw sp , sp^2 , sp^3 hybrid orbitals.


Ans.)

"Hybridization is defined as an intermixing of a set of atomic orbitals of slightly different energies, thereby forming a new set of orbitals having equivalent energies and shapes".

E.g. 1 s-orbital hybridises with 3 p-orbitals to form 4 sp^3 hybrid orbitals.


(a) sp hybrid orbital

1 s-orbital hybridises with 1 p-orbital to form 2 sp hybrid orbitals. sp hybrid orbital is having linear shape. The formation of sp orbital is:




(b) sp^2 hybrid orbital

1 s-orbital hybridises with 2 p-orbitals to form 3 sp^2 hybrid orbitals. The shape of sp^2 orbital is trigonal planar.




(c) sp^3 hybrid orbital

1 s- orbital hybridises with 3 p- orbitals to form 4 sp^3 hybrid orbitals. The shape of sp^3 orbital is tetrahedron.




Q-25) Whether the hybridisation of Al- atom change for the reaction given below?




Ans.)

The ground state of valence orbital of Al -atom is:




In excited state the orbital picture of Al- atom is:



Thus, Al -atom in $AlCl_3$ undergoes sp^2 hybridisation and forms trigonal planar geometry. For the formation of $AlCl_4^-$ the vacant $3p_z$ orbital will also get involved. Thus, sp^2 hybridisation is converted into sp^3 hybridisation and forms a tetrahedral structure.


Q-26) Determine that there is any change in hybridisation of N-Atom and B-atom because of the reaction given below:



Ans.)

N- atom in NH_3 is having sp^3 hybridization. Orbital picture of N- atom is shown below:





B- atom in NF_3 is having sp^2 hybridisation . Orbital picture of B- atom is shown below:



On the reaction of NH_3 and BF_3 , $\text{F}_3\text{B.NH}_3$ is obtained as product, as hybridization of B-atom is changed to sp^3 . Although, hybridization of N-atoms remains unchanged.

Q-27) Explain the formation of double and triple bond in C_2H_2 and C_2H_4 with the help of diagrams.


Ans.)

C_2H_2

In formation of ethyne(C_2H_2) molecule, C- atom is having sp hybridization with 2 $2p$ - orbitals in unhybridized state.


1 sp orbital of each C- atom overlaps the inter-nuclear axis and forms C-C sigma bond. The 2^{nd} sp orbital of each C- atom overlaps half-filled $1s$ orbital so as to form a sigma bond.

The triple bond between the 2 C- atoms has 1 sigma bond and 2 Pie bonds. This is because 2 unhybridized $2p$ - orbitals overlaps with the $2p$ - orbital of other C- atom, thus forming 2 pie bonds.



C_2H_4


Electronic configuration of carbon atom in excited state is given below:



In the formation of C_2H_4 (ethane) molecule 1 sp^2 orbital of C- atom overlaps sp^2 orbital of other C- atom. Thus, forming a C-C sigma bond.

The 2 remaining sp^2 orbital of every C- atom forms sp^2 -s σ bond with 2 H- atoms.

One c -atom having unhybridized orbital overlaps with the unhybridized orbital of other C -atom and forms a π bond.



Q-28) Find out the number of π and σ bonds in the molecules given below:

(i) C_2H_4

(ii) C_2H_2

Ans.)

Single bond is formed as the axis of bonding orbital overlaps. Thus, it forms a σ bond. By sidewise overlapping of orbital double and triple bonds i.e. multiple bonds are formed. π bond is always present in the multiple bonds. Triple bond consist of 2 π and 1 σ bond.


(i) C_2H_2



Thus, there are 2 π bonds and 3 σ in C_2H_2 .

(ii) C_2H_4

Thus, there are 1 π bonds and 5 σ in C_2H_4 .



Q-29) Take x- axis as inter-nuclear axis and find which of the orbital given below will not form σ bond and why?

(i) $1s$ and $2p_x$

(ii) $1s$ and $1s$

(iii) $1s$ and $2s$

(iv) $2p_y$ and $2p_y$

Ans.)

(iv) $2p_y$ and $2p_y$

$2p_y$ and $2p_y$ orbitals won't form a σ as it will undergo lateral overlapping and will form a π bond.

Q-30) Give the hybrid orbitals used by the C- atoms in the molecules given below:

(i) $\text{CH}_3\text{-CH=CH}_2$

(ii) $\text{CH}_3\text{-CH}_3$


(iii) $\text{CH}_3\text{-CHO}$

(iv) CH_3COOH

(v) $\text{CH}_3\text{CH}_2\text{-OH}$


Ans.)

(i) $\text{CH}_3\text{-CH=CH}_2$




Here, C_3 and C_2 are having sp^2 hybridization and C_1 is having sp^3 hybridization.

(ii) $\text{CH}_3\text{-CH}_3$




Here, C_1 and C_2 are having sp^3 hybridization.

(iii) $\text{CH}_3\text{-CHO}$




Here, C_1 is having sp^3 hybridization and C_2 is having sp^2 hybridization.

(iv) CH_3COOH



Here, C_1 is having sp^3 hybridization and C_2 is having sp^2 hybridization.

(v) $\text{CH}_3\text{CH}_2\text{-OH}$



Here, C_1 and C_2 are having sp^3 hybridization.


Q-31) Explain lone pairs and bond pairs of electrons with an example.

Ans.)


A covalent bond is formed when 2 atoms combine with each other by sharing their valence electrons.

"The shared pairs of electrons present between the bonded atoms are called bond pairs". Each and every electron cannot participate in bonding. "The pairs of electrons which do not participate in bonding are called lone pairs".

E.g. a) Ethane is having 7 bond pairs but zero lone pair.



b) Water is having 2 bond pairs and 2 lone pairs on O-atom.



Q-32) State difference between pi bond and sigma bond.

Ans.)

Sr. No.	Pi bond	Sigma bond
	π bond is formed by lateral overlap of p orbitals.	σ bond is formed by head-on overlap of orbitals.

1	Pi bond is formed by lateral overlapping of orbitals.	Sigma bond is formed by end to end overlapping of orbitals.
2	It is comparatively weak bond.	It is comparatively strong bond.
3	There is only one overlapping orbital is p-p.	The overlapping orbitals are s-s, s-p, p-p.
4	Rotation around pi- bond is restricted.	Rotation is possible around sigma bond.
5	Electron cloud is not symmetrical about the line joining 2 nuclei.	Electron cloud is symmetrical about the line joining 2 nuclei.
6	It is having 2 electron clouds one above the plane of atomic nuclei and one below the plane of atomic nuclei.	It is having 1 electron cloud and that is symmetrical about the inter-nuclear axis.

Q-33) Using valence bond theory, explain the formation of hydrogen molecule.

Ans.)

Assuming 2 H- atoms X and Y with nuclei N_X and N_Y and electrons e_X and e_Y respectively.

When X and Y are far from each other then there is no interaction between them. As soon as they come closer, the attractive force and repulsive force becomes active.

The repulsive forces are:


- Between electrons of both the atoms i.e. e_X and e_Y .
- Between nuclei of both the atoms i.e. N_X and N_Y .

The attractive forces are:


- Between the electron and nucleus of the same atom i.e. $N_X - e_X$ and $N_Y - e_Y$.
- Between the electron of one atom and nucleus of other atom i.e. $N_X - e_Y$ and $N_Y - e_X$.

The repulsive force pushes the 2 atoms apart whereas the attractive force tends to bring them together.

Repulsive forces:



Attractive forces:





The values of repulsive forces are less than that of attractive forces. Thus, 2 atoms approach each other. Thus, there is decrease in potential energy. At the end a stage is reached when the repulsive forces balance the attractive forces and the system achieves the minimum energy, which leads to formation of H_2 molecule.

Q-34) What are the conditions that are required for linear combination of atomic orbitals to form molecular orbitals.

Ans.)

The conditions that are required for linear combination of atomic orbitals to form molecular orbitals are as follows:

- (i) The joining of atomic orbitals must have approximately same energy. This implies in a homo-nuclear molecule, the 1s-orbital of one atom can join with the 1s-orbital of another atom, but cannot join with the 2s-orbital.
- (ii) The joining atomic orbitals must have legitimate orientations to ensure the maximum overlap.
- (iii) The overlapping must be in a large extent.


Q-35) Explain why molecule Be_2 doesn't exist by using molecular orbital theory.

Ans.)

Electronic configuration of Be:



Molecular orbital electronic configuration of Be_2 is:



Thus, bond order of Be_2 : $0.5(N_b - N_a)$.

N_a : No. of electrons in the bonding orbitals

N_b : No. of electrons in the anti-bonding orbitals

Therefore, bond order of $Be_2 = 0.5(4 - 4) = 0$

Zero value of bond order indicates that given molecule is unstable. Thus, Be_2 doesn't exist.

Q-36) Compare relative stability of the species given below and also write their magnetic properties.

(a) O_2



(b) O_2^- (peroxide)

(c) O_2^{2-} (superoxide)

(d) O_2^+

Ans.)

O_2 contain 16 electrons i.e. 8 electrons from each O- atom.

Electronic configuration of O_2 is:

$$[\sigma(1s)]^2[\sigma^*(1s)]^2[\sigma(2s)]^2[\sigma^*(2s)]^2[\sigma(2p_z)]^2[\pi(2p_x)]^2[\pi(2p_y)]^2[\pi^*(2p_x)]^1[\pi^*(2p_y)]^1$$

As 1s- orbital of each O- atom does not involve in the bonding,

No. of bonding electrons = $N_b = 8$

No. of anti-bonding electrons = $N_a = 4$

Now,

$$\text{Bond order} = 0.5(8 - 4) = 2$$

Electronic configuration of O_2^{2-} (peroxide) is:

$$KK[\sigma(2s)]^2[\sigma^*(2s)]^2[\sigma(2p_z)]^2[\pi(2p_x)]^2[\pi(2p_y)]^2[\pi^*(2p_x)]^2[\pi^*(2p_y)]^2$$

No. of bonding electrons = $N_b = 8$

No. of anti-bonding electrons = $N_a = 6$

Now,

$$\text{Bond order} = 0.5(8 - 6) = 1$$

Electronic configuration of O_2^- (superoxide) is:

$$KK[\sigma(2s)]^2[\sigma^*(2s)]^2[\sigma(2p_z)]^2[\pi(2p_x)]^2[\pi(2p_y)]^2[\pi^*(2p_x)]^2[\pi^*(2p_y)]^1$$

No. of bonding electrons = $N_b = 8$

No. of anti-bonding electrons = $N_a = 5$

Now,

$$\text{Bond order} = 0.5(8 - 5) = 1.5$$

Electronic configuration of O_2^+ is:

$$KK[\sigma(2s)]^2[\sigma^*(2s)]^2[\sigma(2p_z)]^2[\pi(2p_x)]^2[\pi(2p_y)]^2[\pi^*(2p_x)]^1$$

No. of bonding electrons = $N_b = 8$

No. of anti-bonding electrons = $N_a = 3$

Now,

$$\text{Bond order} = 0.5(8 - 3) = 2.5$$

As, bond dissociation energy $\propto$ bond order

Hence, higher the bond order, higher stability will be there.

Arrangement according to decreasing order of stability is given as:

$$O_2^+ > O_2 > O_2^- > O_2^{2-}$$

Q-37) What is the significance of positive (+) and negative (-) sign in representing the molecular orbitals?

Ans.)

Generally molecular orbital are represented by the 'wave function'.

Positive (+) sign in representing a molecular orbital indicates positive wave function.

Negative (-) sign in representing a molecular orbital indicates negative wave function.

Q-38) Which hybridization is PCl_5 is having? Why are equatorial bonds are shorter than axial bonds?


Ans.)

The electronic configuration of outer orbital of phosphorus in excited state and in ground state is given below:


Ground State:




Excited State:



Phosphorus atom is having sp^3d hybridization. These orbitals are filled due to donation of electron pairs by 5 Cl^- atoms: PCl_5



The 5 sp^3d hybrid orbitals present here are directed towards 5 corners of trigonalbipyramidal. Thus, the geometry of PCl_5 is given below:



PCl_5 contains 5 P- Cl sigma bonds. Out of which 3 P-Cl bond lie in only 1 plane and they are making 120° with each other. And as these bonds lie in 1 plane they are known equatorial bonds.

Out of 2 remaining P-Cl bonds one bond lie above the equatorial plane and one bond lie below the equatorial bond. And they are making 90° with each other. These bonds are called axial bond.


Equatorial bond pairs repel axial bond pairs to a large extent so, equatorial bonds are slightly shorter than axial bonds.

Q-39) Give definition of H- bond (Hydrogen bond). Is H- bond stronger or weaker compared to Van der Waals force?

Ans.)

"H-bond is defined as an attractive force acting between the hydrogen attached to an electronegative atom of one molecule and an electronegative atom of a different molecule".


As there is difference in the electronegativities between the atoms, thus, the bond pair electronegative atom and hydrogen atom is drifted away from H- atom. Therefore, hydrogen atom gets electropositive w.r.t. the other atom and procures a positive charge.



The value of H- bond is minimum in gaseous state and maximum in the solid state.

Two types of hydrogen bond are there:

- (a) Intramolecular hydrogen bond e.g., o- nitrophenol
- (b) Inter molecular hydrogen bond e.g., HF, H₂O etc.



H- bonds are stronger than Van der Waals forces as H- bond are regarded as extreme form of the dipole- dipole interaction.

Q-40) What do you mean by bond order? Also calculate the bond order for:

- (a) O₂
- (b) O₂⁺ (superoxide)
- (c) O₂⁻
- (d) N₂

Ans.)

Bond Order is defined as 0.5 times the difference between the # of electrons

Bond Order: It is defined as 0.5 times the difference between the "No. of electrons present in bonding orbitals and No. of electrons present in anti-bonding orbitals" of a molecule.

$$\text{Bond Order} = 0.5(N_b - N_a);$$

N_a : No. of anti-bonding electrons

N_b : No. of bonding electrons

O_2 contain 16 electrons i.e. 8 electrons from each O- atom.

Electronic configuration of O_2 is:

$$[\sigma(1s)]^2[\sigma^*(1s)]^2[\sigma(2s)]^2[\sigma^*(2s)]^2[\sigma(2p_z)]^2[\pi(2p_x)]^2[\pi(2p_y)]^2[\pi^*(2p_x)]^1[\pi^*(2p_y)]^1$$

As 1s- orbital of each O- atom does not involve in the bonding,

$$\text{No. of bonding electrons} = N_b = 8$$

$$\text{No. of anti-bonding electrons} = N_a = 4$$

Now,

$$\text{Bond order} = 0.5(8 - 4) = 2$$

Electronic configuration of O_2^- (superoxide) is:

$$KK[\sigma(2s)]^2[\sigma^*(2s)]^2[\sigma(2p_z)]^2[\pi(2p_x)]^2[\pi(2p_y)]^2[\pi^*(2p_x)]^2[\pi^*(2p_y)]^1$$

$$\text{No. of bonding electrons} = N_b = 8$$

$$\text{No. of anti-bonding electrons} = N_a = 5$$

Now,

$$\text{Bond order} = 0.5(8 - 5) = 1.5$$

Electronic configuration of O_2^+ is:

$$KK[\sigma(2s)]^2[\sigma^*(2s)]^2[\sigma(2p_z)]^2[\pi(2p_x)]^2[\pi(2p_y)]^2[\pi^*(2p_x)]^1$$


$$\text{No. of bonding electrons} = N_b = 8$$

$$\text{No. of anti-bonding electrons} = N_a = 3$$

Now,

$$\text{Bond order} = 0.5(8 - 3) = 2.5$$

Electronic configuration of N_2 is:




No. of bonding electrons = $N_b = 10$

No. of anti-bonding electrons = $N_a = 4$

Now,

$$\text{Bond order} = 0.5(10 - 4) = 3$$

Class 11-12 - Introduction to Chemical Bonding



Noble Gases

2 He, 10 Ne, 18 Ar, 36 Kr, 54 Xe, 86 Rn, 118 Uuo

stable atom