

1. SOME BASIC CONCEPTS OF CHEMISTRY

Chemistry is the branch of Science that deals with the properties, structure and composition of matter. There are a large number of branches for Chemistry. Some of them are:

1. Inorganic Chemistry
2. Organic Chemistry
3. Physical Chemistry
4. Analytical Chemistry
5. Polymer Chemistry
6. Biochemistry
7. Medicinal Chemistry
8. Industrial Chemistry
9. Hydrochemistry
10. Electrochemistry
11. Green Chemistry etc.

Matter: Matter is anything that occupies space, has a definite mass and can be perceived by any of our sense organs. Based on the physical state we can divide matter into different categories.

1. Solid state
2. Liquid state
3. Gaseous state
4. Plasma state
5. Bose-Einstein condensate
6. Fermionic condensate

Classification of matter

Based on the chemical composition of matter can be divided into two categories – pure substances and mixtures.

Pure substances contain only one type of particles. E.g. sodium (Na), Potassium (K), Hydrogen (H), Oxygen (O), Helium (He), Carbon dioxide (CO₂), water (H₂O), ammonia (NH₃), cane sugar (C₁₂H₂₂O₁₁) etc. These are further divided into two – **elements and compounds**.

Elements are pure substances which contain only one type of particles. These particles may be atoms or molecules. The term element was first introduced by Robert Boyle, the father of modern Chemistry. Now there are about 117 elements. Some elements exist as monoatomic and some others are polyatomic. E.g. Hydrogen, Nitrogen, Oxygen (diatomic), Sodium, Potassium, Lithium, Calcium (monoatomic), Phosphorus, Sulphur (polyatomic) etc.

Compounds are pure substances which contain more than one type of atoms. E.g. CO₂, H₂O, NH₃, H₂SO₄ etc.

Mixtures contain more than one type of particles. E.g. all types of solutions, gold ornaments, sea water, muddy water, air etc.

There are two types of mixtures – homogeneous and heterogeneous mixtures. Mixtures having uniform composition throughout are called **homogeneous mixtures**. E.g. all type of solutions, air etc.

Mixtures having different compositions at different parts are called **heterogeneous mixtures**. E.g. sea water, soil etc.

Mass and Weight

Mass is the amount of matter present in a body. It is a constant quantity. Its SI unit is kilogram (kg). Weight is the gravitational force acting on a body. It is a variable quantity. i.e. it changes with place. Its SI unit is newton (N).

Volume (V)

It is the space occupied by a body. Its SI unit is m³. Other units are cm³, mL, L etc.

$$1 \text{ m}^3 = 10^6 \text{ cm}^3$$

$$1 \text{ cm}^3 = 1 \text{ mL}$$

$$1 \text{ L} = 10^3 \text{ cm}^3 (\text{mL})$$

$$1 \text{ dm}^3 = 10^3 \text{ cm}^3$$

Density (d)

It is the amount of mass per unit volume.

i.e. density = mass/volume. Its SI unit is kg/m³. But it is commonly expressed in g/cm³.

Temperature (T)

It is the degree of hotness or coldness of a body. It is commonly expressed in degree Celsius (°C). Other units are degree Fahrenheit (°F), Kelvin (K) etc. its SI unit is Kelvin (K).

Degree Celsius and degree Fahrenheit are related as:

$$^{\circ}\text{F} = 9/5(^{\circ}\text{C}) + 32$$

Degree celsius and Kelvin are related as:

$$\text{K} = ^{\circ}\text{C} + 273.15$$

Precision and Accuracy

Precision refers to the closeness of various measurements for the same quantity. But, accuracy is the agreement of a particular value to the true value of the result.

Significant Figures

Every experimental measurement has some amount of uncertainty associated with it. The uncertainty in the experimental or the calculated values is indicated by mentioning the number of significant figures. **Significant figures are meaningful digits which are known with certainty.** The uncertainty is indicated by writing the certain digits and the last uncertain digit.

There are certain rules for determining the number of significant figures. These are:

1. All non-zero digits are significant. For example in 295 cm, there are three significant figures and in 0.25 mL, there are two significant figures.
2. Zeros preceding to first non-zero digit are not significant. Such zero indicates the position of decimal point. Thus, 0.03 has one significant figure and 0.0052 has two significant figures.
3. Zeros between two non-zero digits are significant. Thus, 2.005 has four significant figures.
4. Zeros at the end or right of a number are significant if they are on the right side of the decimal point; otherwise, they are not significant. For example, 0.200 g has three significant figures.
5. Exact numbers have an infinite number of significant figures. For example, in 2 balls or 20 eggs, there are infinite significant figures since these are exact numbers and can be represented by writing infinite number of zeros after placing a decimal i.e., $2 = 2.000000$ or $20 = 20.000000$
6. When numbers are written in scientific notation, the number of digits between 1 and 10 gives the number of significant figures. For e.g. 4.01×10^2 has three significant figures, and 8.256×10^{-3} has four significant figures.

LAWS OF CHEMICAL COMBINATIONS

The combination of elements to form compounds is governed by the following five basic laws:

1. **Law of Conservation of Mass (Law of indestructibility of matter):** This law was proposed by **Antoine Lavoisier**. It states that *matter can neither be created nor destroyed. We can only convert one form of matter into another form. Or, in a chemical reaction, the total mass of reactants is equal to the total mass of products.* Chemical equations are balanced according to this law.

Illustration



Here 4 g of H₂ combines with 32 g of O₂ to form 36 g of water.

Total mass of reactants = 4 + 32 = 36g

Total mass of products = 36 g

2. **Law of Definite Proportions (Law of definite composition):** This law was proposed by **Joseph Proust**. It states that *a given compound always contains exactly the same proportion of elements by weight. Or, the same compound always contains the same elements combined in a fixed ratio by mass.*

Illustration: Carbon dioxide can be formed in the atmosphere by various methods like respiration, burning of fuels, reaction of metal carbonates and bicarbonates with acid etc. All these samples of CO_2 contain only two elements Carbon and Oxygen combined in a mass ratio 3:8.

3. **Law of Multiple Proportions:** This law was proposed by **John Dalton**. It states that *if two elements can combine to form more than one compound, the different masses of one of the elements that combine with a fixed mass of the other element, are in small whole number ratio.*

Illustration: Hydrogen combines with oxygen to form two compounds – water and hydrogen peroxide.

Hydrogen + Oxygen \rightarrow Water

2g 16g 18g

Hydrogen + Oxygen \rightarrow Hydrogen Peroxide

2g 32g 34g

Here, the masses of oxygen (i.e. 16 g and 32 g) which combine with a fixed mass of hydrogen (2g) bear a simple ratio, i.e. 16:32 or 1: 2.

4. **Gay Lussac's Law of Gaseous Volumes:** This law was proposed by **Gay Lussac**. It states that *when gases combine to form gaseous products, their volumes are in simple whole number ratio at constant temperature and pressure.*

Illustration: H_2 combines with O_2 to form water vapour according to the equation $2\text{H}_2(\text{g}) + \text{O}_2(\text{g}) \rightarrow 2\text{H}_2\text{O}(\text{g})$. If 100 mL of hydrogen combine with 50 mL of oxygen, we get 100 mL of water vapour. Thus, the volumes of hydrogen and oxygen which combine together (i.e. 100 mL and 50 mL) bear a simple ratio of 2:1.

5. **Avogadro's Law:** This law was proposed by Amedeo Avogadro. It states that equal volumes of all gases at the same temperature and pressure should contain equal number of moles or molecules.

Illustration: If we take 10L each of NH_3 , N_2 , O_2 and CO_2 at the same temperature and pressure, all of them contain the same number of moles and molecules.

DALTON'S ATOMIC THEORY

The term atom was first used by John Dalton from the Greek word a-tomio (means indivisible). He proposed the first atomic theory. The important postulates of this theory are:

1. Matter is made up of minute and indivisible particles called atoms.
2. Atoms can neither be created nor be destroyed.
3. Atoms of same element are identical in their properties and mass. While atoms of different elements have different properties and mass.
4. Atoms combined to form compound atoms called molecules.
5. When atoms combine, they do so in a fixed ratio by mass.

Dalton's theory could explain the laws of chemical combination.

Atoms and Molecules

Atom is the smallest particle of an element. Molecules are the smallest particle of a substance. A molecule has all the properties of that substance.

Types of molecules

Based on the type of atoms, there are two types of molecules – homonuclear molecule and heteronuclear molecule. A molecule containing only one type of atom is called *homonuclear molecule*.

e.g. H_2 , O_2 , N_2 , O_3 (ozone) etc

Heteronuclear molecules contain different types of atoms. E.g. CO_2 , H_2O , $\text{C}_6\text{H}_{12}\text{O}_6$, NH_3 etc.

Based on the no. of atoms there are three types of molecules – monoatomic, diatomic and polyatomic molecules.

Monoatomic molecules contain only one atom. E.g. all metals, noble gases like He, Ne, Ar etc.

Diatomic molecules contain 2 atoms. E.g. H_2 , O_2 , N_2 , halogens (F_2 , Cl_2 , Br_2 and I_2)

Polyatomic molecules contain more than two atoms. E.g. ozone (O_3), Phosphorus (P_4), Sulphur (S_8) etc.

Atomic mass

Atomic mass of an element is a number that expresses how many times the mass of an atom of the element is greater than $1/12^{\text{th}}$ the mass of a C^{12} atom.

For e.g. atomic mass of Nitrogen is 14, which means that mass of one N atom is 14 times greater than $1/12^{\text{th}}$ the mass of a C^{12} atom.

Atomic mass unit (amu): $1/12^{\text{th}}$ the mass of a C^{12} atom is called atomic mass unit (amu).

$$\begin{aligned}\text{i.e. } 1 \text{ amu} &= 1/12 \times \text{mass of a } \text{C}^{12} \text{ atom} \\ &= 1.66 \times 10^{-24} \text{ g} \\ &= 1.66 \times 10^{-27} \text{ kg}\end{aligned}$$

Today, 'amu' has been replaced by 'u' which is known as unified mass.

Average atomic mass: All most all the elements have isotopes. So we can calculate an average atomic mass of an element by considering the atomic mass of the isotopes and their relative abundance. For e.g. chlorine has two isotopes Cl^{35} and Cl^{37} in the ratio 3:1. So the average atomic mass $\text{Cl} = (3 \times 35 + 1 \times 37)/4 = 35.5$

Molecular mass:

Molecular mass is the sum of atomic masses of the elements present in a molecule. It is obtained by multiplying the atomic mass of each element by the number of its atoms and adding them together.

For e.g. molecular mass of H_2SO_4 is calculated as: $2 \times 1 + 32 + 4 \times 16 = 98 \text{ u}$.

Formula mass:

In the case of ionic compounds (like NaCl), there is no discrete (separate) molecules. Here the positive ions and the negative ions are arranged in a three-dimensional structure. So we can calculate only formula mass by taking molecular formula of the compound.

Mole concept

Mole is the unit of amount of substance. It is defined as *the amount of substance that contains as many particles as there are atoms in exactly 12 g C^{12} isotope*. **1 mole of any substance contains 6.022×10^{23} atoms**. This number is known as **Avogadro number or Avogadro constant** (N_A or N_0).

1 mol of hydrogen atoms = 6.022×10^{23} atoms

1 mol of water molecules = 6.022×10^{23} water molecules

1 mol of sodium chloride = 6.022×10^{23} formula units of sodium chloride

Molar mass: The mass of one mole of a substance in gram is called its molar mass (gram molecular mass). The molar mass in grams is numerically equal to molecular mass in u.

Molar mass of oxygen = 32g

Molar mass of hydrogen = 2g etc.

Molar volume: It is the volume of 1 mole of any substance. At standard temperature and pressure (STP), molar volume of any gas = 22.4 L (or, 22400 mL). i.e. 22.4 L of any gas at STP contains 1 mole of the gas or 6.022×10^{23} molecules of the gas and its mass = molar mass.

For e.g. 22.4 L of hydrogen gas = 1 mole of H_2 = 6.022×10^{23} molecules of hydrogen = 2 g of H_2

Percentage composition

It is the percentage of each elements present in 100g of a substance.

i.e. percentage composition (mass percent) of an element = $\frac{\text{Mass of that element in the compound} \times 100}{\text{Molar mass of the compound}}$

It is helpful in checking the purity of a given sample. Also by knowing the percentage composition, we can calculate the empirical and molecular formula of a compound.

Empirical and Molecular formulae

Empirical formula is the simplest formula which gives only the ratio of different elements present in the compound. But molecular formula is the actual formula that gives the exact number of different elements present in the sample. For e.g. the empirical formula of glucose is CH_2O but its molecular formula is $\text{C}_6\text{H}_{12}\text{O}_6$. By knowing the percentage composition, we can calculate the empirical and molecular formula of a compound.

Stoichiometry and Stoichiometric calculations

The word 'stoichiometry' is derived from two greek words – stoicheion (meaning element) and metron (meaning measure). Thus stoichiometry deals with the calculations involving the masses or the volumes of reactants and the products.

Chemical Equation

It is the representation of a chemical reaction by symbols and formulae. Here the reactants are written in the left hand side and the products, on the right hand side. (The substances which participate in a chemical reaction are called **reactants** and the substances which are formed as a result of a reaction are called **products**).

A chemical equation should be balanced and the physical states of reactants and products are written in brackets.

The following informations are obtained from a chemical equation.

1. An idea about the reactants and products and their physical states.
2. An idea about the masses of reactants and products.
3. An idea about the number moles and molecules of reactants and products.
4. An idea about the volumes of reactants and products at STP.

Limiting reagent (Limiting reactant):

The reagent which limits a reaction or the reagent which is completely consumed in a chemical reaction is called limiting reagent or limiting reactant.

For e.g. in the reaction $2\text{SO}_2(\text{g}) + \text{O}_2(\text{g}) \longrightarrow 2\text{SO}_3(\text{g})$, 2 moles of SO_2 reacts completely with 1 mole of O_2 to form 2 moles of SO_3 . If we take 10 moles each of SO_2 and O_2 , we get only 10 moles of SO_3 because 10 moles of SO_2 requires only 5 moles of O_2 for the complete reaction. So here SO_2 is the limiting reagent and 5 moles of O_2 remains unreacted.

Reactions in solutions

Solutions are homogeneous mixture containing 2 or more components. The component which is present in larger quantity is called solvent and the other components are called solutes. Or, the substance which is dissolved is called solute and the substance in which solute is dissolved is called solvent.

For e.g. in NaCl solution, NaCl is the solute and water is the solvent.

A solution containing only 2 components are called **binary solution**. If the solvent is water, it is called *aqueous solution*.

One of the most important terms related to a solution is its concentration. *It is defined as the amount of solute present in a given volume of solution.* Concentration can be expressed in the following ways:

1. **Mass percent (w/w or m/m):** It is defined as the number of parts solute present in 100 parts by mass of solution.

$$\text{i.e. Mass \% of a component} = \frac{\text{Mass of solute} \times 100}{\text{Mass of solution}}$$

2. **Mole fraction:** It is defined as the ratio of the number of moles of a particular component to the total number of moles of solution.

$$\text{i.e. Mole fraction of a component} = \frac{\text{Number of moles of the component}}{\text{Total number of moles of all the components}}$$

For example, in a binary solution, if the number of moles of A and B are n_A and n_B respectively, then the mole fraction of A (x_A) = $n_A / (n_A + n_B)$ and that of the component B (x_B) = $n_B / (n_A + n_B)$

$$x_A + x_B = 1$$

i.e the sum of the mole fractions of all the components in a solution is always equal to 1.

3. **Molarity (M):** It is defined as the number of moles of solute dissolved per litre of solution.

$$\text{i.e. Molarity (M)} = \frac{\text{Number of moles of solute (n)}}{\text{Volume of solution in litre (V)}}$$

1 M NaOH solution means 1 mole of NaOH is present in 1 L of solution.

4. **Molality (m):** It is defined as the number of moles of the solute present per kilogram (kg) of the solvent.

$$\text{i.e. Molality (m)} = \frac{\text{Number of moles of solute}}{\text{Mass of solvent in kg}}$$

Among the above concentration terms, molarity depends on temperature because it is related to volume, which changes with temperature. All the others are temperature independent.
