

# MINERAL NUTRITION

## Points To Remember :

- **Autotrophs** : An organism that synthesizes its required nutrients from simple and inorganic substances.
- **Heterotrophs** : An organism that cannot synthesize its own nutrients and depends on others.

## Essential Mineral elements :

- More than sixty elements found in different plants.
- Some plants accumulate selenium, some other gold.

## Criteria for Essentiality :

- Element absolutely necessary for normal growth and reproduction.
- In the absence of the element the plant cannot complete its life cycle.
- Role of the element can not be replaced by any other elements.
- The element must be directly involved in the metabolism of plant.

**Macronutrients** : are generally present in the plant's tissues in large amount (in excess of 10 mmole  $\text{Kg}^{-1}$  of dry matter).

**Micronutrients** : or trace elements are needed in very small amounts (less than 10 mmole  $\text{Kg}^{-1}$  of dry matter)

## Four groups of essential elements :

- As components of **biomolecules** and forms structural elements of cells (e.g. carbon, hydrogen, oxygen and nitrogen)
- As components of **energy-related** chemical compounds in plants. (magnesium in chlorophyll and phosphorus in ATP)
- Element that **activate** or **inhibit enzymes** ( $\text{Mg}^{2+}$ ,  $\text{Zn}^{2+}$ )
- Alter the **osmotic potential** of a cell. ( $\text{K}^+$ )

## Role of macro and micro-nutrients :

### Nitrogen :

- Absorbed in the form of  $\text{NO}_2^-$  or  $\text{NH}_4^+$
- Required by meristematic tissue and metabolically active tissue.
- Constituent of proteins, nucleic acids, vitamins and hormones.

### Phosphorus :

- Absorbed in the form of  $\text{H}_2\text{PO}_4^-$  or  $\text{HPO}_4^{2-}$ .
- Constituents of cell membrane certain proteins, all nucleic acids and required in **phosphorylation reaction**.

### Potassium :

- Absorbed as potassium ion ( $\text{K}^+$ )
- Required in meristematic tissues.
- Maintain cation and anion balance in cell.
- **Opening and closing** of stomata.
- Activation of enzyme.
- Maintenance of turgidity of cells.

### Calcium :

- Absorbed in the form of calcium ions ( $\text{Ca}^{2+}$ ).
- Required by meristematic and differentiating tissues.
- Used in synthesis of cell wall particularly as calcium pectate in **middle lamella**.
- Required during formation of **mitotic spindle**.
- Involved in normal functioning of cell membrane.
- Activate certain enzyme.
- Important role in regulating metabolic activity.

### Magnesium :

- Absorbed in the form of  $\text{Mg}^{2+}$ .
- Activates enzymes of respiration photosynthesis.
- Involved in the synthesis of DNA and RNA.
- Constituent of the **ring structure of chlorophyll**.
- Maintain **ribosome structure**.

### Sulphur :

- Absorbed in the form of sulphate  $\text{SO}_4^{2-}$ .
- Present in two amino acids **cystine** and **methionine**
- Main constituent of several coenzyme, vitamins and ferredoxin.

### Iron :

- Obtained in the form of ferric ions ( $\text{Fe}^{3+}$ ).
- Required in larger amount in comparison to other elements.
- Constituent of proteins involved in the transfer of electron like **ferredoxin** and **cytochromes**.
- Activates catalase enzyme.
- Essential for formation of chlorophyll.

### Manganese :

- Absorbed in the form of manganous ions ( $Mn^{2+}$ ).
- Activates many enzymes of photosynthesis, respiration and nitrogen metabolism.
- **Photolysis of water** and evolution of oxygen during light reaction.

### Zinc :

- Obtained in the form of  $Zn^{2+}$ .
- Activates enzymes like **carboxylase**.
- Required in **synthesis of auxin**.

### Copper :

- Absorbed in the form of cupric ions ( $Cu^{2+}$ ).
- Essential for overall metabolism.
- Associated with enzyme involved in redox reactions.

### Boron :

- Absorbed in the form of  $BO_3^{3-}$  or  $B_4O_7^{2-}$ .
- Required in **uptake and utilization of  $Ca^{2+}$** .
- Pollen germination.
- Cell elongation.
- Cell differentiation.
- Carbohydrate translocation.

### Molybdenum :

- Obtained in the form of molybdate ions ( $MoO_4^{2-}$ ).
- Component of enzyme like **nitrogenase** and **nitrate reductase**.
- Required in nitrogen metabolism.

### Chlorine :

- Absorbed in the form of chloride anion ( $Cl^-$ ).
- Along with  $Na^+$  and  $K^+$  it determines the solute concentration.
- Maintain anion cation balance of the cell.
- Essential for **photolysis of water** during light reaction of photosynthesis.

### Deficiency symptoms of essential elements :

- Critical concentration: the concentration of the essential element below which plant growth is retarded.
- The element is said to be deficient when present below the critical concentration.
- For the elements that are actively mobilized within the plant that show the deficiency symptoms in the older tissues. (nitrogen, potassium and magnesium)

- The deficiency symptoms tend to appear first in the young tissues whenever the elements are relatively immobile and are not transported out of the mature organs.(sulphur and calcium)
- Deficiency symptom includes chlorosis, necrosis, and stunted growth, premature fall of leaves and buds, and inhibition of cell division.
- **Chlorosis:** is the loss of chlorophyll.
- **Necrosis:** death of cells and tissues.

#### Toxicity of Micronutrients :

- Micronutrient required in low amount.
- Moderate decrease causes the deficiency symptoms.
- Moderate increase causes toxicity.
- Any mineral ion concentration in tissues that reduces the dry weight of the tissues by 10 percent is considered **toxic**.

#### Nitrogen cycle :

- **Nitrogen fixation:** conversion of molecular nitrogen into ammonia.
- **Biological nitrogen fixation:** Conversion of atmospheric into organic compounds by living organisms.
- **Ammonification:** decomposition of organic nitrogen of dead plants and animals into ammonia is called Ammonification. (*Nitrosomonas* bacteria)
- **Nitrification.** Ammonia oxidized into nitrite by *Nitrosomonas* and *Nitrococcus* bacteria. The nitrite further oxidized to nitrate with the help of *Nitrobacter*. These steps are called nitrification.
- **Assimilation:**
  - Nitrates absorbed by plant from soil and transported to the leaves.
  - In the leaves nitrates reduced to form ammonia that finally forms the amine group of amino acids.
- **Denitrification:** Nitrate in the soil is also reduced to molecular nitrogen. This process is carried by bacteria like *Pseudomonas* and *Thiobacillus*.

#### Biological nitrogen fixation :

- Reduction of nitrogen to ammonia by living organisms is called biological nitrogen fixation.
- The enzyme nitrogenase which catalyses the process are present in prokaryotes, called nitrogen fixer.
- Nitrogen fixing microbes could be free-living or symbiotic.
- Free-living nitrogen fixing aerobic microbes are *Azotobacter* and *Beijernickia*.
- Free-living nitrogen fixing anaerobic microbes are *Rhodospirillum*.
- A number of cyanobacteria like *Anabaena* and *Nostoc* are free-living nitrogen fixer.

### Symbiotic nitrogen fixation :

- Best example of symbiotic nitrogen fixation is observed in legume-Rhizobium bacteria.
- Rhizobium form root nodules in leguminous plants.
- **Frankia** also produces nitrogen-fixing nodules on the roots of non-leguminous plants (e.g. Alnus).
- Both Rhizobium and Frankia are free living in soil, but as symbiont, can fix atmospheric nitrogen.
- The root nodules contain pink coloured pigment contains a protein called **leg-haemoglobin**.

### Nodule formation :

- Nodule formation involves a sequence of multiple interactions between Rhizobium and roots of the host plant.
- *Rhizobia* multiply and colonize the surroundings of roots and get attached to the epidermal and root hair cells.
- An infection thread is produced carrying the bacteria into the cortex of root.
- Bacteria released from the thread into the cells which differentiated into special nitrogen fixing cells.
- Nodule develops vascular connection for exchange of nutrients.
- The nodule contains an enzyme called **nitrogenase**.
- Nitrogenase is a Mo-Fe protein and catalyses the conversion of atmospheric nitrogen to ammonia.
- Nitrogenase is highly sensitive to molecular oxygen; it requires anaerobic condition.
- Nodule contains a special protein, called **leg-haemoglobin**.
- Leg-haemoglobin acts as **oxygen scavenger** and provides anaerobic condition to the bacteria inside the nodules; protect the enzyme nitrogenase from oxidation.
- Ammonia synthesis by nitrogenase is energetically expensive process; 8 ATP required synthesizing each molecule of  $\text{NH}_3$ .

### Fate of ammonia :

- At physiological pH, the ammonia is protonated to form  $\text{NH}_4^+$ .
- Most of plant assimilated nitrate and ammonium ions.
- **Reductive amination**: the ammonia reacts with  $\alpha$ -ketoglutaric acid and forms Glutamic acid.
- **Transamination**: it involves the transfer of amino group from one amino acid to the keto group of a keto acid.
- Glutamic acid is the main amino acid from which by the process of transamination other amino acids are synthesized.
- Two important amides - asparagines and glutamine found in the protein of plant.
- They are formed from two amino acids namely aspartic acid and Glutamic acid respectively.