

K.RAMALINGAM PG Asst.in Chemistry
SHRI RENGHA EDUCATIONAL INSTITUTIONS
RAYARPALAYAM
NAMAKKAL MAIN ROAD
9788855343

PUBLIC EXAMINATION 2019

CHEMISTRY

QUESTION TYPE B

1.B)First Order

2.A)Diethyl ether

3.A)(ii)and (iii)

4.D)(1)-(iii),(2)-(i),(3)-(iv),(4)-(ii)

5.D)Pentagonal bipyramidal

6.A)4.90BM

7.D)Strong reducing agents

8.C)dynamic

9.C) C_1-C_2

10.C) $\text{ohm}^{-1}\text{m}^{-1}$

11.C) $w-P\Delta V$

12.A)-OH

13.A)Ionic

14.B) $C_6H_5NH-NH-C_6H_5$

15.D) CH_3COCH_3

PUBLIC EXAMINATION 2019

CHEMISTRY

QUESTION TYPE A

1. A)Diethyl ether

2. B)First Order

3. B) $C_6H_5NH-NH-C_6H_5$

4. D)Pentagonal bipyramidal

5. D) CH_3COCH_3

6. C) C_1-C_2

7. C) $w-P\Delta V$

8. A)-OH

9. A)(ii)and (iii)

10. D)(1)-(iii),(2)-(i),(3)-(iv),(4)-(ii)

11. D)Strong reducing agents

12. A)Ionic

13. C)dynamic

14. A)4.90BM

15. C) $\text{ohm}^{-1}\text{m}^{-1}$

16. The higher the nuclear charge of protons in the nucleus, the higher is the ionization energy.

Because of the higher nuclear charge, the electrons are bound with more force and hence higher energy will be required for their removal.


For instance, magnesium has higher nuclear charge (12 protons) as compared to sodium (11 protons). Hence ionization energy in case of magnesium is higher as compared to sodium.

17. These elements show the oxidation states of +2, +3, +4, +5 and +6. Out of these, +4 oxidation state is most common state.

18. A small quantity of Radioactive CO_2^* containing radioactive oxygen O^{18} is mixed with ordinary carbon dioxide and the process is carried out.

It has been found that oxygen gas evolved along with sugar formation is non-radioactive.


Therefore O_2 produced comes from water and not from carbon dioxide. So the correct mechanism is as follows.



19. Glassy or vitreous state is a condition in which certain substance can exist, lying between the solid and liquid states.

20. Complete reduction of glucose with concentrated hydriodic acid in the presence of red phosphorus produces n-hexane as the major product.

This indicates that the six carbon atoms in the glucose molecule form an



This is used as Urinary antiseptic in medicine, in the name of 'Urotropine'.

24. II reaction will go faster than I reaction. This is due to the activation energy is minimum the reaction goes faster

25.


$$\begin{aligned} \text{Electronegativity of fluorine} &= \frac{(\text{IP})_F + (\text{EA})_F}{2 \times 2.8} \\ &= \frac{17.4 + 3.62}{5.6} = \frac{21.02}{5.6} = 3.75 \end{aligned}$$

K.RAMALINGAM PG Asst.in Chemistry
SHRI RENGA EDUCATIONAL INSTITUTIONS
RAYARPALAYAM
NAMAKKAL MAIN ROAD
9788855343

26.

Lanthanides		Actinides	
i)	Binding energies of 4f electrons are higher.	i)	Binding energies of 5f electrons are lower.
ii)	Maximum oxidation state exhibited by lanthanides is +4 e.g. Ce ⁴⁺	ii)	Due to lower binding energies they show higher oxidation states such as +4, +5 and +6. Uranium exhibits +6 oxidation state in UF ₆ and UO ₂ Cl ₂
iii)	4f electrons have greater shielding effect.	iii)	5f electrons have poor shielding effect.

27. Let 'a' and 'b' be the number of α β particles emitted during the change



Comparing the mass numbers,

$$232 = 208 + 4a + b \times 0$$

$$4a = 232 - 208$$

$$= 24$$

$$a = 6$$

Comparing the atomic numbers

$$90 = 82 + 2 \times a + (-1)b$$

$$= 82 + 2a - b$$

$$2a - b = 90 - 82 = 8$$

$$2(6) - b = 8$$

$$b = 12 - 8 = 4$$


Number of α - particle emitted = 6

Number of β - particles emitted = 4

28. 'Q' is defined as the ratio of product of initial concentrations

of products to the product of initial concentrations of reactants under nonequilibrium

conditions. For example, in the equilibrium.



Let [A], [B], [L] and [M] be the actual concentrations present before the occurrence of equilibrium. These concentrations are considered as the nonequilibrium concentration conditions and the reaction quotient

$$Q = \frac{[L]^l [M]^m}{[A]^a [B]^b}$$

'Q' is given as

29. (i) It should have a suitable colour.

(ii) It should be able to fix itself or be capable of being fixed to the fabric.

(iii) It should be fast to light.

30. The standard emf E_o , of a cell is the standard reduction potential of right-hand electrode (cathode) minus the standard reduction potential of the left-hand electrode (anode). That is,

$$E_{\text{cell}} = E_{\text{right}} - E_{\text{left}}$$

$$= \text{Cathode potential} - \text{Anode potential}$$

31. When equal amounts of d-isomer and l-isomer are mixed one gets a

"racemic mixture" and this process is called

racemisation. A racemic mixture

becomes optically inactive. Because, in this mixture rotation towards

clockwise direction by the dextro isomers is


compensated by the rotation

towards the anticlockwise direction by the laevo

isomers. Eg: d, l lactic acid

32.

K.RAMALINGAM PG Asst.in Chemistry
SHRI RENGA EDUCATIONAL INSTITUTIONS
RAYARPALAYAM
NAMAKKAL MAIN ROAD
9788855343



(1, 3, 4, 5, 6, pentahydroxy-2-hexanone)

Fructose molecules have 4 chiral centers of their carbon atoms. A maximum of 2^n optical isomers can occur for a molecule, where 'n' is the number of chiral centers. Since there are 4 chiral centers in fructose, it should have 2^4 = **16 optical isomers.**

33.

$$\text{Log}k_1/k_2 = E_a/2.303 R[1/T_2 - 1/T_1]$$

$$\therefore k_1/k_2 = 100000 \text{ J Mole}^{-1} / 2.303 * 8.314 \text{ JK}^{-1} \text{ mol}^{-1}$$


$$= -0.5745$$

$$\therefore k_1/k_2 \text{ antilog}(-0.5745)$$

$$= 0.2664$$

$$k_2 = 1/0.2664 k_1 = 3.75 \text{ times } k_1.$$

$\therefore k_2$, the rate constant at 35°C will be 3.75 times the rate constant at 25°C .



ii) **“it is impossible to measure simultaneously both the position and velocity (or momentum) of a microscopic particle with absolute accuracy or certainty.”**

Mathematically, uncertainty principle can be put as

$$\Delta x \cdot \Delta p \geq \frac{h}{4\pi}$$

follows.

where, Δx = uncertainty in the position of the particle and Δp = uncertainty in the momentum of the particle.

Or

b)i) One of the familiar characteristic of p-block elements is to show inert pair effect i.e. the tendency of being less availability for ns electron in bonding.

The inert pair effect increases down the group with the increase in atomic number.

ii) 1) Silicones act as excellent insulators for electric motors and other appliances as they can withstand high temperatures.

2) Straight chain polymers of 20 to 500 units are used as silicone fluids. They are water repellent because of the organic side group. These polymers are used in waterproofing textiles, as lubricants and as polish.

3) Silicone rubber retain their elasticity even at low temperatures and resist chemical attack. They are mixed with paints to make them damp-resistant.

35.a)i) $[\text{Ti}(\text{H}_2\text{O})_6]^{3+}$ is coloured because Ti^{3+} one electron present in d orbital The colour of transition metal ions is due to the presence of unpaired electrons in it and the energy gap between two energy levels in the same d-subshell being small. Sc^{3+} ions are also colourless because of the absence of d-electrons.

ii) The articles to be plated with chromium are made the cathode in an electrolytic bath consisting of chromic acid and sulphuric acid while the anode is made of a plate of lead. During electrolysis chromium deposits on the article (cathode). Generally the articles are first plated with nickel and then subjected to chromium plating.

Or


b)i)

1) Every metal atom has two types of valencies

i) Primary valency or ionisable valency

ii) Secondary valency or non ionisable valency

K.RAMALINGAM PG Asst.in Chemistry
SHRI RENGA EDUCATIONAL INSTITUTIONS
RAYARPALAYAM
NAMAKKAL MAIN ROAD
9788855343



b)i) A buffer solution is one which maintains its pH fairly constant even upon the addition of small amounts of acid or base. In other words, a buffer solution resists (or buffers) a change in its pH.

That is, we can add a small amount of an acid or base to a buffer solution and the pH will change very little.

ii) $K_a = 1.8 \times 10^{-5}$

$pK_a = -\log(1.8 \times 10^{-5}) = 4.7447$

$pH = pK_a + \log[\text{salt}]/[\text{acid}]$




... Henderson - Hasselbalch equation

$= 4.7447 + \log 0.5/0.5.$


$pH = 4.7447 + \log 1$

$= 4.7447$

38.a)i) The removal of water from the protonated species is facilitated by anhydrous zinc chloride. Reactivity in this reaction with respect to alcohols is $3^\circ > 2^\circ > 1^\circ$. This reaction is used in Lucas Test.




ii) -OH group, -OMe, group increase the reactivity of the benzene ring with respect to electrophilic attack and is **ortho, para - directing**. With a mixture of con.HNO₃ and con. H₂SO₄ it gives a mixture of ortho and para nitro anisole.




b)i) **Reducing property**

Formic acid is unique because it contains both an aldehyde group and a carboxyl group also. Hence it **can act as a reducing agent**. It reduces Fehling's solution, Tollens reagent and decolourises pink coloured KMnO₄ solution.


(a) Formic acid reduces ammoniacal silver nitrate solution (Tollen's reagent) to metallic silver.



(b) Formic acid reduces Fehling's solution. It reduces blue coloured cupric ions to red coloured cuprous ions.



(c) On the otherhand when sodium formate is heated to 360°C it decomposes to hydrogen and sodium oxalate.



ii) Aniline reacts with chloroform and alcoholic KOH to give an offensive smelling liquid, phenyl isocyanide.

