

Secondary School Certificate Examination

March 2019

Marking Scheme — Mathematics 30/2/1, 30/2/2, 30/2/3

General Instructions:

1. The Marking Scheme provides general guidelines to reduce subjectivity in the marking. The answers given in the Marking Scheme are suggested answers. The content is thus indicative. If a student has given any other answer which is different from the one given in the Marking Scheme, but conveys the meaning, such answers should be given full weightage.
2. Evaluation is to be done as per instructions provided in the marking scheme. It should not be done according to one's own interpretation or any other consideration — Marking Scheme should be strictly adhered to and religiously followed.
3. Alternative methods are accepted. Proportional marks are to be awarded.
4. If a candidate has attempted an extra question, marks obtained in the question attempted first should be retained and the other answer should be scored out.
5. A full scale of marks - 0 to 80 has to be used. Please do not hesitate to award full marks if the answer deserves it.
6. Separate Marking Scheme for all the three sets has been given.
7. As per orders of the Hon'ble Supreme Court. The candidates would now be permitted to obtain photocopy of the Answer book on request on payment of the prescribed fee. All examiners/Head Examiners are once again reminded that they must ensure that evaluation is carried out strictly as per value points for each answer as given in the Marking Scheme.

QUESTION PAPER CODE 30/2/1
EXPECTED ANSWER/VALUE POINTS

SECTION A

$$1. \text{ LCM } (336, 54) = \frac{336 \times 54}{6} \quad \frac{1}{2}$$

$$= 336 \times 9 = 3024 \quad \frac{1}{2}$$

$$2. \frac{3-a}{3a} - \frac{1}{a} = \frac{3-a-3}{3a} = -\frac{1}{3} \quad 1$$

$$3. 2x^2 - 4x + 3 = 0 \Rightarrow D = 16 - 24 = -8 \quad \frac{1}{2}$$

\therefore Equation has NO real roots $\frac{1}{2}$

$$4. \sin^2 60^\circ + 2 \tan 45^\circ - \cos^2 30^\circ = \left(\frac{\sqrt{3}}{2}\right)^2 + 2(1) - \left(\frac{\sqrt{3}}{2}\right)^2 \quad \text{[For any two correct values]} \quad \frac{1}{2}$$

$$= 2 \quad \frac{1}{2}$$

OR

$$\sin A = \frac{3}{4} \Rightarrow \cos A = \sqrt{1 - \frac{9}{16}} = \frac{\sqrt{7}}{4} \quad \frac{1}{2}$$

$$\sec A = \frac{4}{\sqrt{7}} \quad \frac{1}{2}$$

5. Point on x-axis is (2, 0) 1

6. $\triangle ABC$: Isosceles $\triangle \Rightarrow AC = BC = 4$ cm. $\frac{1}{2}$

$$AB = \sqrt{4^2 + 4^2} = 4\sqrt{2} \text{ cm} \quad \frac{1}{2}$$

OR

$$\frac{AD}{BD} = \frac{AE}{CE} \Rightarrow \frac{AD}{7.2} = \frac{1.8}{5.4} \quad \frac{1}{2}$$

$$\therefore AD = \frac{7.2 \times 1.8}{5.4} = 2.4 \text{ cm.} \quad \frac{1}{2}$$

SECTION B

7. Smallest number divisible by 306 and 657 = LCM (306, 657) 1
 LCM (306, 657) = 22338 1

8. A, B, C are collinear \Rightarrow ar. (ΔABC) = 0 $\frac{1}{2}$

$$\therefore \frac{1}{2}[x(6-3) - 4(3-y) - 2(y-6)] = 0 \quad 1$$

$$\Rightarrow 3x + 2y = 0 \quad \frac{1}{2}$$

OR

$$\text{Area of triangle} = \frac{1}{2}[1(6+5) - 4(-5+1) - 3(-1-6)] \quad 1$$

$$= \frac{1}{2}[11+16+21] = \frac{48}{2} = 24 \text{ sq. units.} \quad 1$$

9. P(blue marble) = $\frac{1}{5}$, P(black marble) = $\frac{1}{4}$

$$\therefore \text{P(green marble)} = 1 - \left(\frac{1}{5} + \frac{1}{4}\right) = \frac{11}{20} \quad 1$$

Let total number of marbles be x

$$\text{then } \frac{11}{20} \times x = 11 \Rightarrow x = 20 \quad 1$$

10. For unique solution $\frac{1}{3} \neq \frac{2}{k}$ 1

$$\Rightarrow k \neq 6 \quad 1$$

11. Let larger angle be x°

$$\therefore \text{Smaller angle} = 180^\circ - x^\circ \quad \frac{1}{2}$$

$$\therefore (x) - (180 - x) = 18 \quad \frac{1}{2}$$

$$2x = 180 + 18 = 198 \Rightarrow x = 99 \quad \frac{1}{2}$$

$$\therefore \text{The two angles are } 99^\circ, 81^\circ \quad \frac{1}{2}$$

OR

Let Son's present age be x years

Then Sumit's present age = $3x$ years.

$$\therefore \text{ 5 Years later, we have, } 3x + 5 = \frac{5}{2}(x + 5)$$

$$6x + 10 = 5x + 25 \Rightarrow x = 15$$

\therefore Sumit's present age = 45 years

12. Maximum frequency = 50, class (modal) = 35 – 40.

$$\text{Mode} = L + \left(\frac{f_1 - f_0}{2f_1 - f_0 - f_2} \right) \times h$$

$$= 35 + \frac{50 - 34}{100 - 34 - 42} \times 5$$

$$= 35 + \frac{16}{24} \times 5 = 38.33$$

SECTION C

13. Let $2 + 5\sqrt{3} = a$, where 'a' is a rational number.

$$\text{than } \sqrt{3} = \frac{a - 2}{5}$$

Which is a contradiction as LHS is irrational and RHS is rational

$$\therefore 2 + 5\sqrt{3} \text{ can not be rational}$$

Hence $2 + 5\sqrt{3}$ is irrational.

Alternate method:

Let $2 + 5\sqrt{3}$ be rational

$$\therefore 2 + 5\sqrt{3} = \frac{p}{q}, \text{ p, q are integers, } q \neq 0$$

$$\Rightarrow \sqrt{3} = \left(\frac{p}{q} - 2\right) \div 5 = \frac{p-2q}{5q}$$

LHS is irrational and RHS is rational
which is a contradiction.

$\therefore 2 + 5\sqrt{3}$ is irrational.

OR


$$2048 = 960 \times 2 + 128$$

$$960 = 128 \times 7 + 64$$

$$128 = 64 \times 2 + 0$$

$$\therefore \text{HCF}(2048, 960) = 64$$

14.



Correct Figure

$\triangle APB \sim \triangle DPC$ [AA similarity]

$$\frac{AP}{DP} = \frac{BP}{PC}$$

$$\Rightarrow AP \times PC = BP \times DP$$

OR

Correct Figure

In $\triangle POQ$ and $\triangle ROS$


$$\left. \begin{array}{l} \angle P = \angle R \\ \angle Q = \angle S \end{array} \right\} \text{alt. } \angle s$$

$\therefore \triangle POQ \sim \triangle ROS$ [AA similarity]


$$\therefore \frac{\text{ar}(\triangle POQ)}{\text{ar}(\triangle ROS)} = \left(\frac{PQ}{RS}\right)^2$$

$$= \left(\frac{3}{1}\right)^2 = \frac{9}{1}$$

$$\therefore \text{ar}(\triangle POQ) : \text{ar}(\triangle ROS) = 9 : 1$$



15.



Correct Figure

$$\Delta AOD \cong \Delta AOC \text{ [SAS]}$$

$$\Rightarrow \angle 1 = \angle 2$$

Similarly $\angle 4 = \angle 3$

$$\Rightarrow \angle 1 + \angle 4 = \angle 2 + \angle 3 = \frac{1}{2}(180^\circ)$$

$$\Rightarrow \angle 2 + \angle 3 = 90^\circ \text{ or } \angle AOB = 90^\circ$$

Alternate method:

Correct Figure

$$\Delta OAD \cong \Delta AOC \text{ [SAS]}$$

$$\Rightarrow \angle 1 = \angle 2$$


Similarly $\angle 4 = \angle 3$

$$\text{But } \angle 1 + \angle 2 + \angle 3 + \angle 4 = 180^\circ \quad [\because PQ \parallel RS]$$

$$\Rightarrow \angle 2 + \angle 3 = \angle 1 + \angle 4 = \frac{1}{2}(180^\circ) = 90^\circ$$

$$\therefore \text{ In } \Delta AOB, \angle AOB = 180^\circ - (\angle 2 + \angle 3) = 90^\circ$$

16.



Let the line $x - 3y = 0$ intersect the segment

joining $A(-2, -5)$ and $B(6, 3)$ in the ratio $k : 1$

$$\therefore \text{ Coordinates of P are } \left(\frac{6k - 2}{k + 1}, \frac{3k - 5}{k + 1} \right)$$

$$P \text{ lies on } x - 3y = 0 \Rightarrow \frac{6k - 2}{k + 1} = 3 \left(\frac{3k - 5}{k + 1} \right) \Rightarrow k = \frac{13}{3}$$

\therefore Ratio is $13 : 3$

$$\Rightarrow \text{ Coordinates of P are } \left(\frac{9}{2}, \frac{3}{2} \right)$$

$$\begin{aligned}
 17. \quad & \left(\frac{3 \sin 43^\circ}{\cos 47^\circ} \right)^2 - \frac{\cos 37^\circ \operatorname{cosec} 53^\circ}{\tan 5^\circ \tan 25^\circ \tan 45^\circ \tan 65^\circ \tan 85^\circ} \\
 & = \left(\frac{3 \sin 43^\circ}{\cos (90^\circ - 43^\circ)} \right)^2 - \frac{\cos 37^\circ \cdot \operatorname{cosec} (90^\circ - 37^\circ)}{\tan 5^\circ \tan 25^\circ (1) \tan (90^\circ - 25^\circ) \tan (90^\circ - 5^\circ)} \\
 & = \left(\frac{3 \sin 43^\circ}{\sin 43^\circ} \right)^2 - \frac{\cos 37^\circ \cdot \sec 37^\circ}{\tan 5^\circ \cdot \tan 25^\circ (1) \cot 25^\circ \cot 5^\circ} \\
 & = 9 - \frac{1}{1} = 8
 \end{aligned}$$

$$18. \text{ Radius of quadrant} = OB = \sqrt{15^2 + 15^2} = 15\sqrt{2} \text{ cm.}$$

Shaded area = Area of quadrant - Area of square 1/2

$$= \frac{1}{4} (3.14) [(15\sqrt{2})^2 - (15)^2]$$

$$= (15)^2 (1.57 - 1) = 128.25 \text{ cm}^2$$

OR


$$BD = \sqrt{(2\sqrt{2})^2 + (2\sqrt{2})^2} = \sqrt{16} = 4 \text{ cm}$$

\therefore Radius of circle = 2 cm 1/2

\therefore Shaded area = Area of circle - Area of square 1/2

$$\begin{aligned}
 & = 3.14 \times 2^2 - (2\sqrt{2})^2 \\
 & = 12.56 - 8 = 4.56 \text{ cm}^2
 \end{aligned}$$

19.



Height of cylinder = $20 - 7 = 13$ cm. 1

$$\text{Total volume} = \pi \left(\frac{7}{2} \right)^2 \cdot 13 + \frac{4}{3} \pi \left(\frac{7}{2} \right)^3 \text{ cm}^3$$

$$= \frac{22}{7} \times \frac{49}{4} \left(13 + \frac{4}{3} \cdot \frac{7}{2} \right) \text{ cm}^3$$

$$= \frac{77 \times 53}{6} = 680.17 \text{ cm}^3$$

$$20. \quad x_i: \quad 32.5 \quad 37.5 \quad 42.5 \quad 47.5 \quad 52.5 \quad 57.5 \quad 62.5$$

 $\frac{1}{2}$

$$f_i: \quad 14 \quad 16 \quad 28 \quad 23 \quad 18 \quad 8 \quad 3 \quad \Sigma f_i = 110$$

 $\frac{1}{2}$

$$u_i: \quad -3 \quad -2 \quad -1 \quad 0 \quad 1 \quad 2 \quad 3$$

$$f_i u_i: \quad -42 \quad -32 \quad -28 \quad 0 \quad 18 \quad 16 \quad 9, \quad \Sigma f_i u_i = -59$$

1

$$\text{Mean} = 47.5 - \frac{59 \times 5}{110} = 47.5 - 2.68 = 44.82$$

1

Note: If N is taken as 100, Ans. 44.55

Accept.

If some one write, data is wrong, give full 3 marks.

$$21. \quad \begin{array}{r} 3x^2 - 5 \overline{) 3x^4 - 9x^3 + x^2 + 15x + k} \left(x^2 - 3x + 2 \right. \\ \underline{3x^4} \\ -9x^3 \\ \underline{-9x^3} \\ 6x^2 + k \\ \underline{6x^2 - 10} \\ k + 10 \end{array}$$

2

$$\therefore k + 10 = 0 \Rightarrow k = -10$$

1

OR

$$p(y) = 7y^2 - \frac{11}{3}y - \frac{2}{3} = \frac{1}{3}(21y^2 - 11y - 2)$$

$$= \frac{1}{3}[(7y+1)(3y-2)]$$

1

$$\therefore \text{Zeroes are } 2/3, -1/7$$

 $\frac{1}{2}$

$$\text{Sum of zeroes} = \frac{2}{3} - \frac{1}{7} = \frac{11}{21}$$

$$\frac{-b}{a} = \frac{11}{21} \therefore \text{sum of zeroes} = \frac{-b}{a}$$

1

$$\text{Product of zeroes} = \left(\frac{2}{3}\right)\left(-\frac{1}{7}\right) = -\frac{2}{21}$$

$$\frac{c}{a} = -\frac{2}{3}\left(\frac{1}{7}\right) = -\frac{2}{21} \therefore \text{Product} = \frac{c}{a} \quad \frac{1}{2}$$

22. $x^2 + px + 16 = 0$ have equal roots if $D = p^2 - 4(16)(1) = 0$ 1

$$p^2 = 64 \Rightarrow p = \pm 8 \quad \frac{1}{2}$$

$$\therefore x^2 \pm 8x + 16 = 0 \Rightarrow (x \pm 4)^2 = 0 \quad 1$$

$$x \pm 4 = 0$$


$$\therefore \text{Roots are } x = -4 \text{ and } x = 4 \quad \frac{1}{2}$$

SECTION D

23. For correct, given, to prove, construction and figure $\frac{1}{2} \times 4 = 2$

For correct proof. 2

24. Correct Figure 1



In $\triangle APQ$

$$\frac{PQ}{AP} = \sin 30^\circ = \frac{1}{2} \quad \frac{1}{2}$$


$$PQ = (200)\left(\frac{1}{2}\right) = 100 \text{ m} \quad 1$$

$$PR = 100 - 50 = 50 \text{ m} \quad \frac{1}{2}$$

$$\text{In } \triangle PRD, \frac{PR}{PD} = \sin 45^\circ = \frac{1}{\sqrt{2}}$$

$$PD = (PR)(\sqrt{2}) = 50\sqrt{2} \text{ m} \quad 1$$

25. Total volume = $3.14 (12)^2 (220) + 3.14(8)^2(60) \text{ cm}^3$ 1



$$= 99475.2 + 12057.6 = 111532.8 \text{ cm}^3 \quad 1$$

$$\text{Mass} = \frac{111532.8 \times 8}{1000} \text{ kg} \quad 1$$

$$= 892.262 \text{ kg} \quad 1$$

26. Constructing an equilateral triangle of side 5 cm 1
- Constructing another similar Δ with scale factor $\frac{2}{3}$ 3
- OR
- Constructing two concentric circle of radii 2 cm and 5 cm 1
- Drawing two tangents PA and PB 2
- PA = 4.5 cm (approx) 1
27. Less than 40 less than 50 less than 60 less than 70 less than 80 less than 90 less than 100 $\frac{1}{2}$
- cf. 7 12 20 30 36 42 50 1
- Plotting of points (40, 7), (50, 12), (60, 20), (70, 30), (80, 36), (90, 42) and (100, 50) $1\frac{1}{2}$
- Joining the points to get the curve 1
28. LHS = $\frac{\tan \theta}{1 - \frac{1}{\tan \theta}} + \frac{\frac{1}{\tan \theta}}{1 - \tan \theta} = \frac{\tan^2 \theta}{\tan \theta - 1} - \frac{1}{\tan \theta (\tan \theta - 1)}$ 1
- = $\frac{\tan^3 \theta - 1}{\tan \theta (\tan \theta - 1)} = \frac{(\tan \theta - 1)(\tan^2 \theta + \tan \theta + 1)}{\tan \theta (\tan \theta - 1)}$ 1
- = $\tan \theta + 1 + \cot \theta = 1 + \frac{\sin \theta}{\cos \theta} + \frac{\cos \theta}{\sin \theta} = 1 + \frac{\sin^2 \theta + \cos^2 \theta}{\sin \theta \cos \theta}$ 1
- = $1 + \frac{1}{\sin \theta \cos \theta} = 1 + \operatorname{cosec} \theta \sec \theta = \text{RHS}$ 1
- OR
- Consider
- $\frac{\sin \theta}{\operatorname{cosec} \theta + \cot \theta} - \frac{\sin \theta}{\cot \theta - \operatorname{cosec} \theta} = \frac{\sin \theta}{\operatorname{cosec} \theta + \cot \theta} + \frac{\sin \theta}{\operatorname{cosec} \theta - \cot \theta}$ 1+1
- = $\frac{\sin \theta [\operatorname{cosec} \theta - \cot \theta + \operatorname{cosec} \theta + \cot \theta]}{\operatorname{cosec}^2 \theta - \cot^2 \theta} = \frac{\sin \theta (2 \operatorname{cosec} \theta)}{1} = 2$ $1\frac{1}{2}$
- Hence $\frac{\sin \theta}{\operatorname{cosec} \theta + \cot \theta} = 2 + \frac{\sin \theta}{\cot \theta - \operatorname{cosec} \theta}$ $\frac{1}{2}$
29. Let $-82 = a_n \therefore -82 = -7 + (n - 1)(-5)$ 1
- $\Rightarrow 15 = n - 1$ or $n = 16$ 1

$$\text{Again } -100 = a_m = -7 + (m - 1)(-5) \quad 1$$

$$\Rightarrow (m - 1)(-5) = -93$$

$$m - 1 = \frac{93}{5} \text{ or } m = \frac{93}{5} + 1 \notin \mathbb{N} \quad 1$$

\therefore -100 is not a term of the AP.

OR

$$S_n = 180 = \frac{n}{2} \cdot [90 + (n - 1)(-6)] \quad 1$$

$$360 = 90n - 6n^2 + 6n \Rightarrow 6n^2 - 96n + 360 = 0 \quad 1$$

$$\Rightarrow 6[(n - 6)(n - 10)] = 0 \Rightarrow n = 6, n = 10 \quad 1$$

$$\text{Sum of } a_7, a_8, a_9, a_{10} = 0 \therefore n = 6 \text{ or } n = 10 \quad 1$$

30. Let marks in Hindi be x

$$\text{Then marks in Eng} = 30 - x \quad \frac{1}{2}$$

$$\therefore (x + 2)(30 - x - 3) = 210 \quad 1$$

$$\Rightarrow x^2 - 25x + 156 = 0 \text{ or } (x - 13)(x - 12) = 0 \quad 1$$

$$\Rightarrow x = 13 \text{ or } x = 12$$

$$\therefore 30 - 13 = 17 \text{ or } 30 - 12 = 18 \quad 1$$

\therefore Marks in Hindi & English are

$$(13, 17) \text{ or } (12, 18) \quad \frac{1}{2}$$

QUESTION PAPER CODE 30/2/2
EXPECTED ANSWER/VALUE POINTS

SECTION A

1. Point on x-axis is (2, 0) 1

2. $\sin^2 60^\circ + 2 \tan 45^\circ - \cos^2 30^\circ = \left(\frac{\sqrt{3}}{2}\right)^2 + 2(1) - \left(\frac{\sqrt{3}}{2}\right)^2$ [For any two correct values] $\frac{1}{2}$

$$= 2$$

$\frac{1}{2}$

OR

$$\sin A = \frac{3}{4} \Rightarrow \cos A = \sqrt{1 - \frac{9}{16}} = \frac{\sqrt{7}}{4}$$
 $\frac{1}{2}$

$$\sec A = \frac{4}{\sqrt{7}}$$
 $\frac{1}{2}$

3. ΔABC : Isosceles $\Delta \Rightarrow AC = BC = 4$ cm. $\frac{1}{2}$

$$AB = \sqrt{4^2 + 4^2} = 4\sqrt{2}$$
 $\frac{1}{2}$

OR

$$\frac{AD}{BD} = \frac{AE}{CE} \Rightarrow \frac{AD}{7.2} = \frac{1.8}{5.4}$$
 $\frac{1}{2}$

$$\therefore AD = \frac{7.2 \times 1.8}{5.4} = 2.4 \text{ cm.}$$
 $\frac{1}{2}$

4. $\frac{3-a}{3a} - \frac{1}{a} = \frac{3-a-3}{3a} = -\frac{1}{3}$ 1

5. LCM (336, 54) = $\frac{336 \times 54}{6}$ $\frac{1}{2}$

$$= 336 \times 9 = 3024$$
 $\frac{1}{2}$

6. $a = -4\frac{1}{2}$, $d = 1\frac{1}{2}$, $\therefore a_{21} = -\frac{9}{2} + 20\left(\frac{3}{2}\right)$ $\frac{1}{2}$

$$= \frac{51}{2}$$
 $\frac{1}{2}$

SECTION B

7. For infinitely many solutions,

$$\frac{2}{k+2} = \frac{3}{-3(1-k)} = \frac{7}{5k+1} \quad 1$$

$$\Rightarrow 2k - 2 = k + 2 \text{ or } 5k + 1 = 7k - 7$$

$$\Rightarrow k = 4 \quad \Rightarrow 2k = 8 \quad \Rightarrow k = 4$$

$$\text{Hence } k = 4. \quad 1$$

8. Maximum frequency = 50, class (modal) = 35 – 40. 1/2

$$\begin{aligned} \text{Mode} &= L + \left(\frac{f_1 - f_0}{2f_1 - f_0 - f_2} \right) \times h \\ &= 35 + \frac{50 - 34}{100 - 34 - 42} \times 5 \quad 1 \end{aligned}$$

$$= 35 + \frac{16}{24} \times 5 = 38.33 \quad \frac{1}{2}$$

9. Let larger angle be x°

$$\therefore \text{Smaller angle} = 180^\circ - x^\circ \quad \frac{1}{2}$$

$$\therefore (x) - (180 - x) = 18 \quad \frac{1}{2}$$

$$2x = 180 + 18 = 198 \Rightarrow x = 99 \quad \frac{1}{2}$$

$$\therefore \text{The two angles are } 99^\circ, 81^\circ \quad \frac{1}{2}$$

OR

Let Son's present age be x years

Then Sumit's present age = $3x$ years. 1/2

$$\therefore \text{5 Years later, we have, } 3x + 5 = \frac{5}{2}(x + 5) \quad \frac{1}{2}$$

$$6x + 10 = 5x + 25 \Rightarrow x = 15 \quad \frac{1}{2}$$

$$\therefore \text{Sumit's present age} = 45 \text{ years} \quad \frac{1}{2}$$

10. $P(\text{blue marble}) = \frac{1}{5}$, $P(\text{black marble}) = \frac{1}{4}$

$$\therefore P(\text{green marble}) = 1 - \left(\frac{1}{5} + \frac{1}{4} \right) = \frac{11}{20}$$

Let total number of marbles be x

$$\text{then } \frac{11}{20} \times x = 11 \Rightarrow x = 20$$

11. A, B, C are collinear \Rightarrow ar. $(\Delta ABC) = 0$

$$\therefore \frac{1}{2}[x(6-3) - 4(3-y) - 2(y-6)] = 0$$

$$\Rightarrow 3x + 2y = 0$$

OR

$$\text{Area of triangle} = \frac{1}{2}[1(6+5) - 4(-5+1) - 3(-1-6)]$$

$$= \frac{1}{2}[11+16+21] = \frac{48}{2} = 24 \text{ sq. units.}$$

12. Smallest number divisible by 306 and 657 = LCM (306, 657)

$$\text{LCM (306, 657)} = 22338$$

SECTION C

13. $\frac{XA}{XY} = \frac{2}{5} \Rightarrow \frac{XA}{AY} = \frac{2}{3}$

$$\therefore \text{Coords. of A are } \left(\frac{-8+18}{5}, \frac{-2-18}{5} \right) \text{ i.e. } (2, -4)$$

$$\text{A lies on } 3x + k(y+1) = 0$$

$$\Rightarrow 6 + k(-3) = 0 \Rightarrow k = 2.$$

14. $x^2 + 5x - (a+3)(a-2) = 0$

$$x^2 + (a+3)x - (a-2)x - (a+3)(a-2) = 0$$

$$[x + (a+3)][x - (a-2)] = 0$$

$$\Rightarrow x = (a-2) \text{ or } x = -(a+3)$$

Alternate method:

$$x^2 + 5x - (a^2 + a - 6) = 0$$

$$x = \frac{-5 \pm \sqrt{25 + 4(a^2 + a - 6)}}{2} \quad 1$$

$$= \frac{-5 \pm (2a + 1)}{2} \quad 1$$

$$x = (a - 2), -(a + 3) \quad 1$$

15. $A + 2B = 60^\circ$ and $A + 4B = 90^\circ$ 1+1

Solving to get $B = 15^\circ$ and $A = 30^\circ$ 1

16. Let $2 + 5\sqrt{3} = a$, where 'a' is a rational number. $\frac{1}{2}$

than $\sqrt{3} = \frac{a - 2}{5}$ 1

Which is a contradiction as LHS is irrational and RHS is rational 1

$\therefore 2 + 5\sqrt{3}$ can not be rational $\frac{1}{2}$

Hence $2 + 5\sqrt{3}$ is irrational.

Alternate method:

Let $2 + 5\sqrt{3}$ be rational $\frac{1}{2}$

$\therefore 2 + 5\sqrt{3} = \frac{p}{q}$, p, q are integers, $q \neq 0$

$\Rightarrow \sqrt{3} = \left(\frac{p}{q} - 2\right) \div 5 = \frac{p - 2q}{5q}$ 1

LHS is irrational and RHS is rational

which is a contradiction 1

$\therefore 2 + 5\sqrt{3}$ is irrational. $\frac{1}{2}$

OR

$$2048 = 960 \times 2 + 128$$

$$960 = 128 \times 7 + 64$$


$$128 = 64 \times 2 + 0$$

$$\therefore \text{HCF}(2048, 960) = 64$$

2

1

17.



Correct Figure

$\triangle APB \sim \triangle DPC$ [AA similarity]

$$\frac{AP}{DP} = \frac{BP}{PC}$$

$$\Rightarrow AP \times PC = BP \times DP$$

$\frac{1}{2}$

1

1

$\frac{1}{2}$

OR

Correct Figure

In $\triangle POQ$ and $\triangle ROS$

$$\left. \begin{aligned} \angle P &= \angle R \\ \angle Q &= \angle S \end{aligned} \right\} \text{alt. } \angle s$$

$\therefore \triangle POQ \sim \triangle ROS$ [AA similarity]

$$\therefore \frac{\text{ar}(\triangle POQ)}{\text{ar}(\triangle ROS)} = \left(\frac{PQ}{RS}\right)^2$$

$$= \left(\frac{3}{1}\right)^2 = \frac{9}{1}$$

$$\therefore \text{ar}(\triangle POQ) : \text{ar}(\triangle ROS) = 9 : 1$$


$\frac{1}{2}$

1

1

$\frac{1}{2}$

18.



Correct Figure

$\triangle AOD \cong \triangle COE$ [SAS]

$$\Rightarrow \angle 1 = \angle 2$$

Similarly $\angle 4 = \angle 3$

$$\Rightarrow \angle 1 + \angle 4 = \angle 2 + \angle 3 = \frac{1}{2}(180^\circ)$$

$$\Rightarrow \angle 2 + \angle 3 = 90^\circ \text{ or } \angle AOB = 90^\circ$$

$\frac{1}{2}$


1

$\frac{1}{2}$

$\frac{1}{2}$

$\frac{1}{2}$

Alternate method:



Correct Figure

$\frac{1}{2}$

$\triangle OAD \cong \triangle OAC$ [SAS]

$\Rightarrow \angle 1 = \angle 2$

1

Similarly $\angle 4 = \angle 3$

$\frac{1}{2}$

But $\angle 1 + \angle 2 + \angle 3 + \angle 4 = 180^\circ$ [$\because PQ \parallel RS$]

$\Rightarrow \angle 2 + \angle 3 = \angle 1 + \angle 4 = \frac{1}{2}(180^\circ) = 90^\circ$

$\frac{1}{2}$

\therefore In $\triangle AOB$, $\angle AOB = 180^\circ - (\angle 2 + \angle 3) = 90^\circ$

$\frac{1}{2}$

19. Radius of quadrant = $OB = \sqrt{15^2 + 15^2} = 15\sqrt{2}$ cm.

1

Shaded area = Area of quadrant – Area of square

$\frac{1}{2}$

$= \frac{1}{4}(3.14)[(15\sqrt{2})^2 - (15)^2]$

1

$= (15)^2 (1.57 - 1) = 128.25 \text{ cm}^2$

$\frac{1}{2}$

OR

$BD = \sqrt{(2\sqrt{2})^2 + (2\sqrt{2})^2} = \sqrt{16} = 4 \text{ cm}$

1

\therefore Radius of circle = 2 cm

$\frac{1}{2}$

\therefore Shaded area = Area of circle – Area of square

$\frac{1}{2}$

$= 3.14 \times 2^2 - (2\sqrt{2})^2$


$= 12.56 - 8 = 4.56 \text{ cm}^2$

1

20.

Height of cylinder = $20 - 7 = 13$ cm.

1



Total volume = $\pi \left(\frac{7}{2}\right)^2 \cdot 13 + \frac{4}{3} \pi \left(\frac{7}{2}\right)^2 \text{ cm}^3$

1

$= \frac{22}{7} \times \frac{49}{4} \left(13 + \frac{4}{3} \cdot \frac{7}{2}\right) \text{ cm}^3$

$= \frac{77 \times 53}{6} = 680.17 \text{ cm}^3$

1

$$21. \quad x_i: \quad 32.5 \quad 37.5 \quad 42.5 \quad 47.5 \quad 52.5 \quad 57.5 \quad 62.5$$

 $\frac{1}{2}$

$$f_i: \quad 14 \quad 16 \quad 28 \quad 23 \quad 18 \quad 8 \quad 3 \quad \Sigma f_i = 110$$

 $\frac{1}{2}$

$$u_i: \quad -3 \quad -2 \quad -1 \quad 0 \quad 1 \quad 2 \quad 3$$

$$f_i u_i: \quad -42 \quad -32 \quad -28 \quad 0 \quad 18 \quad 16 \quad 9, \quad \Sigma f_i u_i = -59$$

1

$$\text{Mean} = 47.5 - \frac{59 \times 5}{110} = 47.5 - 2.68 = 44.82$$

1

Note: If N is taken as 100, Ans. 44.55

Accept.

If some one write, data is wrong, give full 3 marks.

$$22. \quad \begin{array}{r} 3x^2 - 5 \overline{) 3x^4 - 9x^3 + x^2 + 15x + k} \quad (x^2 - 3x + 2 \\ \underline{3x^4} \\ -9x^3 \\ \underline{-9x^3} \\ 6x^2 + k \\ \underline{6x^2 - 10} \\ k + 10 \end{array}$$

2

$$\therefore k + 10 = 0 \Rightarrow k = -10$$

1

OR

$$p(y) = 7y^2 - \frac{11}{3}y - \frac{2}{3} = \frac{1}{3}(21y^2 - 11y - 2)$$

$$= \frac{1}{3}[(7y+1)(3y-2)]$$

1

$$\therefore \text{Zeroes are } 2/3, -1/7$$

 $\frac{1}{2}$

$$\text{Sum of zeroes} = \frac{2}{3} - \frac{1}{7} = \frac{11}{21}$$

$$\frac{-b}{a} = \frac{11}{21} \therefore \text{sum of zeroes} = \frac{-b}{a}$$

1

$$\text{Product of zeroes} = \left(\frac{2}{3}\right)\left(-\frac{1}{7}\right) = -\frac{2}{21}$$


$$\frac{c}{a} = -\frac{2}{3}\left(\frac{1}{7}\right) = -\frac{2}{21} \therefore \text{Product} = \frac{c}{a} \quad \frac{1}{2}$$

SECTION D

23. For correct given, to prove, construction and figure 4 × $\frac{1}{2}$ = 2

For correct proof. 2

24. Correct Figure 1



In $\triangle ABP$, $\frac{h}{x} = \tan 60^\circ = \sqrt{3}$...(i) $\frac{1}{2}$

In $\triangle CDP$, $\frac{h}{80 - x} = \tan 30^\circ = \frac{1}{\sqrt{3}}$...(ii) $\frac{1}{2}$

dividing (i) by (ii) we get $\frac{80 - x}{x} = \frac{3}{1}$

$\Rightarrow 3x = 80 - x$ or $4x = 80 \Rightarrow x = 20$ m. 1

and $h = 20\sqrt{3}$ m. $\frac{1}{2}$

\therefore Height of poles is $20\sqrt{3}$ m

and P is at distances 20 m and 60 m from poles $\frac{1}{2}$

25. Let total length of cloth = l m.

\therefore Rate per metre = ₹ $\frac{200}{l}$ $\frac{1}{2}$

$\Rightarrow (l + 5)\left(\frac{200}{l} - 2\right) = 200$ 1

$\Rightarrow (l + 5)(200 - 2l) = 200l \Rightarrow l^2 + 5l - 500 = 0$ 1

$\Rightarrow (l + 25)(l - 20) = \Rightarrow l = 20$ m. 1

\therefore Rate per metre = ₹ $\frac{200}{20} = ₹ 10$ per metre $\frac{1}{2}$

26. Let $-82 = a_n \therefore -82 = -7 + (n - 1)(-5)$ 1
 $\Rightarrow 15 = n - 1$ or $n = 16$ 1
 Again $-100 = a_m = -7 + (m - 1)(-5)$ 1
 $\Rightarrow (m - 1)(-5) = -93$
 $m - 1 = \frac{93}{5}$ or $m = \frac{93}{5} + 1 \notin \mathbb{N}$ 1
 $\therefore -100$ is not a term of the AP.

OR

$$S_n = 180 = \frac{n}{2} \cdot [90 + (n - 1)(-6)]$$
 1

$$360 = 90n - 6n^2 + 6n \Rightarrow 6n^2 - 96n + 360 = 0$$
 1

$$\Rightarrow 6[(n - 6)(n - 10)] = 0 \Rightarrow n = 6, n = 10$$
 1

$$\text{Sum of } a_7, a_8, a_9, a_{10} = 0 \therefore n = 6 \text{ or } n = 10$$
 1

27. LHS = $\frac{\tan \theta}{1 - \frac{1}{\tan \theta}} + \frac{1}{1 - \tan \theta} = \frac{\tan^2 \theta}{\tan \theta - 1} - \frac{1}{\tan \theta (\tan \theta - 1)}$ 1

$$= \frac{\tan^3 \theta - 1}{\tan \theta (\tan \theta - 1)} = \frac{(\tan \theta - 1)(\tan^2 \theta + \tan \theta + 1)}{\tan \theta (\tan \theta - 1)}$$
 1

$$= \tan \theta + 1 + \cot \theta = 1 + \frac{\sin \theta}{\cos \theta} + \frac{\cos \theta}{\sin \theta} = 1 + \frac{\sin^2 \theta + \cos^2 \theta}{\sin \theta \cos \theta}$$
 1

$$= 1 + \frac{1}{\sin \theta \cos \theta} = 1 + \operatorname{cosec} \theta \sec \theta = \text{RHS}$$
 1

OR

Consider

$$\frac{\sin \theta}{\operatorname{cosec} \theta + \cot \theta} - \frac{\sin \theta}{\cot \theta - \operatorname{cosec} \theta} = \frac{\sin \theta}{\operatorname{cosec} \theta + \cot \theta} + \frac{\sin \theta}{\operatorname{cosec} \theta - \cot \theta}$$
 1+1

$$= \frac{\sin \theta [\operatorname{cosec} \theta - \cot \theta + \operatorname{cosec} \theta + \cot \theta]}{\operatorname{cosec}^2 \theta - \cot^2 \theta} = \frac{\sin \theta (2 \operatorname{cosec} \theta)}{1} = 2$$
 1 \frac{1}{2}

$$\text{Hence } \frac{\sin \theta}{\operatorname{cosec} \theta + \cot \theta} = 2 + \frac{\sin \theta}{\cot \theta - \operatorname{cosec} \theta}$$
 \frac{1}{2}

28.	Less than 40	less than 50	less than 60	less than 70	less than 80	less than 90	less than 100	$\frac{1}{2}$
cf.	7	12	20	30	36	42	50	1

Plotting of points (40, 7), (50, 12), (60, 20), (70, 30), (80, 36), (90, 42) and (100, 50) $1\frac{1}{2}$

Joining the points to get the curve 1

29. Constructing an equilateral triangle of side 5 cm 1

Constructing another similar Δ with scale factor $\frac{2}{3}$ 3


OR

Constructing two concentric circle of radii 2 cm and 5 cm 1

Drawing two tangents PA and PB 2

PA = 4.5 cm (approx) 1

30.



Total volume = $3.14 (12)^2 (220) + 3.14(8)^2(60) \text{ cm}^3$ 1

= $99475.2 + 12057.6 = 111532.8 \text{ cm}^3$ 1

Mass = $\frac{111532.8 \times 8}{1000} \text{ kg}$ 1

= 892.262 kg 1

QUESTION PAPER CODE 30/2/3
EXPECTED ANSWER/VALUE POINTS

SECTION A

1. $D = (4\sqrt{3})^2 - 4(4)(3) = 0$ $\frac{1}{2}$

\therefore Roots are real and equal. $\frac{1}{2}$

2. $\sin^2 60^\circ + 2 \tan 45^\circ - \cos^2 30^\circ = \left(\frac{\sqrt{3}}{2}\right)^2 + 2(1) - \left(\frac{\sqrt{3}}{2}\right)^2$ [For any two correct values] $\frac{1}{2}$

$= 2$ $\frac{1}{2}$

OR

$\sin A = \frac{3}{4} \Rightarrow \cos A = \sqrt{1 - \frac{9}{16}} = \frac{\sqrt{7}}{4}$ $\frac{1}{2}$

$\sec A = \frac{4}{\sqrt{7}}$ $\frac{1}{2}$

3. Point on x-axis is (2, 0) 1

4. $\triangle ABC$: Isosceles $\triangle \Rightarrow AC = BC = 4$ cm. $\frac{1}{2}$

$AB = \sqrt{4^2 + 4^2} = 4\sqrt{2}$ cm $\frac{1}{2}$

OR

$\frac{AD}{BD} = \frac{AE}{CE} \Rightarrow \frac{AD}{7.2} = \frac{1.8}{5.4}$ $\frac{1}{2}$

$\therefore AD = \frac{7.2 \times 1.8}{5.4} = 2.4$ cm. $\frac{1}{2}$

5. $2x^2 - 4x + 3 = 0 \Rightarrow D = 16 - 24 = -8$ $\frac{1}{2}$

\therefore Equation has NO real roots $\frac{1}{2}$

6. $\text{LCM}(336, 54) = \frac{336 \times 54}{6}$ $\frac{1}{2}$

$= 336 \times 9 = 3024$ $\frac{1}{2}$

SECTION B

7. $E_1 : \{(1, 5), (2, 5), (3, 5), (4, 5), (5, 5), (6, 5), (5, 1), (5, 2), (5, 3), (5, 4), (5, 6)\}$

$$\therefore P(5 \text{ will come at least once}) = P(E_1) = \frac{11}{36} \quad 1$$

$$P(5 \text{ will not come either time}) = 1 - \frac{11}{36} = \frac{25}{36} \quad 1$$

8. Maximum frequency = 50, class (modal) = 35 – 40. 1/2

$$\text{Mode} = L + \left(\frac{f_1 - f_0}{2f_1 - f_0 - f_2} \right) \times h$$

$$= 35 + \frac{50 - 34}{100 - 34 - 42} \times 5 \quad 1$$

$$= 35 + \frac{16}{24} \times 5 = 38.33 \quad \frac{1}{2}$$

9. Let larger angle be x°

$$\therefore \text{Smaller angle} = 180^\circ - x^\circ \quad \frac{1}{2}$$

$$\therefore (x) - (180 - x) = 18 \quad \frac{1}{2}$$

$$2x = 180 + 18 = 198 \Rightarrow x = 99 \quad \frac{1}{2}$$

$$\therefore \text{The two angles are } 99^\circ, 81^\circ \quad \frac{1}{2}$$

OR

Let Son's present age be x years

Then Sumit's present age = $3x$ years. 1/2

$$\therefore \text{5 Years later, we have, } 3x + 5 = \frac{5}{2}(x + 5) \quad \frac{1}{2}$$

$$6x + 10 = 5x + 25 \Rightarrow x = 15 \quad \frac{1}{2}$$

$$\therefore \text{Sumit's present age} = 45 \text{ years} \quad \frac{1}{2}$$

10. A, B, C are collinear \Rightarrow ar. (ΔABC) = 0

 $\frac{1}{2}$

$$\therefore \frac{1}{2}[x(6-3) - 4(3-y) - 2(y-6)] = 0$$

1

$$\Rightarrow 3x + 2y = 0$$

 $\frac{1}{2}$

OR

$$\text{Area of triangle} = \frac{1}{2}[1(6+5) - 4(-5+1) - 3(-1-6)]$$

1

$$= \frac{1}{2}[11+16+21] = \frac{48}{2} = 24 \text{ sq. units.}$$

1

11. For unique solution $\frac{1}{3} \neq \frac{2}{k}$

1

$$\Rightarrow k \neq 6$$

1

12. Smallest number divisible by 306 and 657 = LCM (306, 657)

1

$$\text{LCM (306, 657)} = 22338$$

1

SECTION C

13.

Any point on y-axis is P(0, y)

1

$$\begin{array}{c} \text{A} \quad \quad \quad \text{P} \quad \quad \quad \text{B} \\ \text{---} \quad \quad \quad \text{---} \quad \quad \quad \text{---} \\ (-1, -4) \quad (0, y) \quad (5, -6) \end{array}$$

Let P divides AB in k : 1

$$\Rightarrow 0 = \frac{5k-1}{k+1} \Rightarrow k = \frac{1}{5} \text{ i.e. } 1:5$$

1

$$\Rightarrow y = \frac{-6k-4}{k+1} = \frac{-\frac{6}{5}-4}{\frac{1}{5}+1} = \frac{-\frac{26}{5}}{\frac{6}{5}} = \frac{-13}{3}$$

1

$$\Rightarrow \text{P is } \left(0, \frac{-13}{3}\right)$$

14. Given expression = $\left(\frac{3 \tan 41^\circ}{\tan 41^\circ}\right)^2 - \left(\frac{\sin 35^\circ \operatorname{cosec} 35^\circ}{\tan 10^\circ \tan 20^\circ (\sqrt{3}) \cot 20^\circ \cot 10^\circ}\right)^2$

 $1 \frac{1}{2}$

$$= 9 - \frac{1}{3} = \frac{26}{3}$$

 $1 \frac{1}{2}$

15. Radius of first sphere = 3 cm $\therefore \frac{4}{3}\pi(3)^3 d = 1$ {d = density} $\frac{1}{2}$

let radius of 2nd sphere be r cm $\therefore \frac{4}{3}\pi(r)^3 \cdot d = 7 \Rightarrow r^3 = 7(3)^3$ $\frac{1}{2}$

$\Rightarrow \frac{4}{3}\pi(3)^3 + \frac{4}{3}\pi \cdot (3)^3 \cdot 7 = \frac{4}{3}\pi R^3$ 1

$\Rightarrow R^3 = (3)^3 (1 + 7) \Rightarrow R = 3(2) = 6$ $\frac{1}{2}$

\therefore Diameter = 12 cm. $\frac{1}{2}$

16. Let $2 + 5\sqrt{3} = a$, where 'a' is a rational number. $\frac{1}{2}$

then $\sqrt{3} = \frac{a-2}{5}$ 1

Which is a contradiction as LHS is irrational and RHS is rational 1

$\therefore 2 + 5\sqrt{3}$ can not be rational $\frac{1}{2}$

Hence $2 + 5\sqrt{3}$ is irrational.

Alternate method:

Let $2 + 5\sqrt{3}$ be rational $\frac{1}{2}$

$\therefore 2 + 5\sqrt{3} = \frac{p}{q}$, p, q are integers, $q \neq 0$

$\Rightarrow \sqrt{3} = \left(\frac{p}{q} - 2\right) \div 5 = \frac{p-2q}{5q}$ 1

LHS is irrational and RHS is rational

which is a contradiction 1

$\therefore 2 + 5\sqrt{3}$ is irrational. $\frac{1}{2}$

OR

$2048 = 960 \times 2 + 128$


$960 = 128 \times 7 + 64$ 2

$$128 = 64 \times 2 + 0$$

$$\therefore \text{HCF}(2048, 960) = 64$$

1

17.



Correct Figure

$\frac{1}{2}$

$\triangle APB \sim \triangle DPC$ [AA similarity]

1

$$\frac{AP}{DP} = \frac{BP}{PC}$$

1


$$\Rightarrow AP \times PC = BP \times DP$$

$\frac{1}{2}$

OR

Correct Figure

$\frac{1}{2}$



In $\triangle POQ$ and $\triangle ROS$

$$\left. \begin{array}{l} \angle P = \angle R \\ \angle Q = \angle S \end{array} \right\} \text{alt. } \angle s$$

$\therefore \triangle POQ \sim \triangle ROS$ [AA similarity]

1

$$\therefore \frac{\text{ar}(\triangle POQ)}{\text{ar}(\triangle ROS)} = \left(\frac{PQ}{RS}\right)^2$$


1

$$= \left(\frac{3}{1}\right)^2 = \frac{9}{1}$$

$\frac{1}{2}$

$$\therefore \text{ar}(\triangle POQ) : \text{ar}(\triangle ROS) = 9 : 1$$

18.



Correct Figure

$\frac{1}{2}$

$\triangle AOD \cong \triangle BOC$ [SAS]

1

$$\Rightarrow \angle 1 = \angle 2$$

$\frac{1}{2}$

Similarly $\angle 4 = \angle 3$


$\frac{1}{2}$

$$\Rightarrow \angle 1 + \angle 4 = \angle 2 + \angle 3 = \frac{1}{2}(180^\circ)$$

$$\Rightarrow \angle 2 + \angle 3 = 90^\circ \text{ or } \angle AOB = 90^\circ$$

$\frac{1}{2}$

Alternate method:



Correct Figure

$$\triangle OAD \cong \triangle AOC \text{ [SAS]}$$

$$\Rightarrow \angle 1 = \angle 2$$

Similarly $\angle 4 = \angle 3$

$$\text{But } \angle 1 + \angle 2 + \angle 3 + \angle 4 = 180^\circ \text{ [}\because \text{PQ} \parallel \text{RS}]$$

$$\Rightarrow \angle 2 + \angle 3 = \angle 1 + \angle 4 = \frac{1}{2}(180^\circ) = 90^\circ$$

$$\therefore \text{ In } \triangle AOB, \angle AOB = 180^\circ - (\angle 2 + \angle 3) = 90^\circ$$

19. Radius of quadrant = $OB = \sqrt{15^2 + 15^2} = 15\sqrt{2}$ cm.

Shaded area = Area of quadrant – Area of square

$$= \frac{1}{4}(3.14)[(15\sqrt{2})^2 - (15)^2]$$

$$= (15)^2 (1.57 - 1) = 128.25 \text{ cm}^2$$

OR

$$BD = \sqrt{(2\sqrt{2})^2 + (2\sqrt{2})^2} = \sqrt{16} = 4 \text{ cm}$$

\therefore Radius of circle = 2 cm

\therefore Shaded area = Area of circle – Area of square

$$= 3.14 \times 2^2 - (2\sqrt{2})^2$$

$$= 12.56 - 8 = 4.56 \text{ cm}^2$$

20. $x^2 + px + 16 = 0$ have equal roots if $D = p^2 - 4(16)(1) = 0$

$$p^2 = 64 \Rightarrow p = \pm 8$$

$$\therefore x^2 \pm 8x + 16 = 0 \Rightarrow (x \pm 4)^2 = 0$$

$$x \pm 4 = 0$$

\therefore Roots are $x = -4$ and $x = 4$

$$\begin{array}{r}
 21. \quad 3x^2 - 5 \overline{) 3x^4 - 9x^3 + x^2 + 15x + k} \quad (x^2 - 3x + 2 \\
 \underline{3x^4 \quad - 5x^2} \\
 -9x^3 + 6x^2 + 15x + k \\
 \underline{-9x^3 \quad + 15x} \\
 6x^2 + k \\
 \underline{6x^2 - 10} \\
 k + 10
 \end{array}$$

2

$$\therefore k + 10 = 0 \Rightarrow k = -10$$

1

OR

$$p(y) = 7y^2 - \frac{11}{3}y - \frac{2}{3} = \frac{1}{3}(21y^2 - 11y - 2)$$

$$= \frac{1}{3}[(7y+1)(3y-2)]$$

1

$$\therefore \text{Zeroes are } 2/3, -1/7$$

 $\frac{1}{2}$

$$\text{Sum of zeroes} = \frac{2}{3} - \frac{1}{7} = \frac{11}{21}$$

$$\frac{-b}{a} = \frac{11}{21} \therefore \text{sum of zeroes} = \frac{-b}{a}$$

1

$$\text{Product of zeroes} = \left(\frac{2}{3}\right)\left(-\frac{1}{7}\right) = -\frac{2}{21}$$

$$\frac{c}{a} = -\frac{2}{3}\left(\frac{1}{7}\right) = -\frac{2}{21} \therefore \text{Product} = \frac{c}{a}$$

 $\frac{1}{2}$

$$22. \quad x_i: \quad 32.5 \quad 37.5 \quad 42.5 \quad 47.5 \quad 52.5 \quad 57.5 \quad 62.5$$

 $\frac{1}{2}$

$$f_i: \quad 14 \quad 16 \quad 28 \quad 23 \quad 18 \quad 8 \quad 3 \quad \Sigma f_i = 110$$

 $\frac{1}{2}$

$$u_i: \quad -3 \quad -2 \quad -1 \quad 0 \quad 1 \quad 2 \quad 3$$

$$f_i u_i: \quad -42 \quad -32 \quad -28 \quad 0 \quad 18 \quad 16 \quad 9, \quad \Sigma f_i u_i = -59$$

1

$$\text{Mean} = 47.5 - \frac{59 \times 5}{110} = 47.5 - 2.68 = 44.82$$

1

Note: If N is taken as 100, Ans. 44.55

Accept.

If some one write, data is wrong, give full 3 marks.

SECTION D

23. For correct given, to prove, const. and figure

$$4 \times \frac{1}{2} = 2$$

For correct proof.


2

24.

In ΔPAC ,

Correct Figure

1



$$\frac{AC}{AP} = \tan 45^\circ = 1$$

1

$$\Rightarrow x + 5 = y$$

$\frac{1}{2}$

$$\text{In } \Delta PAB, \frac{x}{y} = \tan 30^\circ = \frac{1}{\sqrt{3}}$$

$$\frac{x}{x+5} = \frac{1}{\sqrt{3}} \Rightarrow x = \frac{5}{\sqrt{3}-1} = \frac{5(\sqrt{3}+1)}{3} = 6.83$$

$1 \frac{1}{2}$

\therefore Height of tower = 6.83 m

25. Volume of ice-cream in the cylinder = $\pi(6)^2 \cdot 15 \text{ cm}^3$

1

$$\text{Volume of ice-cream in one cone} = \frac{1}{3} \pi r^2 \cdot 4r + \frac{2}{3} \pi r^3 \text{ cm}^3$$

(Given $h = 4r$)

1

$$= 2\pi r^3 \text{ cm}^3$$

$\frac{1}{2}$

$$\Rightarrow 10(2\pi r^3) = \pi(6)^2 \times 15$$

1

$$\Rightarrow r^3 = (3)^3 \Rightarrow r = 3 \text{ cm.}$$

$\frac{1}{2}$

26. Let marks in Hindi be x

$$\text{Then marks in Eng} = 30 - x$$

$\frac{1}{2}$

$$\therefore (x + 2)(30 - x - 3) = 210$$

1

$$\Rightarrow x^2 - 25x + 156 = 0 \text{ or } (x - 13)(x - 12) = 0$$

1

$$\Rightarrow x = 13 \text{ or } x = 12$$

$$\therefore 30 - 13 = 17 \text{ or } 30 - 12 = 18 \quad 1$$

\therefore Marks in Hindi & English are

$$(13, 17) \text{ or } (12, 18) \quad \frac{1}{2}$$

27. Let $-82 = a_n \therefore -82 = -7 + (n - 1)(-5)$ 1

$$\Rightarrow 15 = n - 1 \text{ or } n = 16 \quad 1$$

Again $-100 = a_m = -7 + (m - 1)(-5)$ 1

$$\Rightarrow (m - 1)(-5) = -93$$

$$m - 1 = \frac{93}{5} \text{ or } m = \frac{93}{5} + 1 \notin \mathbb{N} \quad 1$$

\therefore -100 is not a term of the AP.

OR

$$S_n = 180 = \frac{n}{2} \cdot [90 + (n - 1)(-6)] \quad 1$$

$$360 = 90n - 6n^2 + 6n \Rightarrow 6n^2 - 96n + 360 = 0 \quad 1$$

$$\Rightarrow 6[(n - 6)(n - 10)] = 0 \Rightarrow n = 6, n = 10 \quad 1$$

$$\text{Sum of } a_7, a_8, a_9, a_{10} = 0 \therefore n = 6 \text{ or } n = 10 \quad 1$$

28. LHS = $\frac{\tan \theta}{1 - \frac{1}{\tan \theta}} + \frac{1}{1 - \tan \theta} = \frac{\tan^2 \theta}{\tan \theta - 1} - \frac{1}{\tan \theta (\tan \theta - 1)}$ 1

$$= \frac{\tan^3 \theta - 1}{\tan \theta (\tan \theta - 1)} = \frac{(\tan \theta - 1)(\tan^2 \theta + \tan \theta + 1)}{\tan \theta (\tan \theta - 1)} \quad 1$$

$$= \tan \theta + 1 + \cot \theta = 1 + \frac{\sin \theta}{\cos \theta} + \frac{\cos \theta}{\sin \theta} = 1 + \frac{\sin^2 \theta + \cos^2 \theta}{\sin \theta \cos \theta} \quad 1$$

$$= 1 + \frac{1}{\sin \theta \cos \theta} = 1 + \operatorname{cosec} \theta \sec \theta = \text{RHS} \quad 1$$

OR

Consider

$$\frac{\sin \theta}{\operatorname{cosec} \theta + \cot \theta} - \frac{\sin \theta}{\cot \theta - \operatorname{cosec} \theta} = \frac{\sin \theta}{\operatorname{cosec} \theta + \cot \theta} + \frac{\sin \theta}{\operatorname{cosec} \theta - \cot \theta} \quad 1+1$$

$$= \frac{\sin \theta [\operatorname{cosec} \theta - \cot \theta + \operatorname{cosec} \theta + \cot \theta]}{\operatorname{cosec}^2 \theta - \cot^2 \theta} = \frac{\sin \theta (2 \operatorname{cosec} \theta)}{1} = 2 \quad 1 \frac{1}{2}$$

$$\text{Hence } \frac{\sin \theta}{\operatorname{cosec} \theta + \cot \theta} = 2 + \frac{\sin \theta}{\cot \theta - \operatorname{cosec} \theta} \quad \frac{1}{2}$$

29.	Less than 40	less than 50	less than 60	less than 70	less than 80	less than 90	less than 100	$\frac{1}{2}$
cf.	7	12	20	30	36	42	50	1

Plotting of points (40, 7), (50, 12), (60, 20), (70, 30), (80, 36), (90, 42) and (100, 50) $1 \frac{1}{2}$

Joining the points to get the curve 1

30. Constructing an equilateral triangle of side 5 cm 1

Constructing another similar Δ with scale factor $\frac{2}{3}$ 3

OR

Constructing two concentric circle of radii 2 cm and 5 cm 1

Drawing two tangents PA and PB 2

PA = 4.5 cm (approx) 1