



ಬೆಂಗಳೂರು ಗ್ರಾಮಾಂತರ ಜಿಲ್ಲಾ ಪಂಚಾಯತ್

ಸಾರ್ವಜನಿಕ ಶಿಕ್ಷಣ ಇಲಾಖೆ

ಉಪನಿರ್ದೇಶಕರ ಕಛೇರಿ (ಆಡಳಿತ), ಬೆಂಗಳೂರು ಗ್ರಾಮಾಂತರ ಜಿಲ್ಲೆ

ಸ್ಫೂರ್ತಿ

ಬಹುಆಯ್ಕೆ ಮಾದರಿ ಪ್ರಶ್ನೆಗಳು

ಎಸ್.ಎಸ್.ಎಲ್.ಸಿ 2020-21

ಗಣಿತ

(ಆಂಗ್ಲ ಮಾಧ್ಯಮ)



- 4) The n^{th} term in an Arithmetic progression with first term 'a' and the common difference 'd' is
 A) $a_n = a + n + 1$ B) $a_n = a + (n-1)d$ C) $a_n = a + (n+1)d$ D) $a_n = a - 1$
- 5) In an Arithmetic progression $a_n = 5n + 2$, then 1st term is
 A) 7 B) 6 C) 5 D) 8
- 6) In an Arithmetic progression $a_n = 3n + 5$, then the 5th term is
 A) 30 B) 28 C) 20 D) 13
- 7) An Arithmetic progression among the following is
 A) 10, 7, 4, 1, B) 5, 8, 12, C) 0, 3, 6, 10, D) 16, 8, 4, 2,
- 8) How many two digit numbers are divisible by 5
 A) 17 B) 18 C) 19 D) 20
- 9) An Arithmetic progression with first term 2 and the common difference 3 is
 A) 2, 5, 9, B) 2, 5, 8, ... C) 3, 5, 7, D) 2, 6, 10,
- 10) In an A.P, $a_n = 2n - 1$ then the product of first three term is
 A) 10 B) 12 C) 15 D) 18
- 11) 10th term of an Arithmetic progression 1, 4, 7, is
 A) 31 B) 27 C) 32 D) 28
- 12) The Sum of first 20 natural numbers is
 A) 210 B) 200 C) 190 D) 205
- 13) First term in the given Arithmetic progression 5, 8, 11, is
 A) 5 B) 8 C) 11 D) 2
- 14) An Arithmetic progression among the following is
 A) -37, -35, -33, -31 B) 2, 4, 8, 16 C) 4, 10, 16, 26 D) -5, -10, +10, +5
- 15) The common difference in an Arithmetic progression 10, 14, 18, is

- A) 10 B) 4 C) 18 D) -4

16) The next term of an Arithmetic progression 13, 10, 7..... Is

- A) 3 B) 7 C) 4 D) -4

17) In an A.P , $s_n = \frac{n}{2}(7n - 1)$, then the common difference is

- A) 3 B) 5 C) 7 D) 9

18) In an Arithmetic progression $a_n = 3n + 5$, then 8th term is

- A) 29 B) 19 C) 43 D) 26

19) In an Arithmetic progression $a_n = 6n - 4$, then the first term is

- A) 4 B) 2 C) -4 D) -2

20) In an Arithmetic progression $a_n = 2n + 3$, then the common difference is

- A) 5 B) 1 C) 3 D) 2

21) In an Arithmetic progression $a_n = 2n$ then the value of S_3 is

- A) 12 B) 2 C) 14 D) 6

22) The fourth term in an Arithmetic progression if $S_4 = 38$ and $S_3 = 24$ is

- A) 5 B) 14 C) 3 D) 8

23) The next four terms of an Arithmetic progression 6 , 9 , 12....are

- A) 16 , 19 , 22 , 25 B) 15 , 18 , 24 , 30
C) 15 , 18 , 21 , 24 D) 15 , 17 , 19 , 21

24) S_{10} in an Arithmetic progression if the sum of first 3 terms is 9 and sum of next 3 terms is 27 is

- A) 30 B) 38 C) 46 D) 36

25) The Sum of first n th terms of an Arithmetic progression with first term 'a' and the common difference 'd' is

- A) $S_n = n[2a + (n-1)d]$ B) $S_n = \frac{n}{2} [a + (n-1)d]$
C) $S_n = \frac{n}{2} [2a + (n+1)d]$ D) $S_n = \frac{n}{2} [2a + (n - 1)d]$

26) An Arithmetic progression with first term 3 and the common difference 4 is

A) 4,7,11,15

B) 3, 7, 11, 15

C) 3, 8, 13, 18

D) 3, 6, 9, 12

27) The sum of first n positive integers is

A) $S_n = \frac{(n+2)n}{2}$

B) $S_n = \frac{n(n+1)}{2}$

C) $S_n = \frac{n(n-1)}{2}$

D) $S_n = \frac{n(n-2)}{2}$

28) The common difference in An Arithmetic progression $2\sqrt{3}, 5\sqrt{3}, 8\sqrt{3}, \dots$ is

A) $3\sqrt{3}$

B) 3

C) $\sqrt{3}$

D) $\sqrt{6}$

29) The 10th term of an Arithmetic progression 4, 7, 10, ... is

A) 36

B) 31

C) 34

D) 40

30) Sum of first 10 odd natural numbers is

A) 120

B) 55

C) 110

D) 100

31) In A.P if $a=5, d=3, a_n=20$ then find the value of 'n'

A) 5

B) 6

C) 3

D) 7

32) The meaning of a_{10} in Arithmetic progression is ...

A) $a+10d$

B) $a+11d$

C) $a+9d$

D) $a-9d$

33) If 2, x, 14 are in Arithmetic progression, then the value of 'x' is

A) 28

B) 16

C) 7

D) 8

34) $\sqrt{7}, \sqrt{28}, \sqrt{63}, \dots$ are in Arithmetic progression, then the next term is

A) $\sqrt{112}$

B) $\sqrt{84}$

C) $\sqrt{98}$

D) $\sqrt{122}$

35) The sum of first 'n' even natural numbers is

A) $n(n+1)$

B) n^2

C) $\frac{n}{2}$

D) $n(n-1)$

36) The sum of first '10' even natural numbers is

A) 100

B) 90

C) 80

D) 110

37) The sum of first 'n' odd natural numbers is

A) 2n

B) n

C) n^2

D) n^3

38) a, b, c and d are in A.P, then c-b is equal to

A) (a-b),

B) (b-c)

C) (d-c)

D) (c-d)

ANSWERS:

1	C	11	D	21	A	31	B
2	B	12	A	22	B	32	C
3	A	13	A	23	C	33	D
4	B	14	A	24	D	34	A
5	A	15	B	25	D	35	A
6	C	16	C	26	B	36	D
7	A	17	C	27	B	37	C
8	C	18	A	28	B	38	D
9	B	19	B	29	B		
10	C	20	D	30	D		

Triangles


Points to remember

- ❖ Two figures are similar if and only if they have same shape but not necessarily the same size.
- ❖ Two polygons of same number of sides are similar If
 - 1) All the corresponding angles are equal.
 - 2) All the corresponding sides are in the same ratio or in a proportion.
- ❖ Two triangles are said to be similar if their corresponding angles are equal or corresponding sides are proportional.

drawn parallel to one side of a triangle, then it divides the other two sides proportionally."

- ❖ AA – Similarity criterion

If two triangles if the corresponding angles are equal, then their corresponding sides will be in proportion and hence the two triangles are similar.



In ΔLMN ಮತ್ತು ΔPQR ಗಳಲ್ಲಿ,

$$(i) \quad \angle L = \angle P, \quad \angle M = \angle Q, \quad \angle N = \angle R$$

$$(ii) \quad \frac{LM}{PQ} = \frac{MN}{QR} = \frac{LN}{PR}$$

$$\Rightarrow \quad \Delta LMN \sim \Delta PQR,$$


- ❖ The areas of similar triangles are proportional to square of the corresponding sides.
- ❖ Pythagoras theorem: In a right angled triangle the square on the hypotenuse is equal to the sum of the squares on the other two sides.
- ❖ Baudhayana theorem: The diagonal of a rectangle. Produces both areas of which its length and breadth produce separately.
- ❖ Pythagorean triplets: The triplets of natural numbers are form a right angled triangle are called pythagorean triplets
Ex:- i) 3, 4, 5 ii) 6, 8, 10
- ❖ Converse of pythagoras theorem:- "If the square on the longest side of a triangle is equal to the sum of the squares on the other two sides then those sides contain a right angle".

Multiple choice questions

1) In $\triangle ABC$, $DE \parallel AB$. If $CD=3\text{cm}$, $EC=4\text{cm}$, $BE=6\text{cm}$, then DA is equal to
 (A) 7.5 cm (B) 3 cm (C) 4.5 cm (D) 6 cm

2) In fig, if $XY \parallel BC$, then $\frac{AX}{XB} =$

- (A) $\frac{AX}{AY}$ (B) $\frac{AX}{AB}$ (C) $\frac{AY}{YC}$ (D) $\frac{AC}{AY}$



3) In a rectangle, length=8cm, breadth=6cm, then the length of its diagonal is equal to

- (A) 9 cm (B) 14 cm (C) 10 cm (D) 12 cm

4) $\triangle ABC \sim \triangle DEF$ and $\frac{BC}{EF} = \frac{3}{5}$ then, $\frac{\text{Area of } \triangle ABC}{\text{Area of } \triangle DEF} =$

- (A) $\frac{3}{5}$ (B) $\frac{9}{25}$ (C) $\frac{27}{125}$ (D) $\frac{6}{10}$

5) $\triangle ABC \sim \triangle PQR$ and $\frac{\text{Area of } \triangle ABC}{\text{Area of } \triangle PQR} = \frac{25}{81}$, then $BC:PQ$ is

- (A) 9: 5 (B) 5: 3 (C) 25: 81 (D) 5: 9

- 6) The length of the altitude of an equilateral triangle of side 10cm is
 (A) $5\sqrt{3}$ cm (B) $10\sqrt{3}$ cm (C) $\sqrt{3}$ cm (D) 75cm

(A) 16:81 (B) 4 : 9 (C) 2 : 3 (D) 8:18


- 8) In triangle PQR , $\angle PQR = 90^\circ$. PQ=12cm & QR=5cm , the length of PR=
 (A) 17 cm (B) 14 cm (C) 10 cm (D) 13 cm

- 9) Pythagorean triplet among these

(A) 4, 5 , 6 (B) 2, 3 , 5 (C) 8, 10, 6 (D) 9, 10, 12


- 10) In fig, D and E are the midpoints of AB and AC
 Respectively. If DE=4 cm then the value BC is

(A) 4 cm (B) 6 cm (C) 8 cm (D) 12 cm




- 11) In fig , if $XY \parallel BC$, then the value of $\frac{AX}{AB} =$

(A) $\frac{AX}{AY}$ (B) $\frac{AX}{XB}$ (C) $\frac{AY}{AC}$ (D) $\frac{AC}{AY}$



- 12) In trapezium ABCD , $AB \parallel CD$ and diagonals intersect at O then the value of $\frac{OD}{OC} =$


(A) $\frac{OB}{OA}$ (B) $\frac{AB}{CD}$ (C) $\frac{OC}{OD}$ (D) $\frac{AC}{BD}$



- 13) Sides of a triangle are of length 2 cm, 3 cm and 4 cm respectively.
 the set of numbers which are similar to the above triangle is

(A) 4, 5, 6 (B) 5, 6, 7 (C) 12, 13, 14 (D) 6, 9, 12

- 14) In fig $\angle ABC = \angle AQP = 90^\circ$, then $\frac{AQ}{AB}$



(A) $\frac{BC}{PQ}$ (B) $\frac{AC}{PQ}$ (C) $\frac{QP}{BC}$ (D) $\frac{AP}{AB}$

15) Corresponding sides in equiangular triangle are

- (A) ಸಮನಾಗಿರುತ್ತವೆ (B) ಸಮಾಂತರದಲ್ಲಿರುತ್ತವೆ
(C) ಸಮಾನುಪಾತದಲ್ಲಿರುತ್ತವೆ (D) ಅಸಮನಾಗಿರುತ್ತವೆ

16) Sides of two similar triangles are in the ratio 2:3. Areas of these triangles are in the ratio

- (A) 9 : 4 (B) 4 : 9 (C) 2 : 3 (D) 3 : 2

17) Areas of two similar triangles are in the ratio 25 : 49. Sides of these triangles are in the ratio

- (A) 4 : 6 (B) 5 : 7 (C) 6 : 7 (D) 7 : 8

18) In triangle PQR , $\angle PQR = 90^\circ$. then the correct relation is

- (A) $PR^2 = PQ^2 - QR^2$ (B) $PQ^2 = QR^2 - PR^2$
(C) $PR^2 = PQ^2 + QR^2$ (D) $QR^2 = PQ^2 - PR^2$

19) Pythagorean triplet among these


- (A) 3, 4, 5 (B) 1, 2, 3 (C) 2, 3, 4 (D) 9, 10, 14

20) Among these which one forms the sides a right angle triangle

- (A) 3, 6, 9 (B) 15, 8, 17 (C) 5, 12, 17 (D) 8, 5, 17

21) In the fig, $\angle ABC = 90^\circ$, $\angle ADCB = 90^\circ$, AD = 8 cm, and CD = 2 cm, then find length of BD

- (A) 4 cm (B) 8 cm (C) 16 cm (D) 10 cm




22. In the given figure, $\angle ABC \sim \angle AYX$, then the ratio of the corresponding sides is :

A) $\frac{AX}{AC} = \frac{AB}{AY} = \frac{CB}{XY}$

B) $\frac{AB}{AY} = \frac{BC}{XY} = \frac{AX}{AC}$

C) $\frac{AB}{AX} = \frac{AC}{AY} = \frac{BC}{XY}$

D) $\frac{AX}{AC} = \frac{AY}{AB} = \frac{XY}{CB}$




23. In the figure, $DE \parallel BC$, $AD : AB = 1 : 2$, $BC = 6$ cm, then DE is

(A) 1 cm

(B) 2 cm

(C) 3 cm

(D) 4 cm





24. In the given figure $\triangle ABC \sim \triangle PQC$. The ratio of their corresponding sides is

A) $\frac{AB}{PQ} = \frac{BC}{PC} = \frac{AC}{QC}$

B) $\frac{AB}{PC} = \frac{BC}{PQ} = \frac{AC}{QC}$

C) $\frac{BC}{PQ} = \frac{AB}{QC} = \frac{AC}{PC}$

D) $\frac{AB}{PQ} = \frac{BC}{QC} = \frac{AC}{PC}$



25. In the figure, $\angle ABC = \angle AQP = 90^\circ$, then $\frac{AQ}{AB}$

- A) $\frac{BC}{PQ}$ (b) $\frac{AC}{PQ}$ (c) $\frac{QP}{BC}$ (d) $\frac{AP}{AB}$


26. Sides of a triangle are of length 2 cm, 3 cm and 4 cm respectively. Which of the sets of numbers are the sides of a triangle, similar to the above triangle ?

- (A) 4, 5, 6 (B) 5, 6, 7 (C) 12, 13, 14 (D) 6, 9, 12

27. Select the set of numbers from the following which can form similar triangles

- (A) 9, 12, 18 and 3, 4, 6 (B) 3, 4, 6 and 9, 10, 12
 (C) 8, 6, 12 and 2, 6, 3 (D) 3, 4, 5 and 2, 4, 10

28. In parallelogram ABCD, P is a point on BC. In $\triangle DCP$ and $\triangle BLP$, DP : PL is equal to :



- (A) DC : BL (B) DC : BP (C) PC : BL (D) PC : PL

29. $\triangle ABC$ has sides of length 5 cm, 6 cm and 7 cm. The perimeter of $\triangle DEF$ is 360 cm. If $\triangle ABC \sim \triangle DEF$ then the ratio of the perimeters of $\triangle ABC$ and $\triangle DEF$ is :

- (A) 1 : 2 (B) 2 : 1 (C) 1 : 20 (D) 20 : 1

30. $\triangle ABC \sim \triangle DEF$, $\angle A = \angle D$ and $\angle B = \angle E$, then $\frac{\text{Area of triangle } ABC}{\text{Area of triangle } DEF}$ is equal to :

- (A) $\frac{AC^2}{DF^2}$ (B) $\frac{AB^2}{DF^2}$ (C) $\frac{AC^2}{EF^2}$ (D) $\frac{BC^2}{DE^2}$

31. $\triangle ABC \sim \triangle DEF$, the area of $\triangle ABC$ is 45 cm^2 and the area of $\triangle DEF$ is 20 cm^2 one side of $\triangle ABC$ is 3.6 cm , then the length of corresponding side of $\triangle DEF$ is:

- (A) 3.4 cm (B) 2.4 cm (C) 1.4 cm (D) 4.4 cm

32. If the perimeters of two similar triangles are in the ratio of $4 : 1$, then the ratio between their areas will be :

- (A) $16 : 1$ (B) $4 : 1$ (C) $2 : 1$ (D) $2 : 1$

33. Which of the following is a correct statement ?

- (A) All the rectangles are similar
(B) All the right angled triangles are similar
(C) All the rhombus are similar
(D) All the equilateral triangles are similar

34. The area of $\triangle ABC = 144 \text{ sq. cm}$ and area of $\triangle PQR = 25 \text{ sq. cm}$. Altitude of $\triangle ABC = 6 \text{ cm}$. If $\triangle ABC \sim \triangle PQR$, then the corresponding altitude of $\triangle PQR$ is

- (A) 2.5 cm (B) 5 cm (C) 12 cm (D) 6 cm

35. Two similar triangles have areas 120 sq. cm and 480 sq. cm respectively. Then the ratio of any pair of corresponding sides is :

- (A) $1 : 4$ (B) $1 : 2$ (C) $4 : 1$ (D) $2 : 3$

36. In $\triangle ABC$, D , E and F are mid points of AB , BC and CA respectively. If area of $\triangle ABC = 60 \text{ sq. cm}$, then area of $\triangle DEF$ is :

- (A) 15 sq. cm (B) 30 sq. cm
(C) 45 sq. cm (D) 60 sq. cm

37. In $\triangle PQR$, $\angle PQR = 90^\circ$. The correct relation with respect to
 (A) $PR^2 = PQ^2 - QR^2$ (B) $PQ^2 = QR^2 - PR^2$
 (C) $PR^2 = PQ^2 + QR^2$ (D) $QR^2 = PQ^2 - PR^2$
38. "If the square of one side of a triangle is equal to the sum of the squares on the other two sides, then those two sides contain a right angle." This statement refers
 (A) Pythagoras theorem (B) Thales theorem
 (C) Converse of Thales theorem
 (D) Converse of Pythagoras theorem.
39. The length of a diagonal of a square of side 5 cm is :
 (A) $5\sqrt{2}$ cm (B) $2\sqrt{5}$ cm (C) 10 cm (D) $10\sqrt{2}$ cm
40. In a rhombus ABCD, diagonals intersect at O. The sum of $AC^2 + BD^2$ are :
 (A) $4 AB^2$ (B) $4 AC^2$ (C) $4 BD^2$ (D) $4 AO^2$
41. A man goes 15 m due west and then 8 m due north. Calculate the distance from the starting point.
 (A) 17m (B) 15m (C) 12m (D) 23m
42. A ladder 17 m long reaches a window of a building 15 m above the ground. The distance of the foot of the ladder from the building is
 (A) 32m (B) 2m (C) 8m (D) 13m

ANSWERS:

1	C	11	C	21	A	31	B	41	A
2	C	12	A	22	D	32	A	42	C
3	C	13	D	23	C	33	D		
4	B	14	C	24	D	34	A		
5	D	15	C	25	C	35	B		
6	A	16	B	26	D	36	A		
7	A	17	B	27	A	37	C		
8	D	18	C	28	A	38	D		
9	C	19	A	29	C	39	A		
10	C	20	B	30	A	40	A		

Points to remember

$$a_1x + b_1y + c_1 = 0, \quad a_2x + b_2y + c_2 = 0$$

Where $a_1, a_2, b_1, b_2, c_1, c_2$ are real numbers

❖ Conditions for solvability (or consistency):

Condition	Solution	Graphical representatio	Consistency / Inconsistency
$\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$	Unique solution		

A) 12 B) 8 C) 10 D) 4

A) $\frac{a_1}{a_2} \neq \frac{b_1}{b_2} = \frac{c_1}{c_2}$ B) $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$ C) $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$ D) $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$

A) $2x - 3y + 6 = 0, 2x + 3y + 6 = 0$

B) $3x - 4y - 6 = 0, 3x - 4y + 6 = 0$

C) $x - y + 10 = 0, x - y + 10 = 0$

D) $5x - 10y + 20 = 0, 5x - 20y + 30 = 0$

6) The inconsistent pair of equations among the following

A) $x - y + 3 = 0, x - y + 6 = 0$

B) $2x - y + 20 = 0, x - 2y + 10 = 0$

C) $3x - 4y + 12 = 0, x - y + 10 = 0$

D) $5x - 10y + 20 = 0, x - 2y + 4 = 0$

A) unique solution B) infinitely many solutions

C) no solution D) have two solutions

A) Consistent pair B) Dependent pair

C) Inconsistent pair D) Straight pair

A) unique solution

B) infinitely many solutions

C) no solution

D) have two solutions

10) The value of x and y for two equations $x + y = 5$ and $2x - y = 4$ is

A) (2, 3)

B) (3, 2)

C) (1, 4)

D) (4, 1)

A) 6 B) 3 C) 4 D) 2

A) 2. B) 4. C) 6. D) 8.

C) coincident

D) inconsistent

A) 1 B) 2 C) 3 D) 4

A) $2x - 3y = 9$. B) $4x + 6y = 3$. C) $6x + 9y = 10$. D) $2x + 3y = 9$.

ANSWERS:

1	C	7	C	13	C
2	D	8	A	14	B
3	A	9	C	15	A
4	B	10	B	16	B
5	C	11	D		
6	A	12	D		

Circles

Points to remember

- ❖ The point where a tangent touches the circles is called the point of contact.
- ❖ In any circle, the radius drawn at the point of contact is perpendicular to the tangent.
- ❖ Only one tangent can be drawn to a circle at any point on it.
- ❖ Tangents drawn at the ends of a diameter are parallel to each other.
- ❖ Length of tangents from an external point to a circle are equal.

Multiple choice questions

A) Chord

B) Secant

C) ~~Diameter~~ Radius

A) Tangent

B) Secant

C) Radius

D) arc of a circle

3) The maximum number of parallel tangents that can be drawn to a circle is

A) 1

B) 2

C) 3

D) Infinitely many

4) The angle subtended between tangent and radius of a circle is

A) 30°

B) 60°

C) 90°

D) 180°

5) The maximum number of tangents that can be drawn to a circle from an external point is

A) 1

B) 2

C) 3


D) 4

6) The number of tangents to a circle passing through a point lying on the circle is

- A) 1 B) 2 C) 3 D) Infinitely many

7) Two concentric circles are of radii 5cm & 3cm with center 'O' and AB is the chord of larger circle which touches the smaller circle at P then the length of the chord AB =

- A) 24cm B) 12cm C) 18cm D) 8 cm



8) The length of a tangent drawn to a circle of radius 6cm from an external point which is 10cm away from the centre is

- A) 8cm B) 16cm C) 10cm D) 5cm

9) If the angle between two tangents of a circle with is 70° then angle between their radii is

- A) 110° B) 70° C) 60° D) 100°

10) If the angle between two radii of a circle is 90° , the angle between tangents drawn at the ends of the radii is

- A) 0° B) 180° C) 60° D) 90°


11) The tangents drawn at the ends of a diameter of a circle are

- A) parallel to each other B) Perpendicular to each other
C) intersects to each other D) Coincides to each other

12) A quadrilateral ABCD is drawn to circumscribe a circle.

AB+CD=

- A) AC+CD B) AD+BC C) AB+BC D) AB+AD




13) PQ is a tangent to a circle with center 'O' touches the circle at point P then $\angle OPQ =$

- A) 30° B) 60° C) 90° D) 180°

14) In the figure BC, CE and EF are the tangents drawn to a circle. If BC = 5 cm, EF = 3cm, then the length of CE =

- A) 3cm b) 5cm C) 8cm D) 2 cm




15) "P" is the external point at a distance of 5cm from the center 'O' of the circle with radius OQ = 3cm and PQ is the tangent touches the circle at Q then the length of PQ =

- A) 4cm B) 7cm C) 8cm D) 2cm

16) If TP and TQ are the two tangents to a circle with center 'O' so that $\angle POQ = 110^\circ$ then $\angle PTQ =$

- A) 110° B) 70° C) 60° D) 100°



17) PQ & PR are tangents to a circle with center 'O' if $\angle QPR = 50^\circ$ then $\angle QOR =$


- A) 130° B) 50° C) 65° D) 90°

18) The tangents drawn at the ends of a diameter of a circle are

- A) parallel to each other B) Perpendicular to each other
 C) intersects to each other D) Coincides to each other

19) In the Figure PB is the tangent to a circle with center 'A' if $\angle ABP = 40^\circ$ then $\angle PAB =$

- A) 90° B) 50° C) 40° D) 60°



20) A circle with radius 3cm is circumscribe in a square ABCD. Then, the perimeter of a square is A) 24cm B) 6cm C) 12cm D) 36 cm

ANSWERS:

1	A	7	D	13	C	19	B
2	A	8	A	14	C	20	C
3	B	9	A	15	A		
4	C	10	D	16	B		
5	B	11	B	17	A		
6	A	12	B	18	A		

Constructions

A) 4cm 5cm C) 4cm 5cm

A) 4cm ಮತ್ತು 7cm B) 4cm ಮತ್ತು 6cm

C) 4.6cm ಮತ್ತು 5.4cm D) 4.2cm ಮತ್ತು 5.8cm

3) Construct a triangle of with sides 6cm , 9cm and 7.5cm respectively. Then construct another triangle similar to the given triangle such that each of its sides are $\frac{2}{3}$ of the corresponding sides of given triangle . The lengths of the new triangle respectively are

A) 4cm , 6cm , 8cm B) 4cm , 6cm , 10cm

C) 4cm , 7cm , 6cm D) 4cm , 6cm , 5cm

ASWERS:

1	2	3
C	B	D

Coordinate geometry

Points to remember

1. Distance between P(x₁, y₁) & Q(x₂, y₂) is given by

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

2. Distance between origin and a point P(x, y) is given by

$$d = \sqrt{x^2 + y^2}$$

$$(x, y) = \left[\frac{m_1x_2 + m_2x_1}{m_1 + m_2}, \frac{m_1y_2 + m_2y_1}{m_1 + m_2} \right]$$

$$= \left[\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right]$$

5. Area of a triangle with vertices (x₁, y₁), (x₂, y₂) & (x₃, y₃) is

$$\Delta = \frac{1}{2} \{x_1 (y_2 - y_3) + x_2 (y_3 - y_1) + x_3 (y_1 - y_2)\}$$

Multiple questions

- 1) The distance between (X₁, Y₁) & (X₂, Y₂) is

A) $\sqrt{(X_2 - X_1)^2 + (Y_2 - Y_1)^2}$ B) $(X_2 - X_1)^2 + (Y_2 - Y_1)^2$
C) $\sqrt{(X_2 - X_1) + (Y_2 - Y_1)}$ D) $(X_2 - X_1) + (Y_2 - Y_1)$

- 2) The distance of a point (X, Y) from the origin is

A) $X^2 + Y^2$ B) $\sqrt{X^2 + Y^2}$ C) $X^2 - Y^2$ D) $\sqrt{X^2 - Y^2}$

$$\left(\frac{X_1+Y_1}{2}, \frac{X_2+Y_2}{2} \right)$$

$$B) \left(\frac{X_1-Y_1}{2}, \frac{X_2-Y_2}{2} \right)$$

$$C) \left(\frac{x_2+x_1}{2}, \frac{y_2+y_1}{2} \right)$$

$$D) \left(\frac{x_2-x_1}{2}, \frac{y_2-y_1}{2} \right)$$

4) The distance between the coordinates (3,4) from its origin is

- A) 6 units, B) 5 units C) 10 units D) 8 units

5) The distance between the coordinates (8,3) and (5,7) is

- A) 5 units B) 11 units C) $2\sqrt{2}$ units D) 4 units

- A) (5, 3) B) (-3, 7) C) (8,1) D) (16,2)

7) The coordinates of the end points of a diameter of a circle are (6,2) & (6,4). Then the coordinates of the center of the circle are

- A) (4,5) B) (6,3) C) (5,4) D) (10, 8)

- A) ± 7 B) ± 4 C) 0 D) ± 3

9) The coordinates of the point which divides the join of (3,2) & (0,5) in the ratio 2:1 are

- A) (1, 4) B) (4,1) C) (3, 7) D) (7,3)

(X_3, Y_3) is

- A) 1 sq.units B) 0 sq.units C) 100 sq.units D) -1 sq.units

- A) 4 B) 5 C) -1 D) $3\sqrt{3}$

- A) 3:4 B) 3:2 C) 2:3 D) 4:3

- 15) The area of a triangle with vertices (1,-1) (-4,6) & (-3,-5) is
 A) 24 sq.units B) 40 sq.units C) 48 sq.units D) 32 sq.units
- 16) The distance of the point P(2, 7) from the x – axis is _____
 A) 2 units B) 7 units C) 9 units D) 11 units
- 17) The distance of the point Q(6, 2) from the Y – axis is _____
 A) 2 units B) 4 units C) 6 units D) 8 units
- 18) Co-ordinates of origin are.
 A) (1, 1) B) (1, 0) C) (0, 1) D) (0, 0)

A (x_1, y_1) , B (x_2, y_2) and C (x_3, y_3) is _____

- A) . $\frac{1}{2} [x_1 (y_2 + y_3) + x_2 (y_3 + y_1) + x_3 (y_1 + y_2)]$
- B) . $\frac{1}{2} [x_1 (y_2 - y_3) + x_2 (y_3 - y_1) + x_3 (y_1 - y_2)]$
- C) . $\frac{1}{2} [x_1 (y_2 - y_3) - x_2 (y_3 - y_1) - x_3 (y_1 - y_2)]$
- D) . $\frac{1}{2} [x_1 (y_2 + y_3) - x_2 (y_3 + y_1) - x_3 (y_1 + y_2)]$

A) . $\left[\frac{m_1 x_2 + m_2 x_1}{m_1 + m_2}, \frac{m_1 y_2 + m_2 y_1}{m_1 + m_2} \right]$

B) . $\left[\frac{m_1 x_2 - m_2 x_1}{m_1 + m_2}, \frac{m_1 y_2 - m_2 y_1}{m_1 + m_2} \right]$

C) . $\left[\frac{m_1x_2+m_2x_1}{m_1-m_2} , \frac{m_1y_2+m_2y_1}{m_1-m_2} \right]$ D). $\left[\frac{m_1x_2-m_2x_1}{m_1-m_2} , \frac{m_1y_2-m_2y_1}{m_1-m_2} \right]$

21) The area of the triangle whose vertices are A **(3, 3)**, B**(3, 5)** and C **(2, 4)** is

- A) 0 units B) 1 units C) 2 units D) 3 units

22) The perpendicular distance of point A (6, 9) from the x-axis and y-axis is

- A) 6 units, 9units B) 9 units , 6units
 C) 6 units ,15 units D) 15 units, 9units

23) The coordinates of the point on the x- axis will be in the form _____

- A) (0, y) B) (x, 0) C) (0, 0) D) (x, y)

24) The coordinates of the point on the y- axis will be in the form _____

- A) (0, y) B) (x, 0) C) (0, 0) D) (x, y)

25) Y- coordinate on x-axis is _____

- A) 0 B) 1 C) 2 D) 3

26) X-coordinate on y-axis is _____

- A) 3 B) 2 C) 1 D) 0

27) The point among the following which lie on x-axis is

- A) (2, 0) B) (0, 2) C) (2, 3) D) (0, -2)

28) The point among the following which lie on y-axis is

- A) (3, 0) B) (0, -4) C) (-2, 0) D) (4, 6)

ANSWERS:

1	A	7	B	13	D	19	B	25	A
2	B	8	B	14	A	20	A	26	D
3	C	9	A	15	A	21	B	27	A
4	B	10	B	16	B	22	B	28	B
5	A	11	C	17	C	23	B		
6	C	12	C	18	D	24	A		

Quadratic equations

Points to remember

1. Standard form of a quadratic equation is $ax^2 + bx + c = 0, a \neq 0$

2. Quadratic formula to find roots of $ax^2 + bx + c = 0$ are

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

3. Nature of roots of a quadratic equation

- i. The discriminant to find nature of roots of a quadratic equation $ax^2 + bx + c = 0$ is given by $\Delta = b^2 - 4ac$
- ii. If $\Delta = b^2 - 4ac = 0$ then roots are real and equal.
- iii. If $\Delta = b^2 - 4ac > 0$ (+ve) then roots are real and distinct.
- iv. If $\Delta = b^2 - 4ac < 0$ (-ve) No real roots.

1. Which of the following is not a quadratic equation

- A) $x^2 + 3x - 5 = 0$ (B) $x^2 + x^3 + 2 = 0$ (C) $3 + x + x^2 = 0$ (D) $x^2 - 9 = 0$

2. The quadratic equation has degree

- (A) 0 (B) 1 (C) 2 (D) 3

3. The standard form of a quadratic equation is

- A) $ax^2 + bx + c = 0$ B) $ax + b = c$ C) $ax^3 + bx^2 + c = 0$ D) $ax - b = 0$

4. Pure quadratic equation among the following is

- A) $x^2 + 2x + 2 = 0$ B) $x^2 + 5x + 6 = 0$

- C) $x^2 + 9 = x$ D) $x^2 - 9 = 0$

5. Sridhara's quadratic formula for the quadratic equation $ax^2 + bx + c = 0$ is

A) $-b \pm \sqrt{b^2 - 4ac}$

B) $\frac{-b}{2a} \pm \sqrt{b^2 - 4ac}$

C) $\frac{-b \pm \sqrt{b^2 - 4ac}}{4a}$

D) $\frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

6. Formula to find the discriminant of the quadratic equation $ax^2 + bx + c = 0$ is

A) $b^2 - 4ac$

B) $\frac{b}{2a}$

C) $\frac{-b}{2a}$

D) $\frac{-b}{4ac}$

7. The maximum number of roots of a quadratic equation is

A) one

B) two

C) three

(D) depends on the given equation

8. Sum of a number and its reciprocal is $\frac{17}{4}$, the number is

A) 4

B) 3

C) 5

D) 2

9. The values of X in given Quadratic equation $X^2 + 1 = 101$ are

A) ± 1 B) ± 10 C) ± 11 D) $\pm \sqrt{10}$

10. The roots of the Quadratic equation $x^2 = 49$ are

A) 7 and -7

B) 24 and 5

C) 8 and -8

D) 7 and 0

11. If the roots of the equation $ax^2 + bx + c = 0$ are equal, then the value of C is

A) $b^2 - 4a$ B) $b^2 \times 4a$ C) $\frac{b^2}{4a}$ D) $\frac{4a}{b^2}$

12. The discriminant of the given quadratic equation $2x^2 - 4x + 3 = 0$ is

A) -8

B) 8

C) 0

D) 1

13. If the roots of the quadratic equation $x^2 + mx + 4 = 0$ are equal, then the value of m is

A) 2

B) 4

C) 6

D) 5

14. "The product of two consecutive positive integers is 30" this statement can be expressed as

A) $x(x+2) = 30$ B) $x(x-2) = 30$ C) $(x-3)x = 30$ D) $x(x+1) = 30$

15. Which of the following is an example for quadratic equation

- A) $x(x+3) + 5 = x^2$ B) $x(x-3)=5$ C) $2x^2 + 2x = 2(x^2 - 5)$ D) $(x+1)x=x(x-3)$

16. The nature of roots of the equation $x^2 + 4x + 4 = 0$ is

- A) Real and Equal B) Real and distinct C) No real D) Different roots

17. " The sum of squares of two consecutive odd positive integers is 34" this can expressed as

A) $x^2 + (x+1)^2 = 34$ B) $x^2 + (x+3)^2 = 34$

C) $(x+1)^2 + (x+2)^2 = 34$ D) $x^2 + (x+2)^2 = 34$

18. If -5 is a root of the quadratic equation $2x^2 + px - 15 = 0$, then the value of p is

- (A) 3 (B) 5 (C) 7 (D) 1

19. The sum of the reciprocals of Rehman's ages 3 years ago and 5 years from now is $\frac{1}{3}$. The present age of Rehman is:

- (A) 7 years (B) 10 years (C) 5 years (D) 6 years

20. What number should be added to $x^2 + 6x$ to make it a perfect square?

- A) 36 (B) 18 (C) 9 (D) 72


ANSWERS:

1	B	6	A	11	C	16	A
2	C	7	B	12	A	17	D
3	A	8	A	13	B	18	C
4	D	9	B	14	D	19	A
5	D	10	A	15	B	20	A

UNIT : TRIGONOMETRY

Points to remember

TRIGONOMETRIC RATIOS :



$\sin A$	$\cos A$	$\tan A$	$\operatorname{cosec} A$	$\sec A$	$\cot A$
$\frac{\text{Opposite}}{\text{Hypotenuse}}$	$\frac{\text{Adjacent}}{\text{Hypotenuse}}$	$\frac{\text{Opposite}}{\text{Adjacent}}$	$\frac{\text{Hypotenuse}}{\text{Opposite}}$	$\frac{\text{Hypotenuse}}{\text{Adjacent}}$	$\frac{\text{Adjacent}}{\text{Opposite}}$

TRIGONOMETRICAL RATIOS TABLE FOR STANDARD ANGLES

$\angle A$	0°	30°	45°	60°	90°
$\sin A$	0	$\frac{1}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{\sqrt{3}}{2}$	1
$\cos A$	1	$\frac{\sqrt{3}}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{1}{2}$	0
$\tan A$	0	$\frac{1}{\sqrt{3}}$	1	$\sqrt{3}$	N.D
$\operatorname{cosec} A$	N.D	2	$\sqrt{2}$	$\frac{2}{\sqrt{3}}$	1
$\sec A$	1	$\frac{2}{\sqrt{3}}$	$\sqrt{2}$	2	N.D
$\cot A$	N.D	$\sqrt{3}$	1	$\frac{1}{\sqrt{3}}$	0

TRIGONOMETRIC RATIOS OF COMPLEMENTARY ANGLES

$\sin A$	$\cos A$	$\tan A$	$\operatorname{cosec} A$	$\sec A$	$\cot A$
$\cos(90-A)$	$\sin(90-A)$	$\cot(90-A)$	$\sec(90-A)$	$\operatorname{cosec}(90-A)$	$\tan(90-A)$

TRIGONOMETRIC IDENTITY

- $\sin^2 A + \cos^2 A = 1$
- $\tan^2 A + 1 = \sec^2 A$
- $1 + \cot^2 A = \operatorname{cosec}^2 A$

INVERSE OF TRIGONOMETRIC VALUES

$\frac{1}{\sin A}$	$\operatorname{CoSec} A$
$\frac{1}{\cos A}$	$\sec A$
$\frac{1}{\tan A}$	$\cot A$
$\frac{1}{\operatorname{cosec} A}$	$\sin A$
$\frac{1}{\sec A}$	$\cos A$
$\frac{1}{\cot A}$	$\tan A$

SOME IMPORTANT MULTIPLE CHOICE QUESTIONS

- The value of $\tan 45^\circ$ is _____
A) $\sqrt{3}$ B) 0 C) 1 D) $\frac{1}{\sqrt{3}}$
- If $2\sin 2\theta = \sqrt{3}$ then the value of ' θ ' is _____
A) 90° B) 60° C) 30° D) 45°
- $\sin(90-\theta)$ is equal to _____
A) $\cos \theta$ B) $\tan \theta$ C) $\sec \theta$ D) $\cot \theta$
- $\tan \theta - \cot(90-\theta)$ is equal to _____
A) 0 B) 1 C) 2 D) 3
- If $\sin \theta = \frac{x}{y}$ then the value of $\cos \theta$ is _____
A) $\frac{y}{\sqrt{y^2-x^2}}$ B) $\frac{y}{x}$ C) $\frac{x}{\sqrt{y^2-x^2}}$ D) $\frac{\sqrt{y^2-x^2}}{y}$
- If $13\sin \theta = 12$ then $\operatorname{cosec} \theta =$ _____
A) $\frac{5}{12}$ B) $\frac{13}{5}$ C) $\frac{12}{13}$ D) $\frac{13}{12}$
- The value of $\sin 30^\circ + \cos 60^\circ$ is _____
A) 1 B) $\frac{3}{2}$ C) $\frac{1}{4}$ D) $\frac{2}{3}$
- The value of $\sin 60^\circ \cdot \cos 30^\circ + \sin 30^\circ \cdot \cos 60^\circ$ is _____
A) 4 B) 3 C) 2 D) 1
- If $\tan A = \frac{4}{3}$ then $\cot A =$ _____
A) $\frac{4}{3}$ B) $\frac{3}{4}$ C) $\frac{4}{5}$ D) $\frac{5}{3}$
- The value of $\tan 45^\circ + \cot 45^\circ - \sin 90^\circ$ is _____
A) 3 B) 2 C) 1 D) 0
- The value of $\frac{\sin 18^\circ}{\cos 72^\circ}$ is _____
A) 0 B) 1 C) 2 D) 3
- The value of $\frac{\tan 26^\circ}{\cot 64^\circ}$ is _____
A) 0 B) 1 C) 2 D) 3

13) The value of $\frac{\tan 65^\circ}{\cot 25^\circ}$ is _____


- A) 0 B) 1 C) 2 D) 3

14) The value of $\cos 48^\circ - \sin 42^\circ$ is _____

- A) 0 B) 1 C) $\frac{1}{2}$ D) -1

15) In the figure the value of $\sin \theta$ is _____

- A) $\frac{5}{13}$ B) $\frac{12}{13}$
C) $\frac{13}{12}$ D) $\frac{13}{5}$



16) $\sin^2 \theta + \cos^2 \theta =$ _____

- A) -1 B) 1 C) 0 D) 2

17) In a $\triangle ABC$, $\hat{B} = 90^\circ$, $AB = 24$ cm and $BC = 7$ cm then $\tan C =$ _____

- A) $\frac{12}{7}$ B) $\frac{24}{7}$ C) $\frac{7}{12}$ D) $\frac{7}{24}$

18) $1 - \cos^2 A =$ _____

- A) $\sin^2 A$ B) $\tan^2 A$ C) $1 - \sin^2 A$ D) $\sec^2 A$

19) $\sec^2 \theta - 1 =$ _____

- A) $\tan^2 \theta$ B) $\tan^2 \theta + 1$ C) $\cot^2 \theta - 1$ D) $\cos^2 \theta$

20) The value of $9\sec^2 A - 9\tan^2 A$ is _____

- A) 1 B) 9 C) 8 D) 0

21) $\tan \theta \cdot \cot \theta =$ _____

- A) $\cot \theta$ B) $\tan \theta$ C) $\tan^2 \theta$ D) 1

22) If $\sqrt{3} \tan \theta = 1$ then the value of ' θ ' is _____

- A) 0° B) 30° C) 45° D) 60°

- 23) The value of $\sin^2 30^\circ - \cos^2 30^\circ$ is _____
 A) $-\frac{1}{2}$ B) $\frac{\sqrt{3}}{2}$ C) $\frac{3}{2}$ D) $-\frac{2}{3}$
- 24) The value of $\frac{1 - \tan^2 45^\circ}{1 + \tan^2 45^\circ}$ is _____
 A) $\tan 90^\circ$ B) 1 C) $\sin 45^\circ$ D) 0
- 25) If we express $\operatorname{cosec} \theta$ in the form of $\cot \theta$, then $\operatorname{cosec} \theta =$ _____
 A) $\cot \theta$ B) $\cot^2 \theta - 1$ C) $\sqrt{1 + \cot^2 \theta}$ D) $1 + \cot^2 \theta$
- 26) If $\sin A = \frac{1}{2}$ and $\cos B = \frac{1}{2}$ then $A + B =$ _____
 A) 0° B) 30° C) 60° D) 90°
- 27) In a $\triangle ABC$ if $\hat{C} = 90^\circ$, then $\sin(A + B) =$ _____
 A) 0 B) $\frac{1}{2}$ C) $\frac{1}{\sqrt{2}}$ D) 1
- 28) If $\sin \theta = \frac{1}{2}$ and $\cos \theta = \frac{\sqrt{3}}{2}$, then $\tan \theta =$ _____
 A) $\sqrt{3}$ B) $\frac{1}{2}$ C) $\frac{\sqrt{2}}{3}$ D) $\frac{1}{\sqrt{3}}$
- 29) In a $\triangle ABC$ $\hat{B} = 90^\circ$ $\tan A = \frac{1}{\sqrt{3}}$, then $\sin A =$ _____
 A) 0 B) $\frac{1}{2}$ C) $\frac{\sqrt{3}}{2}$ D) $\frac{1}{\sqrt{2}}$
- 30) In $\sin A$, $0 \leq A \leq 90^\circ$ then the least value of $\sin A$ is _____
 A) -1 B) 0 C) $\frac{1}{\sqrt{2}}$ D) $\frac{1}{2}$
- 31) If $\sin 2A = 2 \sin A$ then the value of A is _____
 A) 30° B) 45° C) 0° D) 90°
- 32) In a $\triangle ABC$, $\hat{B} = 90^\circ$ and $\hat{C} = 30^\circ$ then $AB : AC$ is _____
 A) 1 : 2 B) 2 : 1 C) $\sqrt{3} : 2$ D) $2 : \sqrt{3}$

- 33) If $\tan A = \cot B$, then $A + B =$ _____
 A) 90° B) 60° C) 30° D) 0°
- 34) The value of $\tan 1^\circ \cdot \tan 2^\circ \cdot \tan 3^\circ \dots \dots \dots \tan 89^\circ$ is _____
 A) 0 B) 1 C) 2 D) $\frac{1}{2}$
- 35) $\frac{2\tan 30^\circ}{1+\tan^2 30^\circ} =$ _____
 A) $\sin 60^\circ$ B) $\cos 60^\circ$ C) $\tan 60^\circ$ D) $\sin 30^\circ$
- 36) $\frac{1+\tan^2 A}{1+\cot^2 A} =$ _____
 A) $\sec^2 A$ B) -1 C) $\cot^2 A$ D) $\tan^2 A$
- 37) If $\sin A - \cos A = 0$ then the value of $\sin^4 A + \cos^4 A$ is _____
 A) 2 B) 1 C) $\frac{3}{4}$ D) $\frac{1}{2}$
- 38) if $\sin \theta + \sin^2 \theta = 1$ then the value of $\cos^2 \theta + \cos^4 \theta$ is _____
 A) -1 B) 1 C) 0 D) 2

ANSWERS:

1	C	11	B	21	D	31	C
2	C	12	B	22	B	32	A
3	A	13	B	23	A	33	A
4	A	14	A	24	D	34	B
5	D	15	B	25	C	35	A
6	D	16	B	26	D	36	D
7	A	17	B	27	D	37	D
8	D	18	A	28	D	38	B
9	B	19	A	29	B		
10	C	20	B	30	B		

12. Applications of trigonometry

Multiple choice questions

- 1) The shadow of a tower is equal to its height at 10-45 a.m. The sun's altitude is
(A) 30° (B) 45° (C) 60° (D) 90°
- 3) The angle of elevation of the top of a tower from a point 20 metres away from its base is 45° . The height of the tower is
(A) 10 m (B) 20 m (C) 30 m (D) $20\sqrt{3}$ m
- 4) If the length of the shadow of a tower increases, then the angle of elevation of the sun
(A) is also increases (B) decreases
(C) remains unaffected (D) increases then decreases
- 5) The angle of elevation of the top of a tower is 30° . If the height of the tower is doubled, then the angle of elevation of its top will
(A) also get doubled (B) will get halved
(C) will be less than 60° (D) 30°
- 6) If a pole 6m high casts a shadow $2\sqrt{3}$ m long on the ground, then the sun's elevation is
(A) 60° (B) 45° (C) 30° (D) 90°
- 7)

ANSWERS:

1	2	3	4	5	6	7
B	C	B	B	C	A	A

Statistics

Points to remember

1. Mean for groupd data

Direct method to find mean

$$\text{mean , } \bar{x} = \frac{\sum f_i x_i}{\sum f_i}$$

$$2. \text{ Mode for grouped data } = L + \left[\frac{f_1 - f_0}{2f_1 - f_0 - f_2} \right] \times h$$

$$3. \text{ Median for grouped of } = L + \left[\frac{\frac{N}{2} - cf}{f} \right] \times h$$

Multiple choise questoins

- 1) If the mean value of 'x', 6, 8, 9 and 12 is 8, then the value of 'x' is
A) 4 B) 5 C) 16 D) 10
- 2) The median of the scores 5, 8, 14, 16, 19 and 20 is
A) 14 B) 15 C) 16 D) 17
- 3) The wickets taken by a bowler in 10 cricket matches are as follows :
2, 6, 4, 5, 0, 2, 1, 3, 2, 3 then the mode of the data is
A) 0 B) 1 C) 2 D) 3
- 4) The emperical relationship between the three measures of central tendency is
A) 3 median = mode + 2 mean B) 2 median = mode + 3 mean
C) median = mode + mean D) median = mode – mean
- 5) Class mark for the class 10 – 25 is

- A) 10.5 B)12.5 C)15.5 D) 17.5

6) Size of the class interval 40-50 is

- ಗಾತ್ರ A) 10 B) 15 C) 20 D) 25

7) Modal class for the given distribution is

CI	1-3	3-5	5-7	7-9	9-11
F	7	8	2	2	1

- A) 1-3 B) 3-5 C) 5-7 D) 9-11

8) The frequency(f_0) of class preceding the modal class for the given distribution is

CI	5-15	15-25	25-35	35-45	45-55	55-65
f	6	11	21	23	14	5

- A) 6 B) 11 C) 21 D) 23

9) The frequency (f_2) of class succeeding the modal class for the given distribution is

CI	1-3	3-5	5-7	7-9	9-11
F	7	8	2	2	1

- A) 7 B) 8 C) 2 D) 1

10) The middle most score in an orderly arranged data is

- A) Mean B) Median C) Mode D) Range

11) The mean of first five odd natural positive integers is

- A) 5 B) 7 C) 9 D) 25

12) The sum of the values of all the observations divided by the total number of observations is

- A) Range B)Mean C) Median D) Mode

13) Cumulative frequency is useful in determining the

- (A) mean (B) median (C) mode (D) range

14) Which of the following is not a measure of central tendency
 (A) mean (B) median (C) mode (D) range

15) Classmark of the class 10 – 20 is
 A) 10 B) 20 C) 15 D) 30

16) Size of the class interval 25-35 is
 A) 25 B) 10 C) 35 D) 15

17) Mode for the data 12,15,14,13,12,15,18,25,16,15,20,18
 A)15 B) 18 C) 12 D) 25

18) Modal class for the distribution

C.I	0-10	10-20	20-30	30-40	40-50	50-60	60-70
f	3	5	8	12	10	6	4

A) 20 – 30 B) 30 – 40 C) 40 – 50 D) 50 – 60

19) Formula to find median is

A) $L + \left[\frac{\frac{N}{2} - cf}{f} \right] Xh$ B) $L + \left[\frac{\frac{N}{2} + cf}{f} \right] Xh$

C) $L + \left[\frac{f_1 - f_0}{2f_1 - f_0 - f_2} \right] Xh$ D) $L + \left[\frac{f_1 - f_0}{2f_1 - f_0 + f_2} \right] Xh$

20) Median class for the distribution

C.I	0-10	10-20	20-30	30-40	40-50	50-60	60-70
f	3	8	16	28	38	46	50

A) 20 – 30 B) 30 – 40 C) 40 – 50 D) 50 – 60

21) Mean for the following scores 12,14,10,13,11 is

A)10 B) 14 C) 12 D) 15

22) The formula used to find mode of the grouped data

A) $1 + \left[\frac{n/2 - cf}{f} \right] \times h$ B) $\frac{\sum f_i x_i}{\sum f_i}$ C) $1 + \left[\frac{f_1 - f_0}{2f_1 - f_0 - f_2} \right] \times h$ D) $1 + \left[\frac{f_1 - f_0}{2f_1 - f_0 - f_2} \right]$

23) If the median is 36 and mean is 18 ,then the value of the mode is

- A) 36 B) 72 C) 18 D) 648

24) If the mean of 12 numbers is 20 then their algebraic sum is

- A) 200 B) 32 C) 240 D) 180

25) If $\sum f_i = 20$, $\sum f_i x_i = 140 + 5k$ and $\bar{X} = 9$ find K

- A) 2 B) 4 C) 8 D) 6

26) The Median of 13,12,5,8,11,9 is

- A) 5 B) 6.5 C) 10 D) 9.5

27) For the given frequency distribution table answer the following question

C-I Marks scored	F No of Students
0-10	11
10-20	9
20-30	8
30-40	12
40-50	10
50-60	10

i. The modal class is

- A) 50-60 B) 30-40 C) 0-10 D) 20-30

ii. upper limit of the modal class is

- A) 20 B) 30 C) 40 D) 50

iii. Value of ' h ' (class mark of modal class) is

- A) 10 B) 20 C) 30 D) 40

iv. The number of Students who scored more than 40 marks

A) 32

B) 12

C) 20

D) 17

ANSWERS:

1	B	7	B	13	B	19	A	25	C
2	B	8	C	14	D	20	B	26	C
3	C	9	C	15	C	21	C	27(i)	B
4	A	10	B	16	B	22	C	27(ii)	C
5	D	11	A	17	A	23	B	27(iii)	A
6	A	12	B	18	B	24	C	27(iv)	C

Surface area and volume

Points to remember

List of formulae

Sl. no	Solid	C.S.A	T.S.A	Volume
1	Cube	$4a^2$	$6a^2$	a^3
2	Cuboid	$2lb + 2lh$	$2lb + 2lh + 2bh$	lbh
3	Cylinder	$2\pi rh$	$2\pi r^2 + 2\pi rh$	$\pi r^2 h$
4	Cone	πrl	$\pi r^2 + \pi rl$	$\frac{1}{3}\pi r^2 h$
5	Frustum of cone	$\pi(r_1 + r_2)l$	$\pi r_1^2 + \pi r_2^2 + \pi(r_1 + r_2)l$	$\frac{1}{3}\pi(r_1^2 + r_2^2 + r_1 r_2)h$
6	Sphere	$4\pi r^2$	$4\pi r^2$	$\frac{4}{3}\pi r^3$
7	Hemisphere	$2\pi r^2$	$3\pi r^2$	$\frac{2}{3}\pi r^3$

Important points

- Area of combination of solids is the sum of areas of visible faces.
- Volume of combination of solids is the sum of its constituent solids.
- A solid is converted from one shape in to another , their volumes remain same.

Multiple choice questions.

1) If the area of the circular base of a cylinder is 22 cm^2 and its height is 10 cm , then the volume of the cylinder is

A) 2200 cm^2

B) 2200 cm^3

C) 220 cm^3

D) 220 cm^2

- 2) The formula used to find the curved surface area of the frustrum of a cone whose radii are r_1 and r_2 and slant height 'l' is
 A) $\pi(r_1 + r_2) l$ B) $\pi(r_1 - r_2) l$ C) $\pi(r_1 \times r_2) l$ D) $\pi(r_1 \div r_2) l$
- 3) 2 cubes each of volume 27cm^3 are joined end to end . The volume of the resulting cuboid is
 A) 27cm^3 B) 54cm^3 C) 108cm^3 D) 216cm^3
- 4) The volume of a cylinder is 300m^3 then the volume of a cone having the same radius and height as that of the cylinder is
 A) 900m^3 B) 600m^3 C) 150m^3 D) 100m^3
- 5) If two solid hemisphere of same radius are joined together along with their bases . Then the surface area of their new solid is
 A) $2\pi r^2$ B) $3\pi r^2$ C) $4\pi r^2$ D) $6\pi r^2$
- 6) Volume of a sphere with radius 'r' is
 A) $\frac{3}{4}\pi r^3$ B) $\frac{3}{2}\pi r^3$ C) $\frac{2}{3}\pi r^3$ D) $\frac{4}{3}\pi r^3$
- 7) A cylinder and a cone are of same base , radius and of same height . The ratio of the volume of the cylinder to that of the cone is
 A) 2:1 B) 3:1 C) 2:3 D) 3:2
- 8) The curved surface area of a right circular cylinder is 440cm^2 and its radius is 7cm, its height is
 A) 3.5 cm B) 7cm C) 10cm D) 14cm
- 9) A cylindrical pencil sharpened at one edge is the combination of
 A) Two cylinders
 B) A hemisphere and a cylinder
 C) Frustrum of a cone and a cylinder
 D) a cone and a cylinder
- 10) A cone made of modelling clay whose height is 24cm and radius of base 6 cm is reshaped into sphere, then the radius of sphere is
 A) 3 cm B) 6cm C) 12cm D) 24 cm
- 11) The surface area of a sphere of radius 7 cm is
 A) 154cm^2 B) 308cm^2 C) 616cm^2 D) 770cm^2
- A) Cuboid B) Cylinder C) Sphere D) Rightcircular cone

13) A toy is in the form of a cone mounted on a hemisphere of same radius . The total surface area of the toy is

- A) $\pi r l + 2\pi r^2$ B) $\pi r l + \pi r^2$ C) $2\pi r l + \pi r^2$ D) $2\pi r l + 2\pi r^2$

14) Formula to find volume of a cylinder is

- A] $\pi r^2 h$ B] $\frac{1}{3} \pi r^2 h$
 C] $\pi r l$ D] $2\pi r h$

15) The solid which is having only one surface is

- A] Sphere B] Hemisphere C] Cylinder D] Cone

16) The ratio of areas of two spheres with the ratio of their radii 2:3 is

- A] 2:3 B] 3:2 C] 4:9 D] 16:9

17) Formula to find total surface area of a hemisphere is

- A] πr^2 B] $3\pi r^2$ C] $2\pi r^2$ D] $4\pi r^2$

18) A Constituent solids in the given combination of solid figure are

- A] cylinder , cone B] Cylinder , cone , hemisphere
 C] Cylinder , cone , sphere D] cube , cone , hemisphere



19) The length of each edge of a cube with its volume 1331 cm³ is

- (A) 12cm (B) 11cm (C) 15cm (D) 13cm

20) A solid formed on revolving a side of a rectangle is

- A) Cuboid B) Cylinder C) Sphere D) Rightcircular cone

Answers:

1	C	6	D	11	C	16	C
2	A	7	B	12	D	17	B
3	B	8	C	13	A	18	B
4	D	9	D	14	A	19	B
5	C	10	B	15	A	20	B