

WARNING	Any malpractice or any attempt to commit any kind of malpractice in the Examination will DISQUALIFY THE CANDIDATE.		
PAPER - II MATHEMATICS - 2021			
Version Code	B2	Question Booklet Serial Number:	7218187
Time: 150 Minutes	Number of Questions : 120	Maximum Marks : 480	
Name of the Candidate	[REDACTED]		
Roll Number	[REDACTED]		
Signature of the Candidate	[REDACTED]		

INSTRUCTIONS TO CANDIDATES

1. Please ensure that the **VERSION CODE** shown at the top of this Question Booklet is same as that shown in the OMR Answer Sheet issued to you. If you have received a Question Booklet with a different Version code, please get it replaced with a Question Booklet with the same Version Code as that of OMR Answer Sheet from the Invigilator. **THIS IS VERY IMPORTANT.**
2. Please fill the items such as Name, Roll Number and Signature in the columns given above. Please also write Question Booklet Serial Number given at the top of this page against item 3 in the OMR Answer Sheet.
3. This Question Booklet contains 120 questions. For each question five answers are suggested and given against (A), (B), (C), (D) and (E) of which only one will be the **'Most Appropriate Answer'**. Mark the bubble containing the letter corresponding to the 'Most Appropriate Answer' in the OMR Answer Sheet, by using either **Blue or Black Ball Point Pen only.**
4. **Negative Marking:** In order to discourage wild guessing the score will be subjected to penalization formula based on the number of right answers actually marked and the number of wrong answer marked. Each correct answer will be awarded **FOUR** marks. **ONE** mark will be deducted for each incorrect answer. More than one answer marked against a question will be deemed as incorrect answer and will be negatively marked.
5. Please read the instructions in the OMR Answer Sheet for marking the answers. Candidates are advised to strictly follow the instruction contained in the OMR Answer Sheet.

IMMEDIATELY AFTER OPENING THE QUESTION BOOKLET, THE CANDIDATE SHOULD VERIFY WHETHER THE QUESTION BOOKLET CONTAINS ALL THE 120 QUESTIONS IN SERIAL ORDER. IF NOT, REQUEST FOR REPLACEMENT.

DO NOT OPEN THE SEAL UNTIL THE INVIGILATOR ASKS YOU TO DO SO.

SEAL

5. The value of $\tan^{-1}\left(\frac{7}{4}\right) - \tan^{-1}\left(\frac{3}{11}\right)$ is equal to

- (A) $\frac{-\pi}{3}$ (B) $\frac{-\pi}{4}$ (C) $\frac{\pi}{4}$ (D) $\frac{\pi}{3}$ (E) π

6. If $0 < \theta < \frac{\pi}{2}$ and $\tan \theta = \frac{\sqrt{5}}{2}$, then $\cos \theta$ is equal to

- (A) $\frac{1}{2}$ (B) $\frac{\sqrt{3}}{2}$ (C) $\frac{1}{3}$ (D) $\frac{2}{3}$ (E) $\frac{\sqrt{5}}{3}$

7. The value of $\sin^2\left(\cos^{-1}\left(\frac{3}{5}\right)\right)$ is equal to

- (A) $\frac{4}{5}$ (B) $\frac{16}{25}$ (C) $\frac{9}{25}$ (D) $\frac{5}{3}$ (E) $\frac{25}{9}$

8. $\cos^4 \frac{\pi}{12} - \sin^4 \frac{\pi}{12}$ is equal to

- (A) $\frac{1}{2}$ (B) $\frac{\sqrt{3}}{2}$ (C) $\frac{\sqrt{3}+1}{2}$ (D) $\frac{\sqrt{3}-1}{2}$ (E) $\frac{\sqrt{2}}{2}$

9. $\tan\left(2 \tan^{-1}\left(\frac{2}{5}\right)\right)$ is equal to
- (A) $\frac{8}{5}$ (B) $\frac{10}{21}$ (C) $\frac{20}{21}$ (D) $\frac{21}{25}$ (E) $\frac{4}{25}$
10. The values of x in the interval $[0, \pi]$ such that $\sin 2x = \frac{\sqrt{3}}{2}$ are
- (A) $\frac{\pi}{6}, \frac{\pi}{3}$ (B) $\frac{\pi}{6}, \frac{2\pi}{3}$ (C) $\frac{\pi}{3}, \frac{2\pi}{3}$ (D) $\frac{\pi}{6}, \frac{5\pi}{6}$ (E) $\frac{\pi}{3}, \frac{5\pi}{6}$
11. If $\sin \alpha + \sin \beta = \frac{\sqrt{6}}{2}$ and $\cos \alpha + \cos \beta = \frac{\sqrt{2}}{2}$, then $\cos(\alpha - \beta)$ is equal to
- (A) $\frac{1}{2}$ (B) $\frac{3}{2}$ (C) $\frac{-1}{2}$ (D) $\frac{-3}{2}$ (E) 0
12. If $ay = x + b$ is the equation of the line passing through the points $(-5, -2)$ and $(4, 7)$, then the value of $2a + b$ is equal to
- (A) 1 (B) 3 (C) 5 (D) -3 (E) -1

13. The y -intercept of the line passing through $(2, 5)$ with slope $\frac{1}{2}$ is equal to
(A) 1 (B) 2 (C) 3 (D) 4 (E) 5
14. The equation of perpendicular bisector of the line segment joining the points $(10, 0)$ and $(0, -4)$ is
 (A) $5x + 2y = 21$ (B) $5x + 2y = 0$ (C) $2x - 5y = 21$
(D) $5x - 2y = 21$ (E) $2x + 3y = 21$
15. The equation of the line which is parallel to $x + \frac{1}{2}y = \frac{3}{2}$ and passing through $(1, 3)$ is
(A) $2x + y = 7$ (B) $2x + y + 5 = 0$ (C) $2x + y = 3$
(D) $2x + y = 6$ (E) $2x + y = 5$
16. If x -intercept of the straight line $ax + 2ay = 30$ is 10, then the y -intercept is
 (A) 5 (B) 10 (C) 15 (D) 20 (E) 30

17. A straight line makes an angle α with the positive direction of x -axis, where $\cos \alpha = \frac{\sqrt{3}}{2}$. If it passes through $(0, -2)$, then its equation is
- (A) $\sqrt{3}x + y + 2 = 0$ (B) $\sqrt{3}y + x + 2 = 0$ (C) $\sqrt{3}y + x + 2\sqrt{3} = 0$
 (D) $\sqrt{3}y - x + 2\sqrt{3} = 0$ (E) $\sqrt{3}x + y - 2\sqrt{3} = 0$
18. The equation of the circle is $3x^2 + 3y^2 + 6x - 4y - 1 = 0$. Then its radius is
- (A) $\frac{1}{3}$ (B) $\frac{4}{3}$ (C) $\frac{2}{3}$ (D) $\frac{16}{3}$ (E) $\frac{8}{3}$
19. The end-points of a diameter of a circle are $(-1, 4)$ and $(5, 4)$. Then the equation of the circle is
- (A) $(x-3)^2 + y^2 = 9$ (B) $(x-3)^2 + (y+4)^2 = 3$ (C) $(x-2)^2 + (y-4)^2 = 9$
(D) $(x+3)^2 + (y+4)^2 = 9$ (E) $(x-3)^2 + (y-4)^2 = 4$
20. The two diameters of a circle are segments of the straight lines $x - y = 5$ and $2x + y = 4$. If the radius of the circle is 5, then the equation of the circle is
- (A) $x^2 + y^2 - 6x + 4y = 12$ (B) $x^2 + y^2 - 3x + 2y = 12$ (C) $x^2 + y^2 - 6x + 2y = 12$
(D) $x^2 + y^2 - 8x + 6y - 18 = 0$ (E) $x^2 + y^2 - 8x + 6y - 7 = 0$

21. The equation of the parabola with vertex $(-6, 2)$, passing through $(-3, 5)$ and having axis parallel to x -axis is
- (A) $(y+2)^2 = 3x+16$ (B) $(x+6)^2 = 3y-6$ (C) $(y+2)^2 = 4x+48$
 (D) $(x-6)^2 = 4y-8$ (E) $(y-2)^2 = 3x+18$
22. One of the vertices of the major axis of an ellipse is $(1, 1)$ and one of the vertices of its minor axis is $(-2, -1)$. If the centre of the ellipse is $(-2, 1)$, then the equation of the ellipse is
- (A) $\frac{(x+2)^2}{9} + \frac{(y-1)^2}{4} = 1$ (B) $\frac{(x+2)^2}{16} + \frac{(y-1)^2}{4} = 1$ (C) $\frac{(x-2)^2}{9} + \frac{(y+1)^2}{4} = 1$
 (D) $\frac{(x-2)^2}{16} + \frac{(y+1)^2}{4} = 1$ (E) $\frac{(x+2)^2}{9} + \frac{(y-1)^2}{2} = 1$
23. The equation of the parabola with focus $(3, 0)$ and directrix $x+3=0$ is
- (A) $y^2 = 3x-9$ (B) $y^2 = 4x-12$ (C) $y^2 = 12x$
 (D) $y^2 = 12x-36$ (E) $y^2 = 12x-9$

24. The eccentricity of the ellipse $\frac{x^2}{36} + \frac{y^2}{16} = 1$ is

- (A) $\frac{\sqrt{5}}{3}$ (B) $\frac{\sqrt{5}}{6}$ (C) $\frac{\sqrt{30}}{6}$ (D) $\frac{\sqrt{10}}{6}$ (E) $\frac{\sqrt{30}}{7}$

25. The foci of a hyperbola are $(8, 3)$ and $(0, 3)$ and eccentricity is $\frac{4}{3}$. Then the length of the transverse axis is

- (A) $\frac{32}{3}$ (B) 4 (C) 8 (D) $\frac{8}{3}$ (E) 6

26. The co-ordinates of the points P and Q are $(2, 6, 4)$ and $(8, -3, 1)$ respectively. If the point R lies on the line segment PQ such that $2|\vec{PR}| = |\vec{RQ}|$, then the co-ordinates of R are

- (A) $(4, -3, 3)$ (B) $(4, 3, -3)$ (C) $(2, -3, 1)$ (D) $(4, 3, 3)$ (E) $(2, 3, 3)$

27. If $|\vec{a}| = 2$, $\vec{b} = 2\hat{i} - \hat{j} - 3\hat{k}$ and the angle between \vec{a} and \vec{b} is $\frac{\pi}{4}$, then $\vec{a} \cdot \vec{b}$ is equal to

- (A) $14\sqrt{2}$ (B) $2\sqrt{7}$ (C) $\sqrt{30}$ (D) $\sqrt{7}$ (E) $\sqrt{14}$

28. If α is the angle made by the vector $\vec{a} = 5\hat{i} + 3\hat{j} + 4\hat{k}$ with the positive x-axis, then $\cos \alpha =$
- (A) $\frac{5}{12}$ (B) $\frac{1}{2}$ (C) $\frac{\sqrt{2}}{2}$ (D) $\frac{\sqrt{5}}{5}$ (E) $\frac{\sqrt{2}}{10}$
29. If $|\vec{a}| = 3, |\vec{b}| = 4$ and $|\vec{a} - \vec{b}| = \sqrt{7}$, then $\vec{a} \cdot \vec{b}$ is equal to
- (A) 7 (B) 8 (C) 9 (D) 10 (E) 12
30. If $\vec{a} = \hat{i} + \lambda\hat{j} - 2\hat{k}$, $\vec{b} = 2\hat{i} - 3\hat{j} + 5\hat{k}$ and $\vec{a} \cdot \vec{b} = -20$, then the value of λ is equal to
- (A) 2 (B) -2 (C) -4 (D) 4 (E) 5
31. If $\vec{a} = \hat{i} - 3\hat{j} + \alpha\hat{k}$, $\vec{b} = \hat{i} - 2\hat{j} + 4\hat{k}$ and $\vec{a} \times \vec{b} = -2\hat{i} + \hat{j} + \beta\hat{k}$, then the value of β is equal to
- (A) -2 (B) 2 (C) -1 (D) 1 (E) -3

32. The values of α so that the vectors $\alpha \hat{i} + (\alpha - 1) \hat{j} + 3 \hat{k}$ and $(\alpha + 2) \hat{i} + \alpha \hat{j} - 2 \hat{k}$ are perpendicular, are
- (A) $\frac{3}{2}, -2$ (B) $2, \frac{3}{2}$ (C) $-2, \frac{-3}{2}$ (D) $2, \frac{-3}{2}$ (E) $-4, \frac{3}{2}$
33. If $|\vec{u}| = 5$, $|\vec{v}| = 4$ and the angle between \vec{u} and \vec{v} is $\frac{\pi}{6}$, then $|\vec{u} \times \vec{v}|$ is equal to
- (A) $10\sqrt{3}$ (B) $10\sqrt{2}$ (C) 20 (D) $5\sqrt{2}$ (E) 10
34. If the point $P(x, 1, 4)$ lies on the line $\vec{r} = \hat{i} + 3\hat{j} + 4\hat{k} + \lambda(2\hat{i} - \hat{j})$, then the value of x is equal to
- (A) 2 (B) -2 (C) 3 (D) -3 (E) 5
35. The equation of the plane through the point $(2, 1, 3)$ and perpendicular to the vector $4\hat{i} + 5\hat{j} + 6\hat{k}$ is
- (A) $4x + 5y + 6z = 28$ (B) $2x + y + 3z = 17$ (C) $4x + 5y + 6z = 33$
(D) $8x + 5y + 18z = 21$ (E) $4x + 5y + 6z = 31$

36. The angle between the line $\vec{r} = \hat{i} + 2\hat{j} + t(3\hat{i} + 2\hat{j} - \hat{k})$ and the plane $2x - 3y - z = 1$ is

- (A) $\sin^{-1}\left(\frac{1}{196}\right)$ (B) $\sin^{-1}\left(\frac{1}{14}\right)$ (C) $\cos^{-1}\left(\frac{1}{14}\right)$ (D) $\cos^{-1}\left(\frac{13}{14}\right)$ (E) $\sin^{-1}\left(\frac{13}{14}\right)$

37. If the line $\vec{r} = 2\hat{i} + \hat{j} + t(3\hat{i} + \hat{j} - 2\hat{k})$ is parallel to the plane $2x + 4y + az = 8$, then the value of a is equal to

- (A) 2 (B) 3 (C) 4 (D) 5 (E) 6

38. The angle between the lines $\vec{r} = \hat{i} + 4\hat{k} + \lambda(2\hat{i} + \hat{j} - \hat{k})$ and $\vec{r} = 2\hat{i} - \hat{j} + 3\hat{k} + \mu(3\hat{i} + \hat{k})$ is

- (A) $\cos^{-1}\left(\frac{\sqrt{5}}{6}\right)$ (B) $\cos^{-1}\left(\frac{\sqrt{15}}{6}\right)$ (C) $\cos^{-1}\left(\frac{1}{12}\right)$ (D) $\cos^{-1}\left(\frac{\sqrt{15}}{15}\right)$ (E) $\cos^{-1}\left(\frac{\sqrt{3}}{30}\right)$

39. The Cartesian equation of the line passing through $(7, 5, 3)$ and perpendicular to the plane $3x + 2y + z = 6$ is

- (A) $\frac{x-7}{3} = \frac{y-5}{2} = \frac{z-3}{1}$ (B) $\frac{x-3}{7} = \frac{y-2}{5} = \frac{z-1}{3}$ (C) $\frac{x-3}{7} = \frac{y-2}{5} = \frac{z}{3}$
(D) $\frac{x-7}{3} = \frac{y-5}{1} = \frac{z-3}{2}$ (E) $\frac{x-4}{4} = \frac{y-3}{3} = \frac{z-2}{2}$

40. The acute angle between the planes $2x - y - 3z = 7$ and $x + 2y + 2z = 0$ is

(A) $\cos^{-1}\left(\frac{-\sqrt{14}}{14}\right)$

(B) $\pi - \cos^{-1}\left(\frac{-\sqrt{14}}{7}\right)$

(C) $\cos^{-1}\left(\frac{\sqrt{14}}{11}\right)$

(D) $\pi - \cos^{-1}\left(\frac{-\sqrt{14}}{21}\right)$

(E) $\pi - \cos^{-1}\left(\frac{\sqrt{14}}{7}\right)$

41. The vector equation of the line joining the points $(2, 1, 3)$ and $(-2, 4, 1)$ is

(A) $\vec{r} = 2\hat{i} + \hat{j} + 3\hat{k} + \lambda(-4\hat{i} + 3\hat{j} - 2\hat{k})$

(B) $\vec{r} = 2\hat{i} + \hat{j} + 3\hat{k} + \lambda(4\hat{i} + 3\hat{j} + 2\hat{k})$

(C) $\vec{r} = -2\hat{i} + \hat{j} + 3\hat{k} + \lambda(-4\hat{i} - 3\hat{j} - 2\hat{k})$

(D) $\vec{r} = 2\hat{i} + \hat{j} + 3\hat{k} + \lambda(3\hat{i} - 4\hat{j} - 2\hat{k})$

(E) $\vec{r} = -4\hat{i} + 3\hat{j} - 2\hat{k} + \lambda(2\hat{i} + \hat{j} + 3\hat{k})$

42. A bag contains 5 yellow, 3 green, 2 blue and 7 white balls. If 4 balls are chosen at random, then the probability that none of them are white is

(A) $\frac{3}{37}$

(B) $\frac{7}{34}$

(C) $\frac{5}{34}$

(D) $\frac{5}{37}$

(E) $\frac{3}{34}$

43. An urn contains 25 marbles which are numbered from 1 to 25 and a marble is chosen at random two times with replacement. Then the probability that both times the marble has the same number is

(A) $\frac{1}{25}$

(B) $\frac{24}{25}$

(C) $\frac{1}{625}$

(D) $\frac{624}{625}$

(E) $\frac{2}{25}$

44. If A and B are two events such that $P(A) = 0.2$, $P(B) = 0.55$ and $P(A \cap B) = 0.1$, then $P(B \cap A^c)$ is equal to
 (A) 0.25 (B) 0.35 (C) 0.45 (D) 0.65 (E) 0.75
45. Two dice are rolled. If A is the event that sum of the numbers is 4 and B is the event that at least one of the dice shows a 3, then $P(A|B)$ is equal to
 (A) $\frac{3}{11}$ (B) $\frac{2}{11}$ (C) $\frac{1}{4}$ (D) $\frac{1}{6}$ (E) $\frac{1}{11}$
46. Assume that n distinct values x_1, x_2, \dots, x_n occur with frequencies f_1, f_2, \dots, f_n respectively. If $\bar{x} = 7$ and $\sum_{i=1}^n f_i x_i = 315$, then $\sum_{i=1}^n f_i =$
 (A) 35 (B) 45 (C) 48 (D) 42 (E) 40
47. The variance of the data x_1, x_2, \dots, x_{50} with $\sum_{i=1}^{50} x_i = 650$ and $\sum_{i=1}^{50} x_i^2 = 10000$ is
 (A) 30 (B) 40 (C) 39 (D) 41 (E) 31

48. If X is a random variable with $E(X) = 6$ and $V(X) = 3$, then $E(X^2)$ is equal to
(A) 33 (B) 36 (C) 39 (D) 42 (E) 27
49. Let $f(x) = \frac{4x+3}{x+2}$. Then the value of $f^{-1}(-2)$ is equal to
(A) $\frac{7}{5}$ (B) $\frac{-7}{6}$ (C) $\frac{-7}{5}$ (D) $\frac{7}{6}$ (E) $\frac{5}{6}$
50. If $f(x) = \begin{cases} 2x & \text{for } x < 1 \\ 5a - x & \text{for } x \geq 1 \end{cases}$ is continuous on \mathbb{R} , then the value of a is equal to
(A) $\frac{1}{5}$ (B) $\frac{2}{5}$ (C) $\frac{3}{5}$ (D) $\frac{4}{5}$ (E) 1
51. $\lim_{t \rightarrow 0} \frac{\sin 2t}{8t^2 + 4t}$ is equal to
 (A) $\frac{1}{2}$ (B) $\frac{2}{5}$ (C) $\frac{1}{6}$ (D) $\frac{1}{3}$ (E) 1



52. $\lim_{x \rightarrow 0} \frac{x}{\sqrt{9-x}-3}$ is equal to
(A) 6 (B) 3 (C) -3 (D) -6 (E) 0
53. Let $f(x) = \begin{cases} 3x+2, & \text{if } x < -2 \\ x^2-3x-1, & \text{if } x \geq -2 \end{cases}$. Then $\lim_{x \rightarrow -2^-} f(x)$ and $\lim_{x \rightarrow -2^+} f(x)$ are respectively
(A) -4, 3 (B) 6, 3 (C) -6, 3 (D) -4, 9 (E) 9, -4
54. $\lim_{x \rightarrow -3} \frac{x^2+16x+39}{2x^2+7x+3}$ is equal to
(A) 2 (B) $\frac{8}{3}$ (C) $\frac{-8}{3}$ (D) -2 (E) 0
55. Let $f(x) = 6\sqrt[3]{x^5}$. If $f'(x) = ax^p$, where a and p are constants, then the value of p is equal to
(A) $\frac{3}{5}$ (B) $\frac{-2}{5}$ (C) $\frac{2}{3}$ (D) $\frac{-2}{3}$ (E) $\frac{2}{5}$

56. Let $y = (\tan x)^{\sin x}$ for $0 < x < \frac{\pi}{2}$. If $\frac{dy}{dx} = (\tan x)^{\sin x} ((\cos x) \log(\tan x) + g(x))$, then

$g(x) =$

(A) $\sin x \sec^2 x$

(B) $\sec x \operatorname{cosec} x$

(C) $\sec x$

(D) $\operatorname{cosec} x$

(E) $\sin x \tan x$

57. If $f(x) = (x^3 + \sin \pi x)^5$, then $f'(1)$ is equal to

(A) 2^5

(B) $5(2^4)$

(C) 15

(D) $5(3 + \pi)$

(E) $5(3 - \pi)$

58. If $h(x) = 4x^3 - 5x + 7$ is the derivative of $f(x)$, then $\lim_{t \rightarrow 0} \frac{f(1+t) - f(1)}{t}$ is equal to

(A) 5

(B) 6

(C) 7

(D) 8

(E) 0

59. Let $f(x) = \begin{cases} e^x, & \text{if } x \leq 1 \\ mx + 6, & \text{if } x > 1 \end{cases}$ be differentiable at $x = 1$. Then the value of m is

(A) 6

(B) e

(C) -6

(D) $-e$

(E) 1

60. $\lim_{t \rightarrow 0} \frac{\tan^2\left(\frac{\pi}{3} + t\right) - 3}{t}$ is equal to
 (A) $4\sqrt{3}$ (B) 24 (C) $16\sqrt{3}$ (D) $8\sqrt{3}$ (E) 16
61. If the tangent line to the graph of a function f at the point $x = 3$ has x -intercept $\frac{5}{3}$ and y -intercept -10 , then $f'(3)$ is equal to
 (A) 3 (B) 5 (C) $\frac{5}{3}$ (D) 6 (E) -10
62. The slope of tangent line to the curve $4x^2 + 2xy + y^2 = 12$ at the point $(1, 2)$ is
 (A) 2 (B) 1 (C) -1 (D) -2 (E) 0
63. Let $f(x) = \sqrt{x} + 5$ for $1 \leq x \leq 9$. Then the value of c whose existence is guaranteed by the Mean Value Theorem is
 (A) 2 (B) 3 (C) 4 (D) 5 (E) 6
64. The derivative of a function f is given by $f'(x) = \frac{x-5}{\sqrt{x^2+4}}$. Then the interval in which f is increasing, is
 (A) $(5, \infty)$ (B) $(0, \infty)$ (C) $(-4, \infty)$ (D) $(-\infty, -4)$ (E) $(-\infty, 5)$

65. Let $f(x) = x^2 \log x, x > 0$. Then the minimum value of f is

- (A) $\frac{1}{\sqrt{e}}$ (B) $2e$ (C) $-2e$ (D) \sqrt{e} (E) $\frac{-1}{2e}$

66. A cube is expanding in such a way that its edge is increasing at a rate of 2 inches per second. If its edge is 5 inches long, then the rate of change of its volume is

- (A) $150 \text{ in}^3/\text{sec}$ (B) $75 \text{ in}^3/\text{sec}$ (C) $50 \text{ in}^3/\text{sec}$
 (D) $30 \text{ in}^3/\text{sec}$ (E) $45 \text{ in}^3/\text{sec}$

67. $\int x^5 e^{1-x^6} dx =$

- (A) $\frac{1}{6} e^{1-x^6} + C$ (B) $-e^{1-x^6} + C$ (C) $\frac{1}{6} e^{1-x^6} + C$
 (D) $\frac{x^5}{5} e^{1-x^6} + C$ (E) $\frac{x^6}{6} e^{1-x^6} + C$

68. $\int (5-4x)e^{-x} dx =$

- (A) $e^{-x}(4x-1)+C$ (B) $e^{-x}(9-4x)+C$ (C) $e^{-x}(4x-5)+C$
 (D) $e^{-x}(4x-9)+C$ (E) $e^{-x}(5-4x)+C$



69. $\int \frac{\cos(\tan x)}{\cos^2 x} dx =$

(A) $(\tan x)\sin(\tan x) + C$

(B) $\sin(\tan x) + C$

(C) $\sec(\tan x) + C$

(D) $(\cos x)\sin(\tan x) + C$

(E) $\cos^2(\tan x) + C$

70. $\int \frac{1}{e^{2x} - 1} dx =$

(A) $2 \log|e^{2x} - 1| - x + C$

(B) $x - \frac{1}{2} \log|e^{2x} - 1| + C$

(C) $x + \frac{1}{2} \log|e^{2x} - 1| + C$

(D) $x - \log|e^{2x} - 1| + C$

(E) $\frac{1}{2} \log|e^{2x} - 1| - x + C$

71. $\int \sin 2x \cos x dx =$

(A) $\frac{-1}{3} \cos^3 x + C$

(B) $\frac{-2}{3} \cos^3 x + C$

(C) $\frac{2}{3} \cos^3 x + C$

(D) $\frac{1}{3} \cos^3 x + C$

(E) $\frac{-4}{3} \cos^3 x + C$

72. $\int \frac{1}{(1 + \cot^2 x)\sin^2 x} dx =$

(A) $\tan^{-1}(\sin x) + C$

(B) $\tan^{-1}(\cos x) + C$

(C) $\cot^{-1}(\sin x) + C$

(D) $\cot^{-1}(\cos x) + C$

(E) $x + C$

73. $\int \frac{4x^9}{x^{10}-10} dx =$

(A) $\frac{1}{5} \log|x^{10}-10| + C$

(B) $\frac{2}{5} \log|x^{10}-10| + C$

(C) $\frac{1}{10} \log|x^{10}-10| + C$

(D) $\frac{-2}{5} \log|x^{10}-10| + C$

(E) $\frac{-1}{10} \log|x^{10}-10| + C$

74. The value of $\int_0^{\sqrt{3}} \frac{6}{9+x^2} dx$ is equal to

(A) $\frac{\pi}{3}$

(B) $\frac{\pi}{6}$

(C) $\frac{\pi}{4}$

(D) $\frac{2\pi}{3}$

(E) 1

75. The value of $\int_{-5}^5 (4-|x|) dx$ is equal to

(A) 18

(B) 10

(C) 12

(D) 16

(E) 15

76. The area of the region bounded by the curves $y=x^2$ and $y=\sqrt{x}$ is (in square units)

(A) $\frac{2}{3}$

(B) $\frac{1}{3}$

(C) $\frac{1}{6}$

(D) $\frac{5}{6}$

(E) 1

77. The value of $\int_0^2 \frac{x^2}{(x^3+1)^2} dx$ is equal to

(A) $\frac{1}{27}$

(B) $\frac{5}{27}$

(C) $\frac{7}{27}$

(D) $\frac{8}{27}$

(E) $\frac{1}{3}$

78. The value of $\int_{\pi/8}^{3\pi/8} \frac{\sin^4 x}{\sin^4 x + \cos^4 x} dx$ is equal to

(A) $\frac{\pi}{4}$

(B) $\frac{\pi}{8}$

(C) $\frac{\pi}{16}$

(D) $\frac{\pi}{2}$

(E) 1

79. The area of the region bounded by $y = 5x$, x -axis and $x = 4$ is (in square units)

(A) 40

(B) 80

(C) 20

(D) 50

(E) 60

80. The general solution of the differential equation $y - xy' = x^2 + y^2$ is

(A) $y = x \tan(C-x)$

(B) $y = \tan x + C$

(C) $y = x^2 \tan x + C$

(D) $y = x \tan x + C$

(E) $y = x \tan x + Cx$

81. The integrating factor of the differential equation $xy' + 2y - 7x^3 = 0$ is
(A) $\log|x|$ (B) x^2 (C) $\frac{1}{x^2}$ (D) $\frac{1}{2}\log|x|$ (E) x
82. The general solution of the differential equation $4xy' + 12x + (2x^2 + 3)y' = 0$ is
(A) $\frac{2x^2 + 3}{y + 3} = C$ (B) $\frac{y - 3}{2x^2 + 3} = C$ (C) $\frac{y + 2}{2x^2 + 3} = C$
(D) $(y - 3)(2x^2 + 3) = C$ (E) $(y + 3)(2x^2 + 3) = C$
83. The constraints of a linear programming problem are $x + 2y \leq 10$ and $6x + 3y \leq 18$. Which of the following points lie in the feasible region?
(A) (0, 6) (B) (4, 3) (C) (5, 7) (D) (1, 7) (E) (1, 3)

www.scienceinstitute.in info@scienceinstitute.in

REPEATERS

One year full time entrance coaching **2021-22**

Admission Started

Medical: **NEET, AIIMS, JIPMER etc.**

Engg: **JEE Main, JEE Advanced, KEAM**

COURSE @ **MANJERI & PERINTHALMANNA**

Ph : 0483 2768340 | 04933 223221

HOSTEL FACILITY AVAILABLE FOR BOYS & GIRLS

84. Let $f: [-4, 2] \rightarrow \mathbb{R}$ be given by $f(x) = \sqrt{16 - x^2}$. Then the range of the function f is
(A) $[0, 2]$ ~~(B) $[0, 2\sqrt{3}]$~~ (C) $[0, 4]$ (D) $[2\sqrt{3}, 4]$ (E) $[-2, 2]$
85. Let $f(x) = x^2$ and $g(x) = \sqrt{9+x}$. Then the value of $(f \circ g - g \circ f)(4)$ is equal to
(A) 6 (B) $\sqrt{6}$ (C) $\sqrt{8}$ (D) 8 (E) 5
86. Let A and B be subsets of the universal set U . If $n(A) = 24$, $n(A \cap B) = 8$ and $n(U) = 63$, then $n(A' \cup B')$ is equal to
(A) 43 (B) 55 (C) 35 (D) 32 (E) 45
87. Let $f(x) = [x]$, $x \in \mathbb{R}$, where $[x]$ denotes the greatest integer $\leq x$. Then the images of the elements -4.6 and 2.7 are respectively
 (A) $-5, 2$ (B) $-5, 3$ (C) $-4, 2$ (D) $-3, 3$ (E) $-4, 3$

88. For any two positive rational numbers m and n , a binary operation $*$ is defined by

$$m * n = \frac{m+n}{3}, \text{ then } \frac{7}{2} * \frac{5}{2} \text{ is equal to}$$

- (A) 4 (B) 6 (C) 2 (D) 8 (E) 9

89. The function $f: \mathbb{R} \rightarrow \mathbb{R}$ given by $f(x) = 7 - 3x$ is

- (A) not one-one (B) not onto (C) even (D) one-one and onto (E) odd

90. A relation R on $\{0, 1, 2\}$ is given by $R = \{(0, 0), (1, 1), (0, 1), (2, 2), (1, 2)\}$. Then the relation R is

- (A) reflexive (B) symmetric (C) transitive
(D) symmetric and transitive (E) equivalence

91. Let z_1, z_2 and z_3 be three distinct points in the complex plane such that the segment joining z_1 and z_2 is perpendicular to the segment joining z_1 and z_3 . If $|z_1 - z_2| = 5$ and $|z_1 - z_3| = 12$ then $|z_2 - z_3|$ is equal to

- (A) 17 (B) 7 (C) 13 (D) 14 (E) 9

92. If $\frac{z}{i} = 11 - 13i$, then $z + \bar{z}$ is equal to

- (A) -22 (B) 22 (C) 25 (D) 26 (E) -26

93. Let $\alpha = 2 - 3i$ be a root of the equation $z^2 - 4z + k = 0$, where k is a real number. If β is the other root, then the value of $\alpha^2 + \beta^2$ is
- (A) 26 (B) -5 (C) 5 (D) 10 (E) -10
94. If $z = 2 - i\sqrt{3}$, then $|z^4|$ is equal to
- (A) 7 (B) $\sqrt{7}$ (C) $7\sqrt{7}$ (D) 49 (E) $49\sqrt{7}$
95. The imaginary part of $z = \frac{2+i}{3-i}$ is
- (A) $\frac{5}{8}$ (B) $\frac{-5}{8}$ (C) $\frac{1}{2}$ (D) $\frac{3}{4}$ (E) $\frac{3}{8}$
96. The area of the triangle on the complex plane formed by the points z , $z + iz$ and iz is 128. Then the value of $|z|$ is
- (A) 12 (B) 16 (C) 18 (D) 17 (E) 19
97. If the real part of the complex number $z = \frac{p+2i}{p-i}$, $p \in \mathbb{R}$, $p > 0$ is $\frac{1}{2}$, then the value of p is equal to
- (A) $\sqrt{2}$ (B) $\sqrt{3}$ (C) $\sqrt{5}$ (D) $\frac{\sqrt{3}}{2}$ (E) 1

98. The value of $\sqrt{-25} + 3\sqrt{-4} + 2\sqrt{-9}$ is equal to
(A) $13i$ (B) $-13i$ (C) $11i$ (D) $-17i$ (E) $17i$
99. The value of $\sum_{k=5}^{36} \frac{1}{k^2 - k}$ is
(A) $\frac{7}{36}$ (B) $\frac{1}{9}$ (C) $\frac{2}{9}$ (D) $\frac{1}{12}$ (E) $\frac{5}{36}$
100. If $a_1, a_2, a_3, \dots, a_n$ are in A. P. with $a_1 = 3$, $a_n = 39$ and $a_1 + a_2 + \dots + a_n = 210$, then the value of n is equal to
(A) 8 (B) 10 (C) 11 (D) 13 (E) 15
101. Let $t_n, n = 1, 2, 3, \dots$ be the n^{th} term of the A. P. 5, 8, 11, Then the value of n for which $t_n = 305$ is
(A) 101 (B) 100 (C) 103 (D) 99 (E) 95
102. If the first term of a G. P. is 1 and the sum of 3rd and 5th terms is 90, then the positive common ratio of the G. P. is
(A) 1 (B) 2 (C) 3 (D) 4 (E) 5

103. In an A.P. the difference between the last and the first terms is 632 and the common difference is 4. Then the number of terms in the A. P. is
(A) 157 (B) 160 (C) 158 (D) 159 (E) 140
104. If the 10th and 12th terms of an A. P. are respectively 15 and 21, then the common difference of the A. P. is
(A) -6 (B) 4 (C) 6 (D) -3 (E) 3
105. The first term of a G. P. is 3 and the common ratio is 2. Then the sum of first eight terms of the G.P. is
(A) 763 (B) 189 (C) 381 (D) 765 (E) 655
106. A covid-19 vaccination reduces the probability of getting covid-19 infection from 0.4 to 0.1. In a city, 45% people are vaccinated. Then the probability that a non-vaccinated person chosen at random in the city gets covid-19 infection is
(A) 0.55 (B) 0.45 (C) 0.32 (D) 0.22 (E) 0.18
107. The number of ways a committee of 3 women and 5 men can be formed from a panel of 8 men and 5 women is
(A) 940 (B) 1120 (C) 560 (D) 760 (E) 520

www.scienceinstitute.in info@scienceinstitute.in

REPEATERS

One year full time entrance coaching **2021-22**

Admission Started

Medical: **NEET, AIIMS, JIPMER etc.**

Engg: **JEE Main, JEE Advanced, KEAM**

COURSE @ **MANJERI & PERINTHALMANNA**

Ph : 0483 2768340 | 04933 223221

HOSTEL FACILITY AVAILABLE FOR BOYS & GIRLS

108. A set contains 9 elements. Then the number of subsets of the set which contains at most 4 elements is
(A) 32 (B) 64 (C) 128 (D) 256 (E) 512
109. If p and q are positive integers such that ${}^{(p+q)}P_2 = 42$ and ${}^{(p-q)}P_2 = 20$, then the values of p and q are respectively
(A) 5, 2 (B) 4, 3 (C) 7, 2 (D) 6, 1 (E) 7, 5
110. The number of 3-digit numbers that can be formed from the digits 0, 2, 3, 5, 7 is (repetition is allowed)
(A) 125 (B) 100 (C) 105 (D) 150 (E) 60
111. If x^{22} is in the $(r+1)^{\text{th}}$ term of the binomial expansion of $(3x^3 - x^2)^9$, then the value of r is equal to
(A) 3 (B) 4 (C) 5 (D) 6 (E) 7

112. The term independent of x in the binomial expansion of $\left(x + \frac{2}{x^3}\right)^{20}$ is

(A) $\binom{20}{5} 2^{15}$

(B) $\binom{20}{15} 2^{10}$

(C) $\binom{20}{10} 2^5$

(D) $\binom{20}{10} 2^{10}$

✓ (E) $\binom{20}{5} 2^5$

113. Let $A + B = \begin{bmatrix} 4 & 1 & 4 \\ 1 & 4 & 4 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 0 & -2 \\ -1 & 3 & 0 \end{bmatrix}$, then $A =$

(A) $\begin{bmatrix} 3 & 1 & 2 \\ 0 & 3 & 4 \end{bmatrix}$

(B) $\begin{bmatrix} 5 & 1 & 2 \\ 0 & 7 & 4 \end{bmatrix}$

(C) $\begin{bmatrix} 3 & -1 & -2 \\ 2 & 1 & 4 \end{bmatrix}$

(D) $\begin{bmatrix} 5 & 1 & 6 \\ 2 & 1 & 4 \end{bmatrix}$

✓ (E) $\begin{bmatrix} 3 & 1 & 6 \\ 2 & 1 & 4 \end{bmatrix}$

114. The value of the determinant $\begin{vmatrix} 4 & 4^2 & 4^3 \\ 3 & 3^2 & 3^3 \\ 2 & 2^2 & 2^3 \end{vmatrix}$ is

(A) 52

(B) -24

(C) 24

(D) 48

✓ (E) -48

115. If $\begin{vmatrix} 1 & 2 & 1 \\ 0 & x & -3 \\ 2 & -1 & x \end{vmatrix} = 0$, then the values of x are

✓ (A) 5, -3

(B) 5, 3

(C) -5, 3

(D) 2, 3

(E) -2, -3

116. If $AB = \begin{bmatrix} 4 & 3 \\ 5 & 4 \end{bmatrix}$ and $A^{-1} = \begin{bmatrix} 3 & -2 \\ -1 & 1 \end{bmatrix}$, then $B =$

(A) $\begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix}$

(B) $\begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$

(C) $\begin{bmatrix} 1 & 2 \\ 1 & 1 \end{bmatrix}$

(D) $\begin{bmatrix} 1 & 1 \\ 1 & 2 \end{bmatrix}$

(E) $\begin{bmatrix} 2 & 1 \\ 1 & 1 \end{bmatrix}$

117. The matrix $\begin{bmatrix} -2 & 1 & 0 \\ 3 & 4 & 1 \\ -4 & \lambda & 0 \end{bmatrix}$ is non-singular for $\lambda \neq$

(A) 2

(B) -2

(C) 4

(D) -4

(E) 0

118. Let $\begin{vmatrix} x-1 & 2 & 1 \\ 2 & x-1 & 2 \\ 1 & x+2 & x-1 \end{vmatrix} = ax^3 + bx^2 + cx + d$, where a, b, c and d are constants. Then the value of d is

(A) -8

(B) 6

(C) 0

(D) -6

(E) 16

119. If the inequality $-13 \leq x \leq 5$ is expressed in the form $|x - a| \leq b$, then the values of a and b are respectively

(A) 4, 8

(B) -4, 9

(C) 4, 9

(D) 5, 9

(E) -5, 9

120. The solution set of the inequality $5(4x + 6) < 25x + 10$ is

(A) $(4, \infty)$

(B) $(-\infty, 4)$

(C) $(-\infty, 5)$

(D) $(5, \infty)$

(E) $(-4, 4)$

REPEATERS 2022

NEET | JEE | KEAM

ONLINE CLASSES 

ONLINE + OFFLINE CLASSES 

APPLY NOW 

www.scienceinstitute.in

Next Batch: 09-08-2021

ONLINE: 9567 300 111

MANJERI: 0483-2768340

PERINTHALMANNA: 9567 300 333

.....
Hostel Facility Available for Boys & Girls



SCIENCE INSTITUTE

THE PREMIER INSTITUTE FOR ENTRANCE COACHING IN THE STATE