

- 1) Deltoid ligament doesn't join med cuneiform.
- 2) Scalenus anterior - wrong statement - subclavian art. lies anteriorly
- 3) Pain along medial aspect of arm through in Ca Breast - intercostobrachial nerve
- 4) Laminae in substantia gelatinosa - 2+3
- 5) Muscle in deep perineal pouch - sphincter urethrae
- 6) Persistent primitive streak - Sacrococcygeal teratoma
- 7) Carpal tunnel syn - median nerve
- 8) Pudendal nerve - S2-4
- 9) Pelvic diaphragm consist of - levator ani
- 10) Lateral displacement of knee prevented by -
- 11) Abduction of toes is by - Dorsal interossei
- 12) Base of 5th metatarsal fracture pull of - Peronius brevis
- 13) Ext rotation in semi flexed knee all except - Fibular Collateral Ligament
- 14) Does not pass under extensor retinaculum - Ant interosseus artery
- 15) Section of fibular nerve - Foot drop
- 16) Not a part of epithalamus is - (parts are 1. pineal, 2. habenular nuclei, trigone & commissure 3. stria medullaris)
- 17) Striae of Gennari seen in - also called the "band" or "line" of Gennari is a band of myelinated axons projecting into layer 4C of the primary visual cortex from the lateral geniculate nucleus
- 18) Porto systemic shunt is present at A/E - Lower anal canal 19) Structure not passing through inf cerebellar peduncle - Pontocerebellar 20) Ophthalmic artery is Branch of - Intra cranial Int. carotid
- 21) Nucleus ambiguus true is - 9th and 10th nerve arise from it.
- 22) Not a part of Internal Capsule - Optic radiation
- 23) Edinger Westphal nucleus - General visceral efferent
- 24) Uterine epithelium develops from - Fusion of paramesonephric ducts
- 25) Boundries of pudendal canal - Between obturator membrane and Facia lunata
- 26) programmed cell death is called - apoptosis
- 27) stable cells are ? liver cells
- 28) christmas tree stain is used for - identification of semen in a forensic sample.
- 29) telomerase is - RNA dependent DNA polymerase
- 30) HGPRT is used for the salvage of which of these - hypoxanthine and guanine