

ANSWER KEY

①

SECOND YEAR HIGHER SECONDARY EXAMINATION MARCH 2022

PART-I/H/III

SUBJECT: STATISTICS

CODE NO: SY 32

VERSION: _____

60 SCORES

2 HOURS

Qn. No	Sub Qns	Answer Key/Value Points	Score	Total Score
1		(a) 0	1	1
2		(b) 1	1	1
3		(d) Walter A. Schewhart	1	1
4		(c) $E(x) = 4$ and $V(x) = 4$	1	1
5.		(a) 1	1	1
6.		(d) Sampling	1	1
7.		(b) 15, 15	1	1
8		(d) 124.9	1	1
9		(c) 1.75	1	1
10.		$3x^2 + 1$	1	1
11.		(d) 16	1	1
12.		(b) 12	1	1
13.		$Z_1 = 1.44$	1	1
14		(i) - (d), (ii) - (c), (iii) - (b), (iv) - (a)	$\frac{1}{2} \times 4$	2
15.		$UCL = \bar{\bar{x}} + A_2 \bar{R} = 16.2 + 0.577 \times 7.4 = 20.47$ $LCL = \bar{\bar{x}} - A_2 \bar{R} = 16.2 - 0.577 \times 7.4 = 11.93$ <p style="text-align: center;">only [If $CL = \bar{\bar{x}} = 16.2$, give 1 score]</p>	$\frac{1}{2} + \frac{1}{2}$ $\frac{1}{2} + \frac{1}{2}$	2.
16	(a)	Type I error - definition	1	2
	(b)	Type II error - definition	1	

Qn. No	Sub Qns	Answer Key/Value Points	Score	Total Score
17		$y - \bar{y} = by_x (x - \bar{x})$ $y - 27 = -1.5(x - 53)$ <p>When $x = 60$, $y - 27 = -1.5(60 - 53)$</p> $y = 16.5$	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$	2
18		<p>Point estimation - Definition/ Example / Explanation</p> <p>Interval estimation - Definition/ Example / Explanation</p>	1 1	2
19.		$Z = \frac{x - \mu}{\sigma} = \frac{x - 68}{\sigma} \sim N(0,1)$ $P(x > 72) = P\left(z > \frac{72 - 68}{\sigma}\right) = 0.1587$ $\frac{4}{\sigma} = 1$ $\sigma = 4$	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$	2
20		<p>Solving the equations</p> $\bar{x} = 1, \bar{y} = 2$	1 $\frac{1}{2} + \frac{1}{2}$	2
21	(a)	$\int 8 dx = 8x + c$ / $8x$	1	3
	(b)	$\int_0^1 (x+1) dx = \left[\frac{x^2}{2} + x \right]_0^1$ $= \left(\frac{1}{2} + 1 \right) - (0)$ $= 1.5$	1 $\frac{1}{2}$ $\frac{1}{2}$	
22		$\mu = 50, \sigma = 10$ $Z = \frac{x - \mu}{\sigma} = \frac{x - 50}{10} \sim N(0,1)$ $P(x < 70) = P\left(z < \frac{70 - 50}{10}\right) = P(z < 2)$ $= 0.5 + 0.4772 = 0.9772$	$\frac{1}{2}$ $\frac{1}{2}$ 1 1	3

Qn. No	Sub Qns	Answer Key/Value Points	Score	Total Score																																		
23.	(a)	Population mean = 12.2	1	3																																		
	(b)	<table border="1" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th>S/No</th> <th>Sample</th> <th>Sample mean</th> </tr> </thead> <tbody> <tr><td>1</td><td>8, 9</td><td>8.5</td></tr> <tr><td>2</td><td>8, 13</td><td>10.5</td></tr> <tr><td>3</td><td>8, 15</td><td>11.5</td></tr> <tr><td>4</td><td>8, 16</td><td>12</td></tr> <tr><td>5</td><td>9, 13</td><td>11</td></tr> <tr><td>6</td><td>9, 15</td><td>12</td></tr> <tr><td>7</td><td>9, 16</td><td>12.5</td></tr> <tr><td>8</td><td>13, 15</td><td>14</td></tr> <tr><td>9</td><td>13, 16</td><td>14.5</td></tr> <tr><td>10</td><td>15, 16</td><td>15.5</td></tr> <tr> <td colspan="2" style="text-align: center;">Total</td> <td>122</td> </tr> </tbody> </table> <p style="text-align: center;"> $E(\text{Sample mean}) = \frac{122}{10} = 12.2$ $= \text{Population mean}$ </p>	S/No		Sample	Sample mean	1	8, 9	8.5	2	8, 13	10.5	3	8, 15	11.5	4	8, 16	12	5	9, 13	11	6	9, 15	12	7	9, 16	12.5	8	13, 15	14	9	13, 16	14.5	10	15, 16	15.5	Total	
S/No	Sample	Sample mean																																				
1	8, 9	8.5																																				
2	8, 13	10.5																																				
3	8, 15	11.5																																				
4	8, 16	12																																				
5	9, 13	11																																				
6	9, 15	12																																				
7	9, 16	12.5																																				
8	13, 15	14																																				
9	13, 16	14.5																																				
10	15, 16	15.5																																				
Total		122																																				
24		<table border="1" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th>Year</th> <th>No. of books</th> <th>3 yearly moving Total</th> <th>3 yearly moving Average</th> </tr> </thead> <tbody> <tr><td>2015</td><td>15</td><td>—</td><td>—</td></tr> <tr><td>2016</td><td>28</td><td>73</td><td>24.33</td></tr> <tr><td>2017</td><td>30</td><td>86</td><td>28.67</td></tr> <tr><td>2018</td><td>28</td><td>84</td><td>28</td></tr> <tr><td>2019</td><td>26</td><td>86</td><td>28.67</td></tr> <tr><td>2020</td><td>32</td><td>—</td><td>—</td></tr> </tbody> </table> <p style="text-align: center;">(3 yearly moving total only give 2 scores)</p>	Year	No. of books	3 yearly moving Total	3 yearly moving Average	2015	15	—	—	2016	28	73	24.33	2017	30	86	28.67	2018	28	84	28	2019	26	86	28.67	2020	32	—	—	3	3						
Year	No. of books	3 yearly moving Total	3 yearly moving Average																																			
2015	15	—	—																																			
2016	28	73	24.33																																			
2017	30	86	28.67																																			
2018	28	84	28																																			
2019	26	86	28.67																																			
2020	32	—	—																																			

Qn. No	Sub Qns	Answer Key/Value Points	Score	Total Score																												
25.		<table border="1" style="margin-bottom: 10px;"> <thead> <tr> <th>R_1</th> <th>R_2</th> <th>$d=R_1-R_2$</th> <th>d^2</th> </tr> </thead> <tbody> <tr> <td>4</td> <td>3</td> <td>1</td> <td>1</td> </tr> <tr> <td>2.5</td> <td>4</td> <td>-1.5</td> <td>2.25</td> </tr> <tr> <td>5</td> <td>1</td> <td>4</td> <td>16</td> </tr> <tr> <td>1</td> <td>5</td> <td>-4</td> <td>16</td> </tr> <tr> <td>2.5</td> <td>2</td> <td>0.5</td> <td>0.25</td> </tr> <tr> <td></td> <td></td> <td></td> <td>35.5</td> </tr> </tbody> </table> <p style="margin-left: 20px;">[Tied ranks need not be considered]</p> $CF = \sum \frac{m^3 - m}{12} = \frac{2^3 - 2}{12} = 0.5$ $F = 1 - \frac{6(\sum d^2 + CF)}{n^3 - n}$ $= 1 - \frac{6(35.5 + 0.5)}{5^3 - 5} = -0.8$	R_1	R_2	$d=R_1-R_2$	d^2	4	3	1	1	2.5	4	-1.5	2.25	5	1	4	16	1	5	-4	16	2.5	2	0.5	0.25				35.5	<p style="text-align: center;">$1\frac{1}{2}$</p> <p style="text-align: center;">$\frac{1}{2}$</p> <p style="text-align: center;">$\frac{1}{2}$</p>	<p style="text-align: center;">3</p>
R_1	R_2	$d=R_1-R_2$	d^2																													
4	3	1	1																													
2.5	4	-1.5	2.25																													
5	1	4	16																													
1	5	-4	16																													
2.5	2	0.5	0.25																													
			35.5																													
26.		$F(x) = P(X \leq x) = \int_{-\infty}^x f(x) dx$ $= \begin{cases} 0, & x < 0 \\ x^2, & 0 < x < 1 \\ 1, & x \geq 1 \end{cases}$	<p style="text-align: center;">2</p> <p style="text-align: center;">1</p>	<p style="text-align: center;">3</p>																												
27.		$R'(x) = 22 - 2x$ $R''(x) = -2$ $R'(x) = 0 \Rightarrow x = 11$ <p style="margin-left: 20px;">Maximum value at $x=11 = 121$</p>	<p style="text-align: center;">1</p> <p style="text-align: center;">$\frac{1}{2}$</p> <p style="text-align: center;">1</p> <p style="text-align: center;">$\frac{1}{2}$</p>	<p style="text-align: center;">3</p>																												

Qn. No	Sub Qns	Answer Key/Value Points	Score	Total Score
28.		$b_{xy} = 1.09$ [Using any relevant formula) $x - \bar{x} = b_{xy} (y - \bar{y})$ $x - 6.14 = 1.09 (y - 4)$	$2\frac{1}{2}$ 1 $\frac{1}{2}$	4.
29.	(a)	$P(1 \leq X \leq 2) = \frac{1}{6} + \frac{2}{6} = \frac{3}{6} = 0.5$	1	4.
	(b)	$E(X) = \sum x P(x)$ $= 1 \times \frac{1}{6} + 2 \times \frac{2}{6} + 3 \times \frac{3}{6} = \frac{14}{6} = 2.33$	$\frac{1}{2}$ $\frac{1}{2}$	
	(c)	$E(X^2) = \sum x^2 P(x) = 6$ $V(X) = E(X^2) - (E(X))^2$ $= 6 - (2.33)^2 = 0.5711$ [If $E(X) = \frac{14}{6}$, $V(X) = 6 - \left(\frac{14}{6}\right)^2 = 0.55$]	1 $\frac{1}{2}$ $\frac{1}{2}$	
30.	(a)	4 conditions $(4 \times \frac{1}{2} = 2)$	2	4.
	(b)	$n_p = 6$, $n_{pq} = 3.6$ $q = \frac{n_{pq}}{n_p} = \frac{3.6}{6} = 0.6$ $p = 1 - q = 1 - 0.6 = 0.4$	$\frac{1}{2} + \frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$	
31	(a)	Assignable factors - Definition / Examples / Explanation	1	

Qn. No	Sub Qns	Answer Key/Value Points	Score	Total Score
		$= \frac{-10}{\sqrt{254} \times \sqrt{100}}$ $= -0.063$ <p>(Cov(x,y) = 0.4, $\sigma_x = 3.19$, $\sigma_y = 2$, $r = 0.063$)</p>	1 1/2	6

35	(a)	<table border="1"> <thead> <tr> <th>P₀</th> <th>P₁</th> <th>I = $\frac{P_1}{P_0} \times 100$</th> </tr> </thead> <tbody> <tr> <td>44</td> <td>48</td> <td>109.1</td> </tr> <tr> <td>38</td> <td>40</td> <td>105.3</td> </tr> <tr> <td>48</td> <td>54</td> <td>112.5</td> </tr> <tr> <td></td> <td></td> <td>326.9</td> </tr> </tbody> </table> <p>Simple AM Index No = $\frac{\sum I}{n}$</p> $= \frac{326.9}{3} = 108.97$	P ₀	P ₁	I = $\frac{P_1}{P_0} \times 100$	44	48	109.1	38	40	105.3	48	54	112.5			326.9	1 1/2 1/2	
P ₀	P ₁	I = $\frac{P_1}{P_0} \times 100$																	
44	48	109.1																	
38	40	105.3																	
48	54	112.5																	
		326.9																	

	(b)	<table border="1"> <thead> <tr> <th>P₀</th> <th>Q₀</th> <th>P₁</th> <th>Q₁</th> <th>P₀Q₀</th> <th>P₁Q₀</th> </tr> </thead> <tbody> <tr> <td>2</td> <td>20</td> <td>5</td> <td>15</td> <td>40</td> <td>100</td> </tr> <tr> <td>4</td> <td>4</td> <td>8</td> <td>5</td> <td>16</td> <td>32</td> </tr> <tr> <td>1</td> <td>10</td> <td>2</td> <td>12</td> <td>10</td> <td>20</td> </tr> <tr> <td>5</td> <td>5</td> <td>10</td> <td>6</td> <td>25</td> <td>50</td> </tr> <tr> <td></td> <td></td> <td></td> <td></td> <td>91</td> <td>202</td> </tr> </tbody> </table> <p>Laspeyres's IN = $\frac{\sum P_1 Q_0}{\sum P_0 Q_0} \times 100$</p> $= \frac{202}{91} \times 100 = 221.98$	P ₀	Q ₀	P ₁	Q ₁	P ₀ Q ₀	P ₁ Q ₀	2	20	5	15	40	100	4	4	8	5	16	32	1	10	2	12	10	20	5	5	10	6	25	50					91	202	2 1 1	6
P ₀	Q ₀	P ₁	Q ₁	P ₀ Q ₀	P ₁ Q ₀																																			
2	20	5	15	40	100																																			
4	4	8	5	16	32																																			
1	10	2	12	10	20																																			
5	5	10	6	25	50																																			
				91	202																																			

Qn. No	Sub Qns	Answer Key/Value Points	Score	Total Score
36.	(a)	$V(T_1) = V(2x_1 - 2x_2 + x_3)$ $= 4\sigma^2 + 4\sigma^2 + \sigma^2 = 9\sigma^2$ $V(T_2) = V(3x_1 - x_2 - x_3)$ $= 9\sigma^2 + \sigma^2 + \sigma^2 = 11\sigma^2$ $V(T_1) < V(T_2)$ $\therefore T_1 \text{ is more efficient than } T_2$	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$	
	(b)	$H_0: \mu = 50, \quad H_1: \mu \neq 50$ $n = 144, \quad s = 20, \quad \bar{x} = 55, \quad \alpha = 0.05$ $Z = \frac{\bar{x} - \mu_0}{\frac{s}{\sqrt{n}}}$ $= \frac{55 - 50}{\frac{20}{\sqrt{144}}} = 3$ $ Z = 3 > Z_{\alpha/2} \quad \text{Reject } H_0$	$\frac{1}{2}$ $\frac{1}{2}$ 1 $\frac{1}{2}$ $\frac{1}{2}$	6

- 1. Dr. MANOJ.K ~~28/04/2022~~
- 2. Dr. Sajish Kumar.M, m.v.k.m.h.s.s. Chittilanchery Sajish
- 3. Dr. Beju. G.V , Govt VHSS Vithura, Beju
- 4. Dasana Kumary.D, NISS Chelvanood, Kollam Dasana
- 5. Smitha.M.S. SNHSS, Pocheekal Smitha
- 6. LATHA. S.G, SMU NSS HSS Kallal Kottayam Lat
- 7. Seby Jose.P m.s.m.h.s.s. kallinga/paramba Seby